

Exempel

Beräkna riktningsderivatan av

$$f(x, y, z) = xy + xe^z$$

i punkten $(2, 1, 0)$ i riktning $(1, 2, 4)$.

$$f(x, y, z) = xy + xe^z$$

$$f(x, y, z) = xy + xe^z$$

OBS! Normera riktningen:

$$\bar{v} = \frac{(1, 2, 4)}{|(1, 2, 4)|} = \frac{1}{\sqrt{21}}(1, 2, 4).$$

$$f(x, y, z) = xy + xe^z$$

OBS! Normera riktningen:

$$\bar{v} = \frac{(1, 2, 4)}{|(1, 2, 4)|} = \frac{1}{\sqrt{21}}(1, 2, 4).$$

$$f'_{\bar{v}} = \nabla f \bullet \bar{v},$$

$$f(x, y, z) = xy + xe^z$$

OBS! Normera riktningen:

$$\bar{v} = \frac{(1, 2, 4)}{|(1, 2, 4)|} = \frac{1}{\sqrt{21}}(1, 2, 4).$$

$$f'_{\bar{v}} = \nabla f \bullet \bar{v},$$

$$\nabla f = (f'_x, f'_y, f'_z) = (y + e^z, x, xe^z),$$

$$f(x, y, z) = xy + xe^z$$

OBS! Normera riktningen:

$$\bar{v} = \frac{(1, 2, 4)}{|(1, 2, 4)|} = \frac{1}{\sqrt{21}}(1, 2, 4).$$

$$f'_{\bar{v}} = \nabla f \bullet \bar{v},$$

$$\nabla f = (f'_x, f'_y, f'_z) = (y + e^z, x, xe^z),$$

$$\nabla f(2, 1, 0) = (2, 2, 2).$$

$$f(x, y, z) = xy + xe^z$$

OBS! Normera riktningen:

$$\bar{v} = \frac{(1, 2, 4)}{|(1, 2, 4)|} = \frac{1}{\sqrt{21}}(1, 2, 4).$$

$$f'_{\bar{v}} = \nabla f \bullet \bar{v},$$

$$\nabla f = (f'_x, f'_y, f'_z) = (y + e^z, x, xe^z),$$

$$\nabla f(2, 1, 0) = (2, 2, 2).$$

$$f'_{\bar{v}}(2, 1, 0) = (2, 2, 2) \bullet \frac{1}{\sqrt{21}}(1, 2, 4)$$

$$f(x, y, z) = xy + xe^z$$

OBS! Normera riktningen:

$$\bar{v} = \frac{(1, 2, 4)}{|(1, 2, 4)|} = \frac{1}{\sqrt{21}}(1, 2, 4).$$

$$f'_{\bar{v}} = \nabla f \bullet \bar{v},$$

$$\nabla f = (f'_x, f'_y, f'_z) = (y + e^z, x, xe^z),$$

$$\nabla f(2, 1, 0) = (2, 2, 2).$$

$$f'_{\bar{v}}(2, 1, 0) = (2, 2, 2) \bullet \frac{1}{\sqrt{21}}(1, 2, 4) = \frac{2+4+8}{\sqrt{21}} = \frac{2\sqrt{7}}{\sqrt{3}}.$$