

Examiner: Xiangfeng Yang (Tel: 013 28 57 88).

Things allowed (Hjälpmedel): a calculator.

Scores rating (Betygsgränser): 8-11 points giving rate 3; 11.5-14.5 points giving rate 4; 15-18 points giving rate 5.

1 (3 points)

In a certain residential suburb, 60% of all households get Internet service from the local cable company, 80% get TV service from that company, and 50% get both services from that company. Now a household is randomly selected.

(1.1) (1p) What is the probability that the household gets at least one of these two services from that company?

(1.2) (1p) What is the probability that the household gets exactly one of these two services from that company?

(1.3) (1p) Given that the household gets Internet service from that company, what is the probability that the household also gets TV service from that company?

2 (3 points)

SJ Express train (Snabbtåg) #520 is scheduled to run 200 minutes for one trip from Linköping central station to Stockholm central station. However, because of many factors, the actual running times of one trip might be different from 200 minutes. Let X be the distribution of actual running times of one trip and assume that $X \sim N(200, 4^2)$.

(2.1) (1p) Find the probability that one trip takes less than 190 minutes, that is, $P(X < 190)$.

(2.2) (1p) Find the probability that one trip takes more than 208 minutes, that is, $P(X > 208)$.

(2.3) (1p) Find the probability $P(195 < X < 205)$.

3 (3 points)

A coin has two sides: head (denoted as 1) and tail (denoted as 0). Let X be the upper side in one flip of the coin, and assume that the coin is not fair with the following probabilities:

$$P(X = 1) = 0.6, \quad P(X = 0) = 0.4.$$

(3.1) (1p) Find the mean $\mu = E(X)$ and the variance $\sigma^2 = V(X)$ of X .

(3.2) (2p) The coin is flipped 100 times, find the probability that one gets at most 65 heads.

4 (3 points)

Let a population X be continuous with a pdf

$$f(x) = \theta \cdot x^{\theta-1}, \quad \text{for } 0 < x < 1,$$

where $\theta > 0$ is an unknown parameter. A sample $\{0.24, 0.66, 0.53, 0.14, 0.82\}$ is taken from this population.

(4.1) (1p) Use the method of moments to find a point estimate $\hat{\theta}_{MM}$ of θ .

(4.2) (2p) Use the maximum-likelihood method to find a point estimate $\hat{\theta}_{ML}$ of θ .

5 (3 points)

Assume that a population is $X \sim N(\mu, \sigma^2)$. A sample from this population has: $n = 16, \bar{x} = 3.1$ and $s = 1.2$.

(5.1) (2p) Assume that $\sigma^2 = 1$. Does the sample provide any evidence that $\mu > 2.8$? Answer this by constructing an appropriate one-sided 95% confidence interval of μ .

(5.2) (1p) Assume that σ^2 is unknown. With a significance level 5%, test the hypotheses

$$H_0 : \sigma^2 = 1 \quad \text{against} \quad H_a : \sigma^2 > 1.$$

6 (3 points)

To understand factors affecting snowfall size in Linköping in each December, researchers study snowfall size Y in terms of temperature x_1 and moisture content x_2 , by linear regression:

$$Y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \varepsilon, \quad \text{where } \varepsilon \sim N(0, \sigma^2).$$

To study such linear regression, researchers use a sample with sample size $n = 16$ and the method of least squares to obtain: the estimated values $\hat{\beta}_i$ of β_i , together with their standard errors denoted as $se(\hat{\beta}_i)$ or $s_{\hat{\beta}_i}$:

$$\hat{\beta}_0 = 1.58, \quad se(\hat{\beta}_0) = 0.14; \quad \hat{\beta}_1 = 8.12, \quad se(\hat{\beta}_1) = 0.28; \quad \hat{\beta}_2 = 0.08, \quad se(\hat{\beta}_2) = 0.02.$$

Researchers want to examine whether or not snowfall size Y really depends on moisture content x_2 . To this end, one wants to check whether or not $\beta_2 \neq 0$, based on the sample.

(6.1) (2p) Is $\beta_2 \neq 0$? Answer this by constructing a 95% confidence interval of β_2 .

(6.2) (1p) With a significance level 5%, perform the hypothesis test

$$H_0 : \beta_2 = 0 \quad \text{against} \quad H_a : \beta_2 \neq 0$$

1. Basic probability

(1.1) Conditional probability $P(A|B) = \frac{P(A \cap B)}{P(B)}$.

(1.2) Total probability $P(B) = \sum_{i=1}^k P(B|A_i)P(A_i)$ where $\{A_i\}$ are disjoint and $\cup_{i=1}^k A_i = S$.

(1.3) Bayes' Theorem $P(A_i|B) = \frac{P(B|A_i)P(A_i)}{\sum_{k=1}^n P(B|A_k)P(A_k)}$ where $\{A_i\}$ are in (1.2).

2. Random variables (r.v.s)

(2.1) Discrete r.v. X has a pmf $p(x) = P(X = x)$ satisfying $p(x) \geq 0$ and $\sum p(x_i) = 1$,

$$\begin{array}{c|cccc} X & x_1 & x_2 & \cdots & x_n & \cdots \\ \hline p(x) & p(x_1) & p(x_2) & \cdots & p(x_n) & \cdots \end{array}$$

Expectation (or *Expected value* or *mean*) $\mu_X = E(X) = \sum x_i p(x_i)$;

Variance $\sigma_X^2 = V(X) = E(X - \mu_X)^2 = E(X^2) - \mu_X^2 = \sum x_i^2 p(x_i) - (\sum x_i p(x_i))^2$.

(2.2) Continuous r.v. X has a pdf $f(x)$ satisfying $f(x) \geq 0$ and $\int_{-\infty}^{\infty} f(x) dx = 1$,

$$P(a < X < b) = \int_a^b f(x) dx.$$

Expectation (or *Expected value* or *mean*) $\mu_X = E(X) = \int_{-\infty}^{\infty} x f(x) dx$;

Variance $\sigma_X^2 = V(X) = E(X - \mu_X)^2 = E(X^2) - \mu_X^2 = \int_{-\infty}^{\infty} x^2 f(x) dx - (\int_{-\infty}^{\infty} x f(x) dx)^2$.

(2.3) Cumulative distribution function (cdf) of a r.v. X is $F(x) = P(X \leq x)$.

(2.4) X and Y are r.v.s, a, b and c are scalars, then

$$E(aX + bY + c) = aE(X) + bE(Y) + c,$$

$$V(aX + bY + c) = a^2V(X) + b^2V(Y) + 2ab \operatorname{cov}(X, Y),$$

$$E(g(X, Y)) = \begin{cases} \sum_{i,j} g(x_i, y_j) \cdot p(x_i, y_j), & \text{for discrete } (X, Y), \\ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) \cdot f(x, y) dx dy, & \text{for continuous } (X, Y). \end{cases}$$

(2.5) • Discrete r.v. (X, Y) has a joint pmf $p(x, y)$ satisfying $p(x, y) \geq 0$ and $\sum_{x_i} \sum_{y_j} p(x_i, y_j) = 1$.

The *marginal pmf* of X is $p_X(x) = \sum_y p(x, y)$;

The *marginal pmf* of Y is $p_Y(y) = \sum_x p(x, y)$;

X and Y are *independent* if $p(x, y) = p_X(x) \cdot p_Y(y)$.

• Continuous r.v. (X, Y) has a joint pdf $p(x, y)$ satisfying $f(x, y) \geq 0$ and $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) dx dy = 1$.

The *marginal pdf* of X is $f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy$;

The *marginal pdf* of Y is $f_Y(y) = \int_{-\infty}^{\infty} f(x, y) dx$;

X and Y are *independent* if $f(x, y) = f_X(x) \cdot f_Y(y)$.

3. Several special r.v.s

(3.1) $X \sim \operatorname{Bin}(n, p)$ has a pmf $p(x) = P(X = x) = \binom{n}{x} \cdot p^x \cdot (1-p)^{n-x}$, $x = 0, 1, 2, \dots, n$.

$$E(X) = n \cdot p, \quad V(X) = n \cdot p \cdot (1-p).$$

(3.2) $X \sim \operatorname{Po}(\lambda)$ has a pmf $p(x) = P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$, $x = 0, 1, 2, \dots$
 $E(X) = \lambda, \quad V(X) = \lambda$.

(3.3) $X \sim \operatorname{Hypergeometric}$ has a pmf $p(x) = P(X = x) = \frac{\binom{M}{x} \binom{N-M}{n-x}}{\binom{N}{n}}$.

(3.4) $X \sim \operatorname{Exp}(\lambda)$ has a pdf

$$f(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0, \\ 0, & \text{otherwise.} \end{cases}$$

(3.5) $E(X) = \frac{1}{\lambda}, \quad V(X) = (\frac{1}{\lambda})^2$.

$X \sim N(\mu, \sigma^2)$ has a pdf

$$f(x) = \frac{1}{\sigma \sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad -\infty < x < \infty.$$

(3.6) $E(X) = \mu, \quad V(X) = \sigma^2$.

$X \sim U(a, b)$ has a pdf

$$f(x) = \begin{cases} \frac{1}{b-a}, & a < x < b, \\ 0, & \text{otherwise.} \end{cases}$$

$$E(X) = \frac{a+b}{2}, \quad V(X) = \frac{(b-a)^2}{12}.$$

4. Central Limit Theorem (CLT)

Suppose that a population has mean $= \mu$ and variance $= \sigma^2$. A random sample $\{X_1, X_2, \dots, X_n\}$ from this population is given. Then for large $n \geq 30$,

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1). \tag{1}$$

• If the population is normal, then (1) holds for any n .

• Note that $\mu = E(\bar{X})$ and $(\sigma/\sqrt{n})^2 = V(\bar{X})$.

5. Several notations in statistics

(5.1) Sample mean: $\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n} = \sum \bar{X}_i$; $\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \sum x_i$.

(5.2) Sample variance:

$$S^2 = \frac{\sum (X_i - \bar{X})^2}{n-1} = \frac{1}{n-1} \left(\sum X_i^2 - \frac{(\sum X_i)^2}{n} \right); \quad s^2 = \frac{\sum (x_i - \bar{x})^2}{n-1} = \frac{1}{n-1} \left(\sum x_i^2 - \frac{(\sum x_i)^2}{n} \right).$$

• Capital letters \bar{X} and S^2 refer to the objects based on random sample (therefore they are in general r.v.s), while small letters \bar{x} and s^2 are the objects based on observations (so they are scalars).

(5.3) A point estimator of θ obtained by Methods of Moments is denoted as $\hat{\theta}_{MM}$.

(5.4) A point estimator of θ obtained by Maximum Likelihood method is denoted as $\hat{\theta}_{ML}$.

6. Confidence Interval (CI)

In this course, three types of confidence intervals are studied depending on the unknown population parameter(s): CI-1 (confidence intervals for population mean(s)), CI-2 (confidence intervals for population variance(s)), and CI-3 (confidence intervals for population proportion(s)).

CI-1: (1 - α) CI of a population mean μ

case 1.1 (any n) If population $X \sim N(\mu, \sigma^2)$ and σ^2 is known, then $\frac{\bar{X}-\mu}{\sigma/\sqrt{n}} \sim N(0, 1)$ and

$$I_\mu = (\bar{x} - z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}, \bar{x} + z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}) := \bar{x} \mp z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}.$$

case 1.2 (n ≥ 30) For any population X , it holds that $\frac{\bar{X}-\mu}{\sigma/\sqrt{n}} \sim N(0, 1)$ and

$$I_\mu = \bar{x} \mp z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \text{ or } I_\mu = \bar{x} \mp z_{\alpha/2} \cdot \frac{\hat{\sigma}}{\sqrt{n}}.$$

case 1.3 (any n) If population $X \sim N(\mu, \sigma^2)$ and σ^2 is unknown, then $\frac{\bar{X}-\mu}{S/\sqrt{n}} \sim T(n-1)$ and

$$I_\mu = \bar{x} \mp t_{\alpha/2}(n-1) \cdot \frac{s}{\sqrt{n}}.$$

CI-1': (1 - α) CI of the difference of two population means $\mu_X - \mu_Y$

case 1.1' (any n_1, n_2) If independent populations $X \sim N(\mu_X, \sigma_X^2)$, $Y \sim N(\mu_Y, \sigma_Y^2)$, and σ_X^2, σ_Y^2 are known, then

$$\frac{(\bar{X}-\bar{Y})-(\mu_X-\mu_Y)}{\sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}} \sim N(0, 1), \text{ and } I_{\mu_X-\mu_Y} = (\bar{x}-\bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}.$$

case 1.2' ($n_1, n_2 \geq 30$) For any independent populations X and Y , it holds that



$$\frac{(\bar{X}-\bar{Y})-(\mu_X-\mu_Y)}{\sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}} \sim N(0, 1) \text{ and}$$

$$I_{\mu_X-\mu_Y} = (\bar{x}-\bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}} \text{ or } I_{\mu_X-\mu_Y} = (\bar{x}-\bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{\sigma}_X^2}{n_1} + \frac{\hat{\sigma}_Y^2}{n_2}}.$$

case 1.3' (any n_1, n_2) If independent populations $X \sim N(\mu_X, \sigma_X^2)$, $Y \sim N(\mu_Y, \sigma_Y^2)$, where σ_X^2, σ_Y^2 are unknown but $\sigma_X^2 = \sigma_Y^2$, then

$$\frac{(\bar{X}-\bar{Y})-(\mu_X-\mu_Y)}{S\sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} \sim T(n_1+n_2-2), \text{ where } S^2 = \frac{(n_1-1)S_X^2 + (n_2-1)S_Y^2}{n_1+n_2-2}, \text{ and}$$

$$I_{\mu_X-\mu_Y} = (\bar{x}-\bar{y}) \mp t_{\alpha/2}(n_1+n_2-2) \cdot s\sqrt{\frac{1}{n_1} + \frac{1}{n_2}}.$$



CI-2: (1 - α) CI of population variance(s) σ^2


• If a population $X \sim N(\mu, \sigma^2)$ and σ^2 is unknown, then $\frac{(n-1)S^2}{\sigma^2} \sim \chi^2(n-1)$, and

$$I_{\sigma^2} = \left(\frac{(n-1)s^2}{\chi_{\alpha/2}^2(n-1)}, \frac{(n-1)s^2}{\chi_{1-\alpha/2}^2(n-1)} \right).$$

• If two independent populations $X \sim N(\mu_X, \sigma^2)$ and $Y \sim N(\mu_Y, \sigma^2)$, and σ^2 is unknown, then $\frac{(n_1+n_2-2)S^2}{\sigma^2} \sim \chi^2(n_1+n_2-2)$, and

$$I_{\sigma^2} = \left(\frac{(n_1+n_2-2)s^2}{\chi_{\alpha/2}^2(n_1+n_2-2)}, \frac{(n_1+n_2-2)s^2}{\chi_{1-\alpha/2}^2(n_1+n_2-2)} \right),$$

where $S^2 = \frac{(n_1-1)S_X^2 + (n_2-1)S_Y^2}{n_1+n_2-2}$.



CI-3: (1 - α) CI of population proportion(s)

• If a (large) population has an unknown proportion p , then $\frac{\hat{p}-p}{\sqrt{p(1-p)/n}} \sim N(0, 1)$ if $n\hat{p} \geq 10$ and $n(1-\hat{p}) \geq 10$ with $\hat{p} = x/n$, and $I_p = \hat{p} \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$.

• If two independent (large) populations have unknown proportions p_1 and p_2 , then

$$\frac{(\hat{p}_1-\hat{p}_2)-(p_1-p_2)}{\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{\hat{p}_2(1-\hat{p}_2)}{n_2}}} \sim N(0, 1)$$

if $n_i\hat{p}_i \geq 10$ and $n_i(1-\hat{p}_i) \geq 10$ for $i = 1, 2$, and $I_{p_1-p_2} = (\hat{p}_1-\hat{p}_2) \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{\hat{p}_2(1-\hat{p}_2)}{n_2}}$.

7. Hypothesis Test (HT)

	H_0 is true	H_0 is false and $\theta = \theta_1$
reject H_0	(type I error or significance level) α	(power) $h(\theta_1)$
don't reject H_0	$1 - \alpha$	(type II error) $\beta(\theta_1) = 1 - h(\theta_1)$

reject $H_0 \Leftrightarrow TS \in C \Leftrightarrow p\text{-value} < \alpha$

χ^2 tests for populations (non-parametric tests)

Suppose that for a random sample of a population X , the n elements of it are classified into k disjoint groups $A_i, 1 \leq i \leq k$. For each group $A_i, 1 \leq i \leq k$, suppose that there are $N_i, 1 \leq i \leq k$ elements inside. Let $p_i = P(A_i)$ assuming a given distribution of X . Note that $p_1 + p_2 + \dots + p_k = 1$ and $N_1 + N_2 + \dots + N_k = n$. One wants to test the hypotheses

$$H_0 : P(A_i) = p_i, \quad 1 \leq i \leq k, \quad H_a : P(A_i) \neq p_i \text{ for some } 1 \leq i \leq k.$$

If n is large in the sense that $np_i \geq 5$ for all $1 \leq i \leq k$, then the test statistic is

$$\sum_{i=1}^k \frac{(N_i - np_i)^2}{np_i} \approx \chi^2(k-1).$$

Therefore the observation of the test statistic is

$$TS = \sum_{i=1}^k \frac{(n_i - np_i)^2}{np_i}, \text{ where } n_i \text{ is the observation of } N_i, 1 \leq i \leq k.$$

For the critical region C , one can take (note that if H_0 is true, then TS should be close to zero)

$$C = (\chi_{\alpha}^2(k-1), \infty).$$

The conclusion would be $TS \in C \iff H_0$ is rejected.

8. Linear and logistic regression

(Multiple) linear regression: $Y = \beta_0 + \beta_1 x_1 + \dots + \beta_k x_k + \varepsilon$, $\varepsilon \sim N(0, \sigma^2)$.

- Y : response variable (which is normal r.v.), $\{x_1, \dots, x_k\}$: predictors (which are scalars).
- sample: $\{(x_{11}, \dots, x_{1k}; y_1), (x_{21}, \dots, x_{2k}; y_2), \dots, (x_{n1}, \dots, x_{nk}; y_n)\}$.
- how to estimate $\beta_j \approx \hat{\beta}_j$: least square method, that is, to minimize $\sum_{i=1}^n (y_i - \hat{y}_i)^2$, where the estimated (multiple) linear regression line \hat{y} is

$$\hat{y} = \hat{\beta}_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k.$$

- $\frac{\hat{\beta}_j - \beta_j}{se(\hat{\beta}_j)} \sim T(n-k-1)$, this helps determine whether or not the real $\beta_j = 0$?
- $\sigma^2 \approx \frac{SSE}{n-k-1}$, this gives an estimation of the size of the error.
- $R^2 = \frac{SSR}{SSY}$, this gives how well the model is (if $R^2 \approx 1$, then the model fits the sample very well).
- How to test $\beta_1 = \dots = \beta_k = 0$? Use the random variable $\frac{SSR/k}{SSE/(n-k-1)} \sim F(k, n-k-1)$.

Logistic regression: Let Y can only take 0 or 1 with $P(Y=1) = p$ and $P(Y=0) = 1-p$.

$$E(Y) = p(x_1, \dots, x_k) = \frac{e^{\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k}}{1 + e^{\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k}}.$$

- Y : response variable (which is Bernoulli r.v. $P(Y=1) = p$ and $P(Y=0) = 1-p$, so $E(Y) = p$), $\{x_1, \dots, x_k\}$: predictors (which are scalars).
- sample: $\{(x_{11}, \dots, x_{1k}; y_1), (x_{21}, \dots, x_{2k}; y_2), \dots, (x_{n1}, \dots, x_{nk}; y_n)\}$.
- how to estimate $\beta_j \approx \hat{\beta}_j$: maximal likelihood method (maximize $\prod_{i=1}^n p(x_{i1}, \dots, x_{ik})^{y_i} (1 - p(x_{i1}, \dots, x_{ik}))^{1-y_i}$).
- $\frac{\hat{\beta}_j - \beta_j}{se(\hat{\beta}_j)} \approx N(0, 1)$ for large $n \geq 30$, this helps determine whether or not the real $\beta_j = 0$?
- Classification of a new object $Y(x_1, \dots, x_k)$ as 1 or 0 according

$$Y(x_1, \dots, x_k) = \begin{cases} 1, & \text{if } \hat{p}(x_1, \dots, x_k) \geq 0.5, \\ 0, & \text{if } \hat{p}(x_1, \dots, x_k) < 0.5, \end{cases}$$

where the estimated logit function $\hat{p}(x_1, \dots, x_k)$ is

$$\hat{p}(x_1, \dots, x_k) = \frac{e^{\hat{\beta}_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k}}{1 + e^{\hat{\beta}_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k}}.$$

9. Tables

(9.1) Table for $N(0, 1)$ standard normal random variable $\Phi(x) = P(N(0, 1) \leq x)$, $x \geq 0$.
There is an important relation $\Phi(-x) = 1 - \Phi(x)$, $x \geq 0$.

x	0	1	2	3	4	5	6	7	8	9
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9564	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9992	0.9993	0.9993	0.9993
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997	0.9997
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998	0.9998	0.9998
3.5	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998
3.6	0.9998	0.9998	0.9998	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.7	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.8	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.9	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
4.0	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

(9.2) Table for $T(f)$ random variable $F(x) = P(T(f) \leq x)$,
where f is a parameter called 'degrees of freedom'.

f	0.75	0.90	0.95	0.975	0.99	0.995	0.9975	0.9995
1	1.00	3.08	6.31	12.71	31.82	63.66	127.32	636.62
2	0.82	1.89	2.92	4.30	6.96	9.92	14.09	31.60
3	0.76	1.64	2.35	3.18	4.54	5.84	7.45	12.92
4	0.74	1.53	2.13	2.78	3.75	4.60	5.60	8.61
5	0.73	1.48	2.02	2.57	3.36	4.03	4.77	6.87
6	0.72	1.44	1.94	2.45	3.14	3.71	4.32	5.96
7	0.71	1.41	1.89	2.36	3.00	3.50	4.03	5.41
8	0.71	1.40	1.86	2.31	2.90	3.36	3.83	5.04
9	0.70	1.38	1.83	2.26	2.82	3.25	3.69	4.78
10	0.70	1.37	1.81	2.23	2.76	3.17	3.58	4.59
11	0.70	1.36	1.80	2.20	2.72	3.11	3.50	4.44
12	0.70	1.36	1.78	2.18	2.68	3.05	3.43	4.32
13	0.69	1.35	1.77	2.16	2.65	3.01	3.37	4.22
14	0.69	1.35	1.76	2.14	2.62	2.98	3.33	4.14
15	0.69	1.34	1.75	2.13	2.60	2.95	3.29	4.07
16	0.69	1.34	1.75	2.12	2.58	2.92	3.25	4.01
17	0.69	1.33	1.74	2.11	2.57	2.90	3.22	3.97
18	0.69	1.33	1.73	2.10	2.55	2.88	3.20	3.92
19	0.69	1.33	1.73	2.09	2.54	2.86	3.17	3.88
20	0.69	1.33	1.72	2.09	2.53	2.85	3.15	3.85
21	0.69	1.32	1.72	2.08	2.52	2.83	3.14	3.82
22	0.69	1.32	1.72	2.07	2.51	2.82	3.12	3.79
23	0.69	1.32	1.71	2.07	2.50	2.81	3.10	3.77
24	0.68	1.32	1.71	2.06	2.49	2.80	3.09	3.75
25	0.68	1.32	1.71	2.06	2.49	2.79	3.08	3.73
26	0.68	1.31	1.71	2.06	2.48	2.78	3.07	3.71
27	0.68	1.31	1.70	2.05	2.47	2.77	3.06	3.69
28	0.68	1.31	1.70	2.05	2.47	2.76	3.05	3.67
29	0.68	1.31	1.70	2.05	2.46	2.76	3.04	3.66
30	0.68	1.31	1.70	2.04	2.46	2.75	3.03	3.65
40	0.68	1.30	1.68	2.02	2.42	2.70	2.97	3.55
50	0.68	1.30	1.68	2.01	2.40	2.68	2.94	3.50
60	0.68	1.30	1.67	2.00	2.39	2.66	2.91	3.46
100	0.68	1.29	1.66	1.98	2.36	2.63	2.87	3.39
∞	0.67	1.28	1.65	1.96	2.33	2.58	2.81	3.29

(9.3) Table for $\chi^2(f)$ random variable $F(x) = P(\chi^2(f) \leq x)$, where f is a parameter.

f	0.0005	0.001	0.005	0.01	0.025	0.05	0.10	0.20	0.30	0.40	0.50
1	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.06	0.15	0.27	0.45
2	0.00	0.00	0.01	0.02	0.05	0.10	0.21	0.45	0.71	1.02	1.39
3	0.02	0.02	0.07	0.11	0.22	0.35	0.58	1.01	1.42	1.87	2.37
4	0.06	0.09	0.21	0.30	0.48	0.71	1.06	1.65	2.19	2.75	3.36
5	0.16	0.21	0.41	0.55	0.83	1.15	1.61	2.34	3.00	3.66	4.35
6	0.30	0.38	0.68	0.87	1.24	1.64	2.20	3.07	3.83	4.57	5.35
7	0.48	0.60	0.99	1.24	1.69	2.17	2.73	3.62	4.67	5.49	6.35
8	0.71	0.86	1.34	1.65	2.18	2.73	3.49	4.59	5.53	6.42	7.34
9	0.97	1.15	1.73	2.09	2.70	3.33	4.17	5.38	6.39	7.36	8.34
10	1.26	1.48	2.16	2.56	3.25	3.94	4.87	6.18	7.27	8.30	9.34
11	1.59	1.83	2.60	3.05	3.82	4.57	5.58	6.99	8.15	9.24	10.34
12	1.93	2.21	3.07	3.57	4.40	5.23	6.30	7.81	9.03	10.18	11.34
13	2.31	2.62	3.57	4.11	5.01	5.89	7.04	8.63	9.93	11.13	12.34
14	2.70	3.04	4.07	4.66	5.63	6.57	7.79	9.47	10.82	12.08	13.34
15	3.11	3.48	4.60	5.23	6.26	7.26	8.55	10.31	11.72	13.03	14.34
16	3.54	3.94	5.14	5.81	6.91	7.96	9.31	11.15	12.62	13.98	15.34
17	3.98	4.42	5.70	6.41	7.56	8.67	10.09	12.00	13.53	14.94	16.34
18	4.44	4.90	6.26	7.01	8.23	9.39	10.86	12.86	14.44	15.89	17.34
19	4.91	5.41	6.84	7.63	8.91	10.12	11.65	13.72	15.35	16.85	18.34
20	5.40	5.92	7.43	8.26	9.59	10.85	12.44	14.58	16.27	17.81	19.34
21	5.90	6.45	8.03	8.90	10.28	11.59	13.24	15.44	17.18	18.77	20.34
22	6.40	6.98	8.64	9.54	10.98	12.34	14.04	16.31	18.10	19.73	21.34
23	6.92	7.53	9.26	10.20	11.69	13.09	14.85	17.19	19.02	20.69	22.34
24	7.45	8.08	9.89	10.86	12.40	13.85	15.66	18.06	19.94	21.65	23.34
25	7.99	8.65	10.52	11.52	13.12	14.61	16.47	18.94	20.87	22.62	24.34
26	8.54	9.22	11.16	12.20	13.84	15.38	17.29	19.82	21.79	23.58	25.34
27	9.09	9.80	11.81	12.88	14.57	16.15	18.11	20.70	22.72	24.54	26.34
28	9.66	10.39	12.46	13.56	15.31	16.93	18.94	21.59	23.65	25.51	27.34
29	10.23	10.99	13.12	14.26	16.05	17.71	19.77	22.48	24.58	26.48	28.34
30	10.80	11.59	13.79	14.95	16.79	18.49	20.60	23.36	25.51	27.44	29.34
40	16.91	17.92	20.71	22.16	24.43	26.51	29.05	32.34	34.87	37.13	39.34
50	23.46	24.67	27.99	29.71	32.36	34.76	37.69	41.45	46.86	51.62	56.66
60	30.34	31.74	35.53	37.48	40.48	43.19	46.46	50.64	53.81	56.62	59.33
100	59.90	61.92	67.33	70.06	74.22	77.93	82.36	87.95	92.13	95.81	99.33

Table for $\chi^2(f)$ random variable $F(x) = P(\chi^2(f) \leq x)$, where f is a parameter.

f	0.60	0.70	0.80	0.90	0.95	0.975	0.99	0.995	0.999	0.9995
1	0.71	1.07	1.64	2.71	3.84	5.02	6.63	7.88	10.83	12.12
2	1.83	2.41	3.22	4.61	5.99	7.38	9.21	10.60	13.82	15.20
3	2.95	3.66	4.64	6.25	7.81	9.35	11.34	12.84	16.27	17.73
4	4.04	4.88	5.99	7.78	9.49	11.14	13.28	14.86	18.47	20.00
5	5.13	6.06	7.29	9.24	11.07	12.83	15.09	16.75	20.52	22.11
6	6.21	7.23	8.56	10.64	12.59	14.45	16.81	18.55	22.46	24.10
7	7.28	8.38	9.80	12.02	14.07	16.01	18.48	20.28	24.32	26.02
8	8.35	9.52	11.03	13.36	15.51	17.53	20.09	21.95	26.12	27.87
9	9.41	10.66	12.24	14.68	16.92	19.02	21.67	23.59	27.88	29.67
10	10.47	11.78	13.44	15.99	18.31	20.48	23.21	25.19	29.59	31.42
11	11.53	12.90	14.63	17.28	19.68	21.92	24.72	26.76	31.26	33.14
12	12.58	14.01	15.81	18.55	21.03	23.34	26.22	28.30	32.91	34.82
13	13.64	15.12	16.98	19.81	22.36	24.74	27.69	29.82	34.53	36.48
14	14.69	16.22	18.15	21.06	23.68	26.12	29.14	31.32	36.12	38.11
15	15.73	17.32	19.31	22.31	25.00	27.49	30.58	32.80	37.70	39.72
16	16.78	18.42	20.47	23.54	26.30	28.85	32.00	34.27	39.25	41.31
17	17.82	19.51	21.61	24.77	27.59	30.19	33.41	35.72	40.79	42.88
18	18.87	20.60	22.76	25.99	28.87	31.53	34.81	37.16	42.31	44.43
19	19.91	21.69	23.90	27.20	30.14	32.85	36.19	38.58	43.82	45.97
20	20.95	22.77	25.04	28.41	31.41	34.17	37.57	40.00	45.31	47.50
21	21.99	23.86	26.17	29.62	32.67	35.48	38.93	41.40	46.80	49.01
22	23.03	24.94	27.30	30.81	33.92	36.78	40.29	42.80	48.27	50.51
23	24.07	26.02	28.43	32.01	35.17	38.08	41.64	44.18	49.73	52.00
24	25.11	27.10	29.55	33.20	36.42	39.36	42.98	45.56	51.18	53.48
25	26.14	28.17	30.68	34.38	37.65	40.65	44.31	46.93	52.62	54.95
26	27.18	29.25	31.79	35.56	38.89	41.92	45.64	48.29	54.05	56.41
27	28.21	30.32	32.91	36.74	40.11	43.19	46.96	49.64	55.48	57.86
28	29.25	31.39	34.03	37.92	41.34	44.46	48.28	50.99	56.89	59.30
29	30.28	32.46	35.14	39.09	42.56	45.72	49.59	52.34	58.30	60.73
30	31.32	33.53	36.25	40.26	43.77	46.98	50.89	53.67	59.70	62.16
40	41.62	44.16	47.27	51.81	55.76	59.34	63.69	66.77	73.40	76.09
50	51.89	54.72	58.16	63.17	67.50	71.42	76.15	79.49	86.66	89.56
60	62.13	65.23	68.97	74.40	79.08	83.30	88.38	91.95	99.61	102.69
100	102.95	106.91	111.67	118.50	124.34	129.56	135.81	140.17	149.45	153.17

(9.4) Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
 If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1 - p) \geq n - k)$.

n	k	0.05	0.10	0.15	0.20	0.25	0.30	0.35	0.40	0.45	0.50
2	0	0.9025	0.8100	0.7225	0.6400	0.5625	0.4900	0.4225	0.3600	0.3025	0.2500
	1	0.9975	0.9900	0.9775	0.9600	0.9375	0.9100	0.8775	0.8400	0.7975	0.7500
3	0	0.8574	0.7290	0.6141	0.5120	0.4219	0.3430	0.2746	0.2160	0.1664	0.1250
	1	0.9928	0.9720	0.9392	0.8960	0.8438	0.7840	0.7183	0.6480	0.5747	0.5000
4	0	0.8145	0.6561	0.5220	0.4096	0.3164	0.2401	0.1785	0.1256	0.0915	0.0625
	1	0.9860	0.9477	0.8905	0.8192	0.7383	0.6517	0.5630	0.4735	0.3910	0.3125
5	0	0.7738	0.5905	0.4437	0.3277	0.2373	0.1681	0.1160	0.0778	0.0503	0.0313
	1	0.9774	0.9185	0.8352	0.7373	0.6328	0.5282	0.4284	0.3370	0.2562	0.1875
6	0	0.7351	0.5314	0.3771	0.2621	0.1780	0.1176	0.0754	0.0467	0.0277	0.0156
	1	0.9672	0.8847	0.7765	0.6554	0.5339	0.4202	0.3191	0.2333	0.1636	0.1094
7	0	0.6983	0.4783	0.3206	0.2097	0.1335	0.0824	0.0490	0.0280	0.0152	0.0078
	1	0.9566	0.8503	0.7166	0.5767	0.4449	0.3294	0.2338	0.1586	0.1024	0.0625
8	0	0.6634	0.4305	0.2725	0.1678	0.1001	0.0576	0.0319	0.0168	0.0084	0.0039
	1	0.9428	0.8131	0.6572	0.5033	0.3671	0.2553	0.1691	0.1064	0.0632	0.0352
9	0	0.6302	0.3874	0.2316	0.1342	0.0751	0.0404	0.0207	0.0101	0.0046	0.0020
	1	0.9288	0.7748	0.5995	0.4362	0.3003	0.1960	0.1211	0.0705	0.0385	0.0195

Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
 If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1 - p) \geq n - k)$.

n	k	0.05	0.10	0.15	0.20	0.25	0.30	0.35	0.40	0.45	0.50
10	0	0.5987	0.3487	0.1969	0.1074	0.0563	0.0282	0.0135	0.0060	0.0025	0.0010
	1	0.9139	0.7361	0.5443	0.3758	0.2440	0.1493	0.0860	0.0464	0.0233	0.0107
11	0	0.5688	0.3138	0.1673	0.0859	0.0422	0.0198	0.0088	0.0036	0.0014	0.0005
	1	0.8981	0.6974	0.4922	0.3221	0.1971	0.1130	0.0606	0.0302	0.0139	0.0059
12	0	0.5404	0.2824	0.1422	0.0687	0.0317	0.0138	0.0057	0.0022	0.0008	0.0002
	1	0.8816	0.6590	0.4435	0.2749	0.1584	0.0850	0.0424	0.0196	0.0083	0.0032

(9.5) Table for Poisson random variable $P(Po(\mu) \leq k)$.

k	μ									
	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0
0	0.9048	0.8187	0.7408	0.6703	0.6065	0.5488	0.4966	0.4493	0.4066	0.3679
1	0.9953	0.9825	0.9631	0.9384	0.9098	0.8781	0.8442	0.8088	0.7725	0.7358
2	0.9998	0.9989	0.9964	0.9921	0.9856	0.9769	0.9659	0.9526	0.9371	0.9197
3	1.0000	0.9999	0.9997	0.9992	0.9982	0.9966	0.9942	0.9909	0.9865	0.9810
4	1.0000	1.0000	1.0000	0.9999	0.9998	0.9996	0.9992	0.9986	0.9977	0.9963
5	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9997	0.9994
6	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999
7	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
k	μ									
	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0
0	0.3329	0.3012	0.2725	0.2466	0.2231	0.2019	0.1827	0.1653	0.1496	0.1353
1	0.6990	0.6626	0.6268	0.5918	0.5578	0.5249	0.4932	0.4628	0.4337	0.4060
2	0.9004	0.8795	0.8571	0.8335	0.8088	0.7834	0.7572	0.7306	0.7037	0.6767
3	0.9743	0.9662	0.9569	0.9463	0.9344	0.9212	0.9068	0.8913	0.8747	0.8571
4	0.9946	0.9923	0.9893	0.9857	0.9814	0.9763	0.9704	0.9636	0.9559	0.9473
5	0.9990	0.9985	0.9978	0.9968	0.9955	0.9940	0.9920	0.9896	0.9868	0.9834
6	0.9999	0.9997	0.9996	0.9994	0.9991	0.9987	0.9981	0.9974	0.9966	0.9955
7	1.0000	1.0000	0.9999	0.9999	0.9998	0.9997	0.9996	0.9994	0.9992	0.9989
8	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9999	0.9998
9	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
k	μ									
	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0
0	0.1225	0.1108	0.1003	0.0907	0.0821	0.0743	0.0672	0.0608	0.0550	0.0498
1	0.3796	0.3546	0.3309	0.3084	0.2873	0.2674	0.2487	0.2311	0.2146	0.1991
2	0.6496	0.6227	0.5960	0.5697	0.5438	0.5184	0.4936	0.4695	0.4460	0.4232
3	0.8386	0.8194	0.7993	0.7787	0.7576	0.7360	0.7141	0.6919	0.6696	0.6472
4	0.9379	0.9275	0.9162	0.9041	0.8912	0.8774	0.8629	0.8477	0.8318	0.8153
5	0.9796	0.9751	0.9700	0.9643	0.9580	0.9510	0.9433	0.9349	0.9258	0.9161
6	0.9941	0.9925	0.9906	0.9884	0.9858	0.9828	0.9794	0.9756	0.9713	0.9665
7	0.9985	0.9980	0.9974	0.9967	0.9958	0.9947	0.9934	0.9919	0.9901	0.9881
8	0.9997	0.9995	0.9994	0.9991	0.9989	0.9985	0.9981	0.9976	0.9969	0.9962
9	0.9999	0.9999	0.9999	0.9998	0.9997	0.9996	0.9995	0.9993	0.9991	0.9989
10	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9999	0.9998	0.9997	0.9996
11	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9999
12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

Table for Poisson random variable $P(Po(\mu) \leq k)$.

k	μ									
	3.2	3.4	3.6	3.8	4.0	4.2	4.4	4.6	4.8	5.0
0	0.0408	0.0334	0.0273	0.0224	0.0183	0.0150	0.0123	0.0101	0.0082	0.0067
1	0.1712	0.1468	0.1257	0.1074	0.0916	0.0780	0.0663	0.0563	0.0477	0.0404
2	0.3799	0.3397	0.3027	0.2689	0.2381	0.2102	0.1851	0.1626	0.1425	0.1247
3	0.6025	0.5584	0.5152	0.4735	0.4335	0.3954	0.3594	0.3257	0.2942	0.2650
4	0.7806	0.7442	0.7064	0.6678	0.6288	0.5898	0.5512	0.5132	0.4763	0.4405
5	0.8946	0.8705	0.8441	0.8156	0.7851	0.7531	0.7199	0.6858	0.6510	0.6160
6	0.9534	0.9421	0.9267	0.9091	0.8893	0.8675	0.8436	0.8180	0.7908	0.7622
7	0.9832	0.9769	0.9682	0.9599	0.9509	0.9419	0.9314	0.9204	0.9087	0.8866
8	0.9943	0.9917	0.9883	0.9840	0.9786	0.9721	0.9642	0.9549	0.9442	0.9319
9	0.9982	0.9973	0.9960	0.9942	0.9919	0.9889	0.9851	0.9805	0.9749	0.9682
10	0.9995	0.9992	0.9987	0.9981	0.9972	0.9959	0.9943	0.9922	0.9896	0.9863
11	0.9999	0.9998	0.9996	0.9994	0.9991	0.9986	0.9980	0.9971	0.9960	0.9945
12	1.0000	0.9999	0.9999	0.9999	0.9998	0.9997	0.9996	0.9993	0.9990	0.9986
13	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9996	0.9993	0.9990	0.9986
14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9997	0.9995
15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9998
k	μ									
	5.2	5.4	5.6	5.8	6.0	6.5	7.0	7.5	8.0	8.5
0	0.0055	0.0045	0.0037	0.0030	0.0025	0.0015	0.0009	0.0006	0.0003	0.0002
1	0.0342	0.0289	0.0244	0.0206	0.0174	0.0113	0.0073	0.0047	0.0030	0.0019
2	0.1088	0.0948	0.0824	0.0715	0.0620	0.0430	0.0286	0.0203	0.0138	0.0093
3	0.2381	0.2133	0.1906	0.1700	0.1512	0.1118	0.0818	0.0591	0.0424	0.0301
4	0.4061	0.3733	0.3422	0.3127	0.2851	0.2237	0.1730	0.1321	0.0996	0.0744
5	0.5809	0.5461	0.5119	0.4783	0.4457	0.3690	0.3007	0.2414	0.1912	0.1496
6	0.7324	0.7017	0.6703	0.6384	0.6063	0.5265	0.4497	0.3782	0.3134	0.2562
7	0.8449	0.8217	0.7970	0.7710	0.7440	0.6728	0.5987	0.5246	0.4530	0.3856
8	0.9181	0.9027	0.8857	0.8672	0.8472	0.7916	0.7291	0.6620	0.5925	0.5231
9	0.9603	0.9512	0.9409	0.9292	0.9161	0.8774	0.8305	0.7764	0.7166	0.6530
10	0.9823	0.9775	0.9718	0.9651	0.9574	0.9332	0.9015	0.8622	0.8159	0.7634
11	0.9927	0.9904	0.9875	0.9841	0.9799	0.9661	0.9467	0.9208	0.8881	0.8487
12	0.9972	0.9962	0.9949	0.9932	0.9912	0.9840	0.9730	0.9573	0.9362	0.9091
13	0.9990	0.9986	0.9980	0.9973	0.9964	0.9929	0.9872	0.9784	0.9658	0.9486
14	0.9999	0.9995	0.9993	0.9990	0.9986	0.9970	0.9943	0.9897	0.9827	0.9726
15	0.9999	0.9998	0.9998	0.9996	0.9995	0.9988	0.9976	0.9954	0.9918	0.9862
16	1.0000	0.9999	0.9999	0.9999	0.9998	0.9996	0.9990	0.9980	0.9963	0.9934
17	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9996	0.9992	0.9984	0.9970
18	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9997	0.9993	0.9987
19	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9997	0.9995
20	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9998
21	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999
22	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

Table for Poisson random variable $P(Po(\mu) \leq k)$.

k	μ																
	9.0	9.5	10.0	11.0	12.0	13.0	14.0	15.0	16.0	17.0							
0	0.0001	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000							
1	0.0012	0.0008	0.0005	0.0002	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000							
2	0.0062	0.0042	0.0028	0.0012	0.0005	0.0002	0.0001	0.0000	0.0000	0.0000							
3	0.0212	0.0149	0.0103	0.0049	0.0023	0.0011	0.0005	0.0002	0.0001	0.0000							
4	0.0550	0.0403	0.0293	0.0151	0.0076	0.0037	0.0018	0.0009	0.0004	0.0002							
5	0.1157	0.0885	0.0671	0.0375	0.0203	0.0107	0.0055	0.0028	0.0014	0.0007							
6	0.2068	0.1649	0.1301	0.0786	0.0458	0.0259	0.0142	0.0076	0.0040	0.0021							
7	0.3239	0.2687	0.2202	0.1432	0.0895	0.0540	0.0316	0.0180	0.0100	0.0054							
8	0.4557	0.3918	0.3328	0.2320	0.1550	0.0998	0.0621	0.0374	0.0220	0.0126							
9	0.5874	0.5218	0.4579	0.3405	0.2424	0.1658	0.1094	0.0699	0.0433	0.0261							
10	0.7060	0.6453	0.5830	0.4599	0.3472	0.2517	0.1757	0.1185	0.0774	0.0491							
11	0.8030	0.7520	0.6968	0.5793	0.4616	0.3532	0.2600	0.1848	0.1270	0.0847							
12	0.8758	0.8364	0.7916	0.6887	0.5760	0.4631	0.3585	0.2676	0.1931	0.1350							
13	0.9261	0.8981	0.8645	0.7813	0.6815	0.5730	0.4644	0.3632	0.2745	0.2009							
14	0.9585	0.9400	0.9165	0.8540	0.7720	0.6751	0.5704	0.4657	0.3675	0.2808							
15	0.9780	0.9665	0.9513	0.9074	0.8444	0.7636	0.6694	0.5681	0.4667	0.3715							
16	0.9889	0.9823	0.9730	0.9441	0.8987	0.8355	0.7559	0.6641	0.5660	0.4677							
17	0.9947	0.9911	0.9857	0.9678	0.9370	0.8905	0.8272	0.7489	0.6593	0.5640							
18	0.9976	0.9957	0.9928	0.9823	0.9626	0.9302	0.8826	0.8195	0.7423	0.6550							
19	0.9989	0.9980	0.9965	0.9907	0.9787	0.9573	0.9235	0.8752	0.8122	0.7363							
20	0.9996	0.9991	0.9984	0.9953	0.9884	0.9750	0.9521	0.9170	0.8682	0.8055							
21	0.9998	0.9996	0.9993	0.9977	0.9939	0.9859	0.9712	0.9469	0.9108	0.8615							
22	0.9999	0.9999	0.9997	0.9990	0.9970	0.9924	0.9833	0.9673	0.9418	0.9047							
23	1.0000	0.9999	0.9999	0.9995	0.9985	0.9960	0.9907	0.9805	0.9633	0.9367							
24	1.0000	1.0000	1.0000	0.9998	0.9993	0.9980	0.9950	0.9888	0.9777	0.9594							
25	1.0000	1.0000	1.0000	0.9999	0.9997	0.9990	0.9974	0.9938	0.9869	0.9748							
26	1.0000	1.0000	1.0000	1.0000	0.9999	0.9995	0.9987	0.9967	0.9925	0.9848							
27	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9994	0.9983	0.9959	0.9912							
28	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9991	0.9978	0.9950							
29	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9996	0.9994	0.9986							
30	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9997	0.9993							
31	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9996							
32	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9996							
33	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9999							
34	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999							
35	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000							