

Examiner: Xiangfeng Yang (Tel: 013 28 57 88).**Things allowed (Hjälpmedel):** a calculator.**Scores rating (Betygsgränser):** 8-11 points giving rate 3; 11.5-14.5 points giving rate 4; 15-18 points giving rate 5.

1 (3 points)

Three students (marked as A , B and C) take an exam. Assume that these three students are independent and the passing exam probabilities for the three students are $P(A) = 0.8$, $P(B) = 0.6$ and $P(C) = 0.1$.

(1.1) (1p) Find the probability that exactly two students pass the exam.

(1.2) (1p) Find the probability that at least one student passes the exam.

(1.3) (1p) Given that at least one student passes the exam, find the probability that C passes the exam.

Solution. (1.1)

$$\begin{aligned} P(\text{exactly two students pass the exam}) &= P(A \text{ and } B \text{ pass}) + P(A \text{ and } C \text{ pass}) + P(B \text{ and } C \text{ pass}) \\ &= P(A \cap B \cap C') + P(A \cap C \cap B') + P(B \cap C \cap A') \\ &= 0.8 \cdot 0.6 \cdot 0.9 + 0.8 \cdot 0.1 \cdot 0.4 + 0.6 \cdot 0.1 \cdot 0.2 \\ &= 0.432 + 0.032 + 0.012 = 0.476. \end{aligned}$$

(1.2)

$$\begin{aligned} P(\text{at least one student passes the exam}) &= P(A \cup B \cup C) = 1 - P(A' \cap B' \cap C') \\ &= 1 - P(\text{no students pass the exam}) \\ &= 1 - P(A') \cdot P(B') \cdot P(C') \\ &= 1 - 0.2 \cdot 0.4 \cdot 0.9 = 1 - 0.072 = 0.928. \end{aligned}$$

(1.3)

$$\begin{aligned} P(C|A \cup B \cup C) &= \frac{P(C \cap (A \cup B \cup C))}{P(A \cup B \cup C)} = \frac{P(C)}{P(A \cup B \cup C)} \\ &= \frac{0.1}{0.928} = 0.1078. \end{aligned}$$

□

2 (3 points)

Let X be a continuous random variable with a probability density function (pdf) as follows

$$f(x) = \begin{cases} c \cdot x^3, & \text{if } 0 \leq x \leq 1 \\ 0, & \text{otherwise.} \end{cases}$$

(2.1) (1p) Find the value of the constant c in order that $f(x)$ is indeed a pdf.

(2.2) (1p) Find the mean $E(X)$ of X and the variance $V(X)$ of X .

(2.3) (1p) Find the 95-th percentile b (that is, find b such that $P(X \leq b) = 0.95$).

(Hint: you might need to use the integral $\int_a^b x^n dx = \left[\frac{1}{n+1} x^{n+1} \right]_{x=a}^{x=b} = \frac{1}{n+1} (b^{n+1} - a^{n+1})$ for any integer $n \geq 0$.)

Solution. (2.1) In order that $f(x)$ becomes a pdf, it is required that

$$1 = \int_{-\infty}^{\infty} f(x) dx = \int_0^1 c \cdot x^3 dx = \left[\frac{c}{4} x^4 \right]_{x=0}^{x=1} = \frac{c}{4} \implies c = 4.$$

(2.2)

$$E(X) = \int_{-\infty}^{\infty} x \cdot f(x) dx = \int_0^1 c \cdot x^4 dx = \left[\frac{c}{5} x^5 \right]_{x=0}^{x=1} = \frac{c}{5} = \frac{4}{5} = 0.8.$$

$$\begin{aligned}
V(X) &= E(X^2) - (E(X))^2 = \int_{-\infty}^{\infty} x^2 \cdot f(x) dx - (E(X))^2 = \int_0^1 c \cdot x^5 dx - (E(X))^2 \\
&= \left[\frac{c}{6} x^6 \right]_{x=0}^{x=1} - (E(X))^2 = \frac{c}{6} - \left(\frac{c}{5} \right)^2 \\
&= \frac{4}{6} - \left(\frac{4}{5} \right)^2 = 0.6667 - 0.64 = 0.0267.
\end{aligned}$$

(2.3)

$$0.95 = P(X \leq b) = \int_{-\infty}^b f(x) dx = \int_0^b c \cdot x^3 dx = \left[\frac{c}{4} x^4 \right]_{x=0}^{x=b} = \frac{c \cdot b^4}{4} \implies b = \left(\frac{4 \cdot 0.95}{c} \right)^{1/4} = 0.9873.$$

□

3 (3 points)

Suppose that the numbers of cars sold in each week by Askling Bil in Linköping follow a Poisson distribution $Po(1)$ with parameter $\lambda = 1$. Assume that the numbers of cars sold in different weeks are independent. Find the probability that there are at most 70 cars sold within one year (52 weeks) by Askling Bil in Linköping.

Solution. Let X_1, X_2, \dots, X_{52} denote the numbers of cars sold in these 52 weeks, then X_1, X_2, \dots, X_{52} are independent and each $X_i \sim Po(1)$. Therefore $\mu = E(X_i) = 1$ and $\sigma^2 = V(X_i) = 1$.

$$\begin{aligned}
P(\text{at most 70 cars sold within one year (52 weeks)}) &= P(X_1 + X_2 + \dots + X_{52} \leq 70) \\
&= P(\bar{X} \leq 70/52) = P\left(\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \leq \frac{70/52 - \mu}{\sigma/\sqrt{n}}\right) \\
&= P(N(0, 1) \leq \frac{70/52 - 1}{1/\sqrt{52}}) = P(N(0, 1) \leq 2.50) = \Phi(2.50) = 0.9938.
\end{aligned}$$

□

4 (3 points)

A population X is a discrete random variable with the following probability mass function (pmf)

X	0	1	2
$p(x)$	$1 - 3\theta$	θ	2θ

where $0 < \theta < 1/3$ is an unknown parameter. A sample $\{X_1, X_2, \dots, X_n\}$ is taken from this population.

(4.1) (2p) Use the method of moments to find a point estimator $\hat{\theta}_{MM}$ of θ .

(4.2) (1p) Find the variance $V(\hat{\theta}_{MM})$ of the point estimator $\hat{\theta}_{MM}$.

Solution. (4.1) The mean $E(X)$ of the population can be computed as $E(X) = 0 \cdot (1 - 3\theta) + 1 \cdot \theta + 2 \cdot 2\theta = 5\theta$. It is then from the equation $E(X) = \bar{X}$ that

$$5\theta = \bar{X} \implies \hat{\theta}_{MM} = \bar{X}/5.$$

(4.2)

$$\begin{aligned}
V(\hat{\theta}_{MM}) &= V(\bar{X}/5) = \frac{1}{25} V(\bar{X}) = \frac{1}{25} V\left(\frac{X_1 + X_2 + \dots + X_n}{n}\right) \\
&= \frac{1}{25} \frac{1}{n^2} V(X_1 + X_2 + \dots + X_n) = \frac{1}{25} \frac{1}{n^2} [V(X_1) + V(X_2) + \dots + V(X_n)] \\
&= \frac{1}{25} \frac{1}{n^2} n V(X) = \frac{V(X)}{25n} = \frac{9\theta - 25\theta^2}{25n},
\end{aligned}$$

where

$$V(X) = E(X^2) - (E(X))^2 = [0^2 \cdot (1 - 3\theta) + 1^2 \cdot \theta + 2^2 \cdot 2\theta] - (5\theta)^2 = 9\theta - 25\theta^2.$$

□

5 (3 points)

- The lengths X of a certain items follow a normal distribution $X \sim N(\mu, \sigma^2)$. In order to study the unknown parameters μ and σ^2 , a sample is taken from X with sample size 9, sample mean 5.2 and sample standard deviation 0.28.
- (5.1) (1p) In order to check whether or not $\mu > 5$, construct an appropriate one-sided 95% confidence interval of μ .
 - (5.2) (1p) With a significance level $\alpha = 5\%$, test the hypotheses $H_0 : \mu = 5$ against $H_a : \mu > 5$.
 - (5.3) (1p) Construct a two-sided 95% confidence interval of σ^2 .

Solution. (5.1) To check whether or not $\mu > 5$, one constructs one-sided confidence interval as follows

$$\begin{aligned} I_\mu &= (\bar{x} - t_{\alpha}(n-1) \cdot s/\sqrt{n}, +\infty) \\ &= (5.2 - t_{0.05}(9-1) \cdot 0.28/\sqrt{9}, +\infty) \\ &= (5.2 - 1.86 \cdot 0.28/\sqrt{9}, +\infty) \\ &= (5.2 - 0.1736, +\infty) = (5.0264, +\infty). \end{aligned}$$

As $5.0264 > 5$, yes, the sample provides evidence showing that $\mu > 5$.

(5.2)

$$TS = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} = 2.1429, \quad C = (t_{\alpha}(n-1), +\infty) = (t_{0.05}(9-1), +\infty) = (1.86, +\infty).$$

It is from $TS \in C$ that H_0 is rejected (namely, $\mu > 5$).

(5.3) The confidence interval is

$$\begin{aligned} I_{\sigma^2} &= \left(\frac{(n-1)s^2}{\chi_{\alpha/2}^2(n-1)}, \frac{(n-1)s^2}{\chi_{1-\alpha/2}^2(n-1)} \right) = \left(\frac{(9-1)0.28^2}{\chi_{0.025}^2(9-1)}, \frac{(9-1)0.28^2}{\chi_{0.975}^2(9-1)} \right) \\ &= \left(\frac{(9-1)0.28^2}{\chi_{0.025}^2(9-1)}, \frac{(9-1)0.28^2}{\chi_{0.975}^2(9-1)} \right) = \left(\frac{0.6272}{17.53}, \frac{0.6272}{2.18} \right) \\ &= (0.0358, 0.2877). \end{aligned}$$

□

6 (3 points)

A coin (with two sides ‘head’ and ‘tail’) has been thrown 100 times and the results are:

Outcome	head	tail
Frequency	55	45

Do the results provide any evidence that the coin is unfair?

(6.1) (1p) Let p be the ‘head’ probability. Test the following hypotheses with a 5% significance level:

$$H_0 : p = 0.5 \quad \text{against} \quad H_a : p \neq 0.5.$$

(6.2) (2p) Let p_1 be the ‘head’ probability and p_2 be the ‘tail’ probability. Perform a χ^2 test for the following hypotheses with a 5% significance level:

$$H_0 : p_1 = 0.5 \text{ and } p_2 = 0.5 \quad H_a : p_1 \neq 0.5 \text{ and } p_2 \neq 0.5$$

Solution. (6.1)

$$\begin{aligned} TS &= \frac{\hat{p} - p_0}{\sqrt{p_0(1-p_0)/n}} = \frac{55/100 - 0.5}{\sqrt{0.5(1-0.5)/100}} = 1, \\ C &= (-\infty, -z_{\alpha/2}) \cup (z_{\alpha/2}, +\infty) = (-\infty, -1.96) \cup (1.96, +\infty). \end{aligned}$$

Since $TS \notin C$, we do NOT reject H_0 (which implies that the results do NOT provide any evidence that the coin is unfair).

(6.2)

$$\begin{aligned} TS &= \sum_{i=1}^k \frac{(n_i - np_i)^2}{np_i} = \frac{(55 - 100 \cdot 0.5)^2}{100 \cdot 0.5} + \frac{(45 - 100 \cdot 0.5)^2}{100 \cdot 0.5} = 1, \\ C &= (\chi_{\alpha}^2(k-1), +\infty) = (\chi_{0.05}^2(2-1), +\infty) = (3.84, +\infty). \end{aligned}$$

Since $TS \notin C$, we do NOT reject H_0 (which implies that the results do NOT provide any evidence that the coin is unfair).

□

1. Basic probability

(1.1) Conditional probability $P(A|B) = \frac{P(A \cap B)}{P(B)}$.

(1.2) Total probability $P(B) = \sum_{i=1}^k P(B|A_i)P(A_i)$ where $\{A_i\}$ are disjoint and $\cup_{i=1}^k A_i = S$.

(1.3) Bayes' Theorem $P(A_j|B) = \frac{P(B|A_j)P(A_j)}{\sum_{i=1}^k P(B|A_i)P(A_i)}$ where $\{A_i\}$ are in (1.2).

2. Random variables (r.v.s)

(2.1) Discrete r.v. X has a pmf $p(x) = P(X = x)$ satisfying $p(x) \geq 0$ and $\sum p(x_i) = 1$,

X	x_1	x_2	\cdots	x_n	\cdots
$p(x)$	$p(x_1)$	$p(x_2)$	\cdots	$p(x_n)$	\cdots

Expectation (or *Expected value* or *mean*) $\mu_X = E(X) = \sum x_i p(x_i)$;
 Variance $\sigma_X^2 = V(X) = E(X - \mu_X)^2 = E(X^2) - \mu_X^2 = \sum x_i^2 p(x_i) - (\sum x_i p(x_i))^2$.

(2.2) Continuous r.v. X has a pdf $f(x)$ satisfying $f(x) \geq 0$ and $\int_{-\infty}^{\infty} f(x)dx = 1$,

$$P(a < X < b) = \int_a^b f(x)dx.$$

Expectation (or *Expected value* or *mean*) $\mu_X = E(X) = \int_{-\infty}^{\infty} xf(x)dx$;

Variance $\sigma_X^2 = V(X) = E(X - \mu_X)^2 = E(X^2) - \mu_X^2 = \int_{-\infty}^{\infty} x^2 f(x)dx - (\int_{-\infty}^{\infty} xf(x)dx)^2$.

(2.3) Cumulative distribution function (cdf) of a r.v. X is $F(x) = P(X \leq x)$.

(2.4) X and Y are r.v.s, a , b and c are scalars, then

$$E(aX + bY + c) = aE(X) + bE(Y) + c,$$

$$V(aX + bY + c) = a^2 V(X) + b^2 V(Y) + 2ab \text{cov}(X, Y),$$

$$E(g(X, Y)) = \begin{cases} \sum_{i,j} g(x_i, y_j) \cdot p(x_i, y_j), & \text{for discrete } (X, Y), \\ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) \cdot f(x, y)dxdy, & \text{for continuous } (X, Y). \end{cases}$$

(2.5) • Discrete r.v. (X, Y) has a joint pmf $p(x, y)$ satisfying $p(x, y) \geq 0$ and $\sum_{x_i} \sum_{y_i} p(x_i, y_i) = 1$.

The marginal pmf of X is $p_X(x) = \sum_y p(x, y)$;

The marginal pmf of Y is $p_Y(y) = \sum_x p(x, y)$;

X and Y are *independent* if $p(x, y) = p_X(x) \cdot p_Y(y)$.

• Continuous r.v. (X, Y) has a joint pdf $f(x, y)$ satisfying $f(x, y) \geq 0$ and $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y)dxdy = 1$.

The marginal pdf of X is $f_X(x) = \int_{-\infty}^{\infty} f(x, y)dy$;

The marginal pdf of Y is $f_Y(y) = \int_{-\infty}^{\infty} f(x, y)dx$;

X and Y are *independent* if $f(x, y) = f_X(x) \cdot f_Y(y)$.

3. Several special r.v.s

(3.1) $X \sim Bin(n, p)$ has a pmf $p(x) = P(X = x) = \binom{n}{x} \cdot p^x \cdot (1-p)^{n-x}$, $x = 0, 1, 2, \dots, n$.

$E(X) = n \cdot p$, $V(X) = n \cdot p \cdot (1-p)$.

(3.2) $X \sim Po(\lambda)$ has a pmf $p(x) = P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$, $x = 0, 1, 2, \dots$.

$E(X) = \lambda$, $V(X) = \lambda$.

(3.3) $X \sim Hypergeometric$ has a pmf $p(x) = P(X = x) = \frac{\binom{M}{x} \binom{N-M}{n-x}}{\binom{N}{n}}$.

(3.4) $X \sim Exp(\lambda)$ has a pdf

$$f(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0, \\ 0, & \text{otherwise.} \end{cases}$$

(3.5) $X \sim N(\mu, \sigma^2)$ has a pdf

$$f(x) = \frac{1}{\sigma \sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, -\infty < x < \infty.$$

$E(X) = \mu$, $V(X) = \sigma^2$

(3.6) $X \sim U(a, b)$ has a pdf

$$f(x) = \begin{cases} \frac{1}{b-a}, & a < x < b, \\ 0, & \text{otherwise.} \end{cases}$$

$$E(X) = \frac{a+b}{2}, \quad V(X) = \frac{(b-a)^2}{12}.$$

4. Central Limit Theorem (CLT)

Suppose that a population has mean = μ and variance = σ^2 . A random sample $\{X_1, X_2, \dots, X_n\}$ from this population is given. Then for large $n \geq 30$,

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1).$$

- If the population is normal, then (1) holds for any n .
- Note that $\mu = E(\bar{X})$ and $(\sigma/\sqrt{n})^2 = V(\bar{X})$.

5. Several notations in statistics

(5.1) Sample mean: $\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n} = \frac{\sum X_i}{n}$; $\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{\sum x_i}{n}$.

(5.2) Sample variance:

$$S^2 = \frac{\sum (X_i - \bar{X})^2}{n-1} = \frac{1}{n-1} \left(\sum X_i^2 - \frac{(\sum X_i)^2}{n} \right); \quad s^2 = \frac{\sum (x_i - \bar{x})^2}{n-1} = \frac{1}{n-1} \left(\sum x_i^2 - \frac{(\sum x_i)^2}{n} \right).$$

- Capital letters X and S^2 refer to the objects based on random sample (therefore they are in general r.v.s), while small letters \bar{x} and s^2 are the objects based on observations (so they are scalars).

(5.3) A point estimator of θ obtained by Maximum Likelihood method is denoted as $\hat{\theta}_{ML}$.

6. Confidence Interval (CI)

In this course, three types of confidence intervals are studied depending on the unknown population parameter(s): CI-1 (confidence intervals for population mean(s)), CI-2 (confidence intervals for population variance(s)), and CI-3 (confidence intervals for population proportion(s)).

CI-1: $(1 - \alpha)$ CI of a population mean μ

case 1.1 (any n) If population $X \sim N(\mu, \sigma^2)$ and σ^2 is known, then $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$ and

$$I_\mu = \left(\bar{x} - z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}, \bar{x} + z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \right) := \bar{x} \mp z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}.$$

case 1.2 ($n \geq 30$) For any population X , it holds that $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$ and

$$I_\mu = \bar{x} \mp z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \text{ or } I_\mu = \bar{x} \mp z_{\alpha/2} \cdot \frac{\hat{\sigma}}{\sqrt{n}}.$$

case 1.3 (any n) If population $X \sim N(\mu, \sigma^2)$ and σ^2 is unknown, then $\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim T(n-1)$ and

$$I_\mu = \bar{x} \mp t_{\alpha/2}(n-1) \cdot \frac{s}{\sqrt{n}}.$$

CI-1': $(1 - \alpha)$ CI of the difference of two population means $\mu_X - \mu_Y$

case 1.1' (any n_1, n_2) If independent populations $X \sim N(\mu_X, \sigma_X^2)$, $Y \sim N(\mu_Y, \sigma_Y^2)$, and σ_X^2, σ_Y^2 are known, then $\frac{(\bar{X} - \bar{Y}) - (\mu_X - \mu_Y)}{\sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}} \sim N(0, 1)$, and $I_{\mu_X - \mu_Y} = (\bar{x} - \bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}$.

case 1.2' ($n_1, n_2 \geq 30$) For any independent populations X and Y , it holds that


$$\frac{(\bar{X} - \bar{Y}) - (\mu_X - \mu_Y)}{\sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}} \sim N(0, 1) \text{ and}$$

$$I_{\mu_X - \mu_Y} = (\bar{x} - \bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}} \text{ or } I_{\mu_X - \mu_Y} = (\bar{x} - \bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{\sigma}_X^2}{n_1} + \frac{\hat{\sigma}_Y^2}{n_2}}.$$

case 1.3' (any n_1, n_2) If independent populations $X \sim N(\mu_X, \sigma_X^2)$, $Y \sim N(\mu_Y, \sigma_Y^2)$, where σ_X^2, σ_Y^2 are unknown but $\sigma_X^2 = \sigma_Y^2$, then

$$\frac{(\bar{X} - \bar{Y}) - (\mu_X - \mu_Y)}{\sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} \sim T(n_1 + n_2 - 2), \text{ where } S^2 = \frac{(n_1 - 1)S_X^2 + (n_2 - 1)S_Y^2}{n_1 + n_2 - 2}, \text{ and}$$

$$I_{\mu_X - \mu_Y} = (\bar{x} - \bar{y}) \mp t_{\alpha/2}(n_1 + n_2 - 2) \cdot s \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}.$$



CI-2: $(1 - \alpha)$ CI of population variance(s) σ^2

- If a population $X \sim N(\mu, \sigma^2)$ and σ^2 is unknown, then $\frac{(n-1)S^2}{\sigma^2} \sim \chi^2(n-1)$, and

$$I_{\sigma^2} = \left(\frac{(n-1)s^2}{\chi_{\alpha/2}^2(n-1)}, \frac{(n-1)s^2}{\chi_{1-\alpha/2}^2(n-1)} \right).$$

If two independent populations $X \sim N(\mu_X, \sigma^2)$ and $Y \sim N(\mu_Y, \sigma^2)$, and σ^2 is unknown, then $\frac{(n_1+n_2-2)S^2}{\sigma^2} \sim \chi^2(n_1 + n_2 - 2)$, and

$$I_{\sigma^2} = \left(\frac{(n_1 + n_2 - 2)s^2}{\chi_{\alpha/2}^2(n_1 + n_2 - 2)}, \frac{(n_1 + n_2 - 2)s^2}{\chi_{1-\alpha/2}^2(n_1 + n_2 - 2)} \right),$$

where $S^2 = \frac{(n_1-1)S_X^2 + (n_2-1)S_Y^2}{n_1 + n_2 - 2}$.

CI-3: $(1 - \alpha)$ CI of population proportion(s)

- If a (large) population has an unknown proportion p , then $\frac{\hat{p} - p}{\sqrt{(1-p)/n}} \sim N(0, 1)$ if $n\hat{p} \geq 10$ and $n(1-\hat{p}) \geq 10$ with $\hat{p} = x/n$, and $I_p = \hat{p} \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$.
- If two independent (large) populations have unknown proportions p_1 and p_2 , then

$$\frac{(\hat{p}_1 - \hat{p}_2) - (p_1 - p_2)}{\sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}}} \sim N(0, 1)$$

if $n_i\hat{p}_i \geq 10$ and $n_i(1 - \hat{p}_i) \geq 10$ for $i = 1, 2$, and $I_{p_1 - p_2} = (\hat{p}_1 - \hat{p}_2) \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{\hat{p}_2(1-\hat{p}_2)}{n_2}}$.

7. Hypothesis Test (HT)

	H_0 is true	H_0 is false and $\theta = \theta_1$
reject H_0	(type I error or significance level) α	(power) $h(\theta_1) = 1 - h(\theta_1)$
don't reject H_0	1 - α	(type II error) $\beta(\theta_1) = 1 - h(\theta_1)$

reject $H_0 \Leftrightarrow TS \in C \Leftrightarrow p\text{-value} < \alpha$

χ^2 tests for populations (non-parametric tests)

Suppose that for a random sample of a population X the n elements of it are classified into k disjoint groups $A_i, 1 \leq i \leq k$. For each group $A_i, 1 \leq i \leq k$, suppose that there are $N_i, 1 \leq i \leq k$ elements inside. Let $p_i = P(A_i)$ assuming a given distribution of X . Note that $p_1 + p_2 + \dots + p_k = 1$ and $N_1 + N_2 + \dots + N_k = n$. One wants to test the hypotheses

$$H_0 : P(A_i) = p_i, \quad 1 \leq i \leq k, \quad H_a : P(A_i) \neq p_i \text{ for some } 1 \leq i \leq k.$$

If n is large in the sense that $np_i \geq 5$ for all $1 \leq i \leq k$, then the test statistic is

$$\sum_{i=1}^k \frac{(N_i - np_i)^2}{np_i} \approx \chi^2(k-1).$$

Therefore the observation of the test statistic is

$$TS = \sum_{i=1}^k \frac{(n_i - np_i)^2}{np_i}, \text{ where } n_i \text{ is the observation of } N_i, 1 \leq i \leq k.$$

For the critical region C , one can take (note that if H_0 is true, then TS should be close to zero)

$$C = (\chi_\alpha^2(k-1), \infty).$$

The conclusion would be $TS \in C \iff H_0$ is rejected.

8. Linear and logistic regression

(Multiple) linear regression: $Y = \beta_0 + \beta_1 x_1 + \dots + \beta_k x_k + \varepsilon, \varepsilon \sim N(0, \sigma^2)$.

\bullet Y : response variable (which is normal r.v.), $\{x_1, \dots, x_k\}$: predictors (which are scalars).

\bullet sample: $\{(x_{11}, \dots, x_{1k}; y_1), (x_{21}, \dots, x_{2k}; y_2), \dots, (x_{n1}, \dots, x_{nk}; y_n)\}$.
 \bullet how to estimate $\beta_j \approx \hat{\beta}_j$: least square method, that is, to minimize $\sum_{i=1}^n (\hat{y}_i - y_i)^2$, where the estimated (multiple) linear regression line \hat{y} is

$$\hat{y} = \beta_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k.$$

\bullet $\frac{\hat{\beta}_j - \beta_j}{s.e(\hat{\beta}_j)} \sim T(n-k-1)$, this helps determine whether or not the real $\beta_j = 0$?

\bullet $\sigma^2 \approx \frac{SSE}{n-k-1}$, this gives an estimation of the size of the error.

$\bullet R^2 = \frac{SSE}{SS_T}$ this gives how well the model is (if $R^2 \approx 1$, then the model fits the sample very well).

\bullet How to test $\beta_1 = \dots = \beta_k = 0$? Use the random variable $\frac{SS_R/k}{SSE/(n-k-1)} \sim F(k, n-k-1)$.

Logistic regression: Let Y can only take 0 or 1 with $P(Y=1) = p$ and $P(Y=0) = 1-p$,

$$E(Y) = p(x_1, \dots, x_k) = \frac{e^{\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k}}{1 + e^{\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k}}.$$

$\bullet Y$: response variable (which is Bernoulli r.v. $P(Y=1) = p$ and $P(Y=0) = 1-p$, so $E(Y) = p$), $\{x_1, \dots, x_k\}$: predictors (which are scalars).

\bullet sample: $\{(x_{11}, \dots, x_{1k}; y_1), (x_{21}, \dots, x_{2k}; y_2), \dots, (x_{n1}, \dots, x_{nk}; y_n)\}$.

\bullet how to estimate $\beta_j \approx \hat{\beta}_j$: maximal likelihood method (maximize $\prod_{i=1}^n p(x_{i1}, \dots, x_{ik})^{y_i} (1 - p(x_{i1}, \dots, x_{ik}))^{1-y_i}$).

\bullet $\frac{\hat{\beta}_j - \beta_j}{s.e(\hat{\beta}_j)} \approx N(0, 1)$ for large $n \geq 30$, this helps determine whether or not the real $\beta_j = 0$?

\bullet Classification of a new object $Y(x_1, \dots, x_k)$ as 1 or 0 according

$$Y(x_1, \dots, x_k) = \begin{cases} 1, & \text{if } \hat{p}(x_1, \dots, x_k) \geq 0.5, \\ 0, & \text{if } \hat{p}(x_1, \dots, x_k) < 0.5, \end{cases}$$

where the estimated logit function $\hat{p}(x_1, \dots, x_k)$ is

$$\hat{p}(x_1, \dots, x_k) = \frac{e^{\hat{\beta}_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k}}{1 + e^{\hat{\beta}_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k}}.$$

9. Tables

(9.1) Table for $N(0,1)$ standard normal random variable $\Phi(x) = P(N(0,1) \leq x)$, $x \geq 0$.

There is an important relation $\Phi(-x) = 1 - \Phi(x)$, $x \geq 0$.

x	0	1	2	3	4	5	6	7	8	9
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7957	0.8023	0.8051	0.8078	0.8106	0.8133	0.8160
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9222	0.9256	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319	0.9330
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964	0.9965
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9977	0.9978	0.9978	0.9979	0.9979	0.9980	0.9981	0.9981
2.9	0.9981	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986	0.9986
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9990	0.9990	0.9990
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9993	0.9993	0.9993	0.9993
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995	0.9995
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997	0.9997
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998	0.9998
3.5	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998
3.6	0.9998	0.9998	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.7	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.8	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.9	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
4.0	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

(9.2) Table for $T(f)$ random variable $F(x) = P(T(f) \leq x)$, where f is a parameter called 'degrees of freedom'.

f	0.75	0.90	0.95	0.975	0.99	0.995	0.9975	0.9995	$F(x)$
1	1.00	3.08	6.31	12.71	31.82	63.66	127.32	636.62	
2	0.82	1.89	2.92	4.30	6.96	9.92	14.09	31.60	
3	0.76	1.64	2.35	3.18	4.54	5.84	7.45	12.92	
4	0.74	1.53	2.13	2.78	3.75	4.60	5.60	8.61	
5	0.73	1.48	2.02	2.57	3.36	4.03	4.77	6.87	
6	0.72	1.44	2.45	3.14	3.71	4.32	5.96	10.47	
7	0.71	1.41	1.89	2.36	3.00	3.50	4.03	5.41	
8	0.71	1.40	1.86	2.31	2.90	3.36	3.83	5.04	
9	0.70	1.38	1.83	2.26	2.82	3.25	3.69	4.78	
10	0.70	1.37	1.81	2.23	2.76	3.17	3.58	4.59	
11	0.70	1.36	1.80	2.20	2.72	3.11	3.50	4.44	
12	0.70	1.36	1.78	2.18	2.68	3.05	3.43	4.32	
13	0.69	1.35	1.77	2.16	2.65	3.01	3.37	4.22	
14	0.69	1.35	1.76	2.14	2.62	2.98	3.33	4.14	
15	0.69	1.34	1.75	2.13	2.60	2.95	3.29	4.07	
16	0.69	1.34	1.75	2.12	2.58	2.92	3.25	4.01	
17	0.69	1.33	1.74	2.11	2.57	2.90	3.22	3.97	
18	0.69	1.33	1.73	2.10	2.55	2.88	3.20	3.92	
19	0.69	1.33	1.73	2.09	2.54	2.86	3.17	3.88	
20	0.69	1.33	1.72	2.09	2.53	2.85	3.15	3.85	
21	0.69	1.32	1.72	2.08	2.52	2.83	3.14	3.82	
22	0.69	1.32	1.72	2.07	2.51	2.82	3.12	3.79	
23	0.69	1.32	1.71	2.07	2.50	2.81	3.10	3.77	
24	0.68	1.32	1.71	2.06	2.49	2.80	3.09	3.75	
25	0.68	1.32	1.71	2.06	2.49	2.79	3.08	3.73	
26	0.68	1.31	1.71	2.06	2.48	2.78	3.07	3.71	
27	0.68	1.31	1.70	2.05	2.47	2.77	3.06	3.69	
28	0.68	1.31	1.70	2.05	2.47	2.76	3.05	3.67	
29	0.68	1.31	1.70	2.05	2.46	2.76	3.04	3.66	
30	0.68	1.31	1.70	2.04	2.46	2.75	3.03	3.65	
40	0.68	1.30	1.68	2.02	2.42	2.70	2.97	3.55	
50	0.68	1.30	1.68	2.01	2.40	2.68	2.94	3.50	
60	0.68	1.30	1.67	2.00	2.39	2.66	2.91	3.46	
100	0.68	1.29	1.66	2.00	2.36	2.63	2.87	3.39	
∞	0.67	1.28	1.65	1.96	2.33	2.58	2.81	3.29	

(9.3) Table for $\chi^2(f)$ random variable $F(x) = P(\chi^2(f) \leq x)$, where f is a parameter.

f	$F(x)$										
	0.0005	0.001	0.005	0.01	0.025	0.05	0.10	0.20	0.30	0.40	0.50
1	0.00	0.00	0.00	0.00	0.00	0.02	0.06	0.15	0.27	0.45	1
2	0.00	0.00	0.01	0.02	0.05	0.10	0.21	0.45	0.71	1.39	2
3	0.02	0.02	0.07	0.11	0.22	0.35	0.58	1.01	1.42	2.37	3
4	0.06	0.09	0.21	0.30	0.48	0.71	1.06	1.65	2.19	2.75	4
5	0.16	0.21	0.41	0.55	0.83	1.15	1.61	2.34	3.00	3.66	5
6	0.30	0.38	0.68	0.87	1.24	1.64	2.20	3.07	3.83	4.57	6
7	0.48	0.60	0.99	1.24	1.69	2.17	2.83	3.82	4.67	5.49	7
8	0.71	0.86	1.34	1.65	2.18	2.73	3.49	4.59	5.53	6.42	8
9	0.97	1.15	1.73	2.09	2.70	3.33	4.17	5.38	6.39	7.36	9
10	1.26	1.48	2.16	2.56	3.25	3.94	4.87	6.18	7.27	8.30	10
11	1.59	1.83	2.60	3.05	3.82	4.57	5.58	6.99	8.15	9.24	11
12	1.93	2.21	3.07	3.57	4.40	5.23	6.30	7.81	9.03	10.18	12
13	2.31	2.62	3.57	4.11	5.01	5.89	7.04	8.63	9.93	11.13	13
14	2.70	3.04	4.07	4.66	5.63	6.57	7.79	9.47	10.82	12.08	14
15	3.11	3.48	4.60	5.23	6.26	7.26	8.55	10.31	11.72	13.03	15
16	3.54	3.94	5.14	5.81	6.91	7.96	9.31	11.15	12.62	13.98	16
17	3.98	4.42	5.70	6.41	7.56	8.67	10.09	12.00	13.53	14.94	17
18	4.44	4.90	6.26	7.01	8.23	9.39	10.86	12.86	14.44	15.89	18
19	4.91	5.41	6.84	7.63	8.91	10.12	11.65	13.72	15.35	16.85	19
20	5.40	5.92	7.43	8.26	9.59	10.85	12.44	14.58	16.27	17.81	20
21	5.90	6.45	8.03	8.90	10.28	11.59	13.24	15.44	17.18	18.77	21
22	6.40	6.98	8.64	9.54	10.98	12.34	14.04	16.31	18.10	19.73	22
23	6.92	7.53	9.26	10.20	11.69	13.09	14.85	17.19	19.02	20.69	23
24	7.45	8.08	9.89	10.86	12.40	13.85	15.66	18.06	19.94	21.65	24
25	7.99	8.65	10.52	11.52	13.12	14.61	16.47	18.94	20.87	22.62	25
26	8.54	9.22	11.16	12.20	13.84	15.38	17.29	19.82	21.79	23.58	26
27	9.09	9.80	11.81	12.88	14.57	16.15	18.11	20.70	22.72	24.54	27
28	9.66	10.39	12.46	13.56	15.31	16.93	18.94	21.59	23.65	25.51	28
29	10.23	10.99	13.12	14.26	16.05	17.71	19.77	22.48	24.58	26.48	29
30	10.80	11.59	13.79	14.95	16.79	18.49	20.60	23.36	25.51	27.44	30
40	16.91	17.92	20.71	22.16	24.43	26.51	29.05	32.34	34.87	37.13	40
50	23.46	24.67	27.99	29.71	32.36	34.76	37.69	41.45	44.31	46.86	50
60	30.34	31.74	35.53	37.48	40.48	43.19	46.46	50.64	53.81	56.62	60
100	59.90	61.92	67.33	70.06	74.22	77.93	82.36	87.95	92.13	95.81	100

Table for $\chi^2(f)$ random variable $F(x) = P(\chi^2(f) \leq x)$, where f is a parameter.

f	$F(x)$									
	0.60	0.70	0.80	0.90	0.95	0.975	0.99	0.995	0.999	0.9995
1	0.71	1.07	1.64	2.71	3.84	5.02	6.63	7.88	10.83	12.12
2	1.83	2.41	3.22	4.61	5.99	7.38	9.21	10.60	13.82	15.20
3	2.95	3.66	4.64	6.25	7.81	9.35	11.34	12.84	16.27	17.73
4	4.04	4.88	5.99	7.78	9.49	11.14	13.28	14.86	18.47	20.00
5	5.13	6.06	7.29	9.24	11.07	12.83	15.09	16.75	20.52	22.11
6	6.21	7.23	8.56	10.64	12.59	14.45	16.81	18.55	22.46	24.10
7	7.28	8.38	9.80	12.02	14.07	16.01	18.48	20.28	24.32	26.02
8	8.35	9.52	11.03	13.36	15.51	17.53	20.09	21.95	26.12	27.87
9	9.41	10.66	12.24	14.68	16.92	19.02	21.67	23.59	27.88	29.67
10	10.47	11.78	13.44	15.99	18.31	20.48	23.21	25.19	29.59	31.42
11	11.53	12.90	14.63	17.28	19.68	21.92	24.72	26.76	31.14	33.14
12	12.58	14.01	15.81	18.55	21.03	23.34	26.22	28.30	32.91	34.82
13	13.64	15.12	16.98	19.81	22.36	24.74	27.69	29.82	34.53	36.48
14	14.69	16.22	18.15	21.06	23.68	26.12	29.14	31.32	36.12	38.11
15	15.73	17.32	19.31	22.31	25.00	27.49	30.58	32.80	37.70	39.79
16	16.78	18.42	20.47	23.54	26.30	28.85	32.00	34.27	39.25	41.31
17	17.82	19.51	21.61	24.77	27.59	30.19	33.41	35.72	40.79	42.88
18	18.87	20.60	22.76	25.99	28.87	31.53	34.81	37.16	42.31	44.43
19	19.91	21.69	23.90	27.20	30.14	32.85	36.19	38.58	43.82	45.97
20	20.95	22.77	25.04	28.41	31.41	34.17	37.57	40.00	45.31	47.50
21	21.99	23.86	26.17	29.62	32.67	35.48	38.93	41.40	46.80	49.01
22	23.03	24.94	27.30	30.81	33.92	36.78	40.29	42.80	48.27	50.51
23	24.07	26.02	28.43	32.01	35.17	38.08	41.64	44.18	49.73	52.00
24	25.11	27.10	29.55	33.20	36.42	39.36	42.98	45.56	51.18	53.48
25	26.14	28.17	30.68	34.38	37.65	40.65	44.31	46.93	52.62	54.95
26	27.18	29.25	31.79	35.56	38.89	41.92	45.64	48.29	54.05	56.41
27	28.21	30.32	32.91	36.74	40.11	43.19	46.96	49.64	55.48	57.86
28	29.25	31.39	34.03	37.92	41.34	44.46	48.28	50.99	56.89	59.30
29	30.28	32.46	35.14	39.09	42.56	45.72	49.59	52.34	58.30	60.73
30	31.32	33.53	36.25	40.26	43.77	46.98	50.89	53.67	59.70	62.16
40	41.62	44.16	47.27	51.81	55.76	59.34	63.69	66.77	73.40	76.09
50	51.89	54.72	58.16	63.17	67.50	71.42	76.15	79.49	86.66	89.56
60	62.13	65.23	68.97	74.40	79.08	83.30	88.38	91.95	99.61	102.69
100	102.95	106.91	111.67	118.50	124.34	129.56	135.81	140.17	149.45	153.17

(9.4) Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
 If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1-p) \geq n-k)$.

n	k	0.05	0.10	0.15	0.20	0.25	0.30	0.35	0.40	0.45	0.50	p
2	0	0.9025	0.8100	0.7225	0.6400	0.5625	0.4900	0.4225	0.3600	0.3025	0.2500	
3	0	0.9575	0.9900	0.9775	0.9600	0.9375	0.9100	0.8775	0.8400	0.7975	0.7500	
4	0	0.8574	0.7290	0.6141	0.5120	0.4219	0.3430	0.2746	0.2160	0.1664	0.1250	
5	0	0.9928	0.9720	0.9392	0.8960	0.8438	0.7840	0.7183	0.6480	0.5747	0.5000	
6	0	0.9999	0.9990	0.9966	0.9920	0.9844	0.9730	0.9571	0.9360	0.9089	0.8750	
7	0	0.8145	0.6561	0.5220	0.4096	0.3164	0.2401	0.1785	0.1296	0.0915	0.0625	
8	0	0.9860	0.9477	0.8905	0.8192	0.7383	0.6517	0.5630	0.4752	0.3910	0.3125	
9	0	0.9995	0.9963	0.9880	0.9728	0.9492	0.9163	0.8735	0.8208	0.7585	0.6875	
10	0	0.7738	0.5905	0.4437	0.3277	0.2373	0.1681	0.1160	0.0778	0.0503	0.0313	
11	0	0.9774	0.9185	0.8352	0.7373	0.6328	0.5282	0.4284	0.3370	0.2562	0.1875	
12	0	0.9988	0.9914	0.9734	0.9421	0.8965	0.8369	0.7648	0.6826	0.5931	0.5000	
13	0	0.9999	0.9995	0.9978	0.9933	0.9844	0.9692	0.9460	0.9130	0.8688	0.8125	
14	0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
15	0	0.7351	0.5314	0.3771	0.2621	0.1780	0.1176	0.0754	0.0467	0.0277	0.0156	
16	0	0.9672	0.8857	0.7765	0.6554	0.5339	0.4202	0.3191	0.2333	0.1636	0.1094	
17	0	0.9978	0.9842	0.9527	0.9011	0.8306	0.7443	0.6471	0.5443	0.4415	0.3438	
18	0	0.9999	0.9987	0.9941	0.9830	0.9624	0.9295	0.8826	0.8208	0.7447	0.6563	
19	0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
20	0	0.6983	0.4783	0.3206	0.2097	0.1335	0.0824	0.0490	0.0280	0.0152	0.0078	
21	0	0.9556	0.8503	0.7166	0.5767	0.4449	0.3294	0.2338	0.1586	0.1024	0.0625	
22	0	0.9962	0.9743	0.9262	0.8520	0.7564	0.6471	0.5323	0.4199	0.3124	0.2266	
23	0	0.9998	0.9973	0.9879	0.9667	0.9294	0.8740	0.8002	0.7102	0.6083	0.5000	
24	0	1.0000	0.9998	0.9988	0.9953	0.9871	0.9712	0.9444	0.9037	0.8471	0.7734	
25	0	1.0000	1.0000	0.9999	0.9996	0.9987	0.9962	0.9910	0.9812	0.9643	0.9375	
26	0	1.0000	1.0000	1.0000	0.9999	0.9998	0.9994	0.9984	0.9963	0.9922	0.9900	
27	0	0.6634	0.4305	0.2725	0.1678	0.1001	0.0576	0.0319	0.0168	0.0084	0.0039	
28	0	1.0000	0.8131	0.6572	0.5033	0.3671	0.2553	0.1691	0.1064	0.0632	0.0352	
29	0	2.0000	0.9619	0.8948	0.7969	0.6785	0.5518	0.4278	0.3154	0.2201	0.1445	
30	0	3.0000	0.9950	0.9786	0.8862	0.8059	0.7064	0.5941	0.4770	0.3633	0.2700	
31	0	4.0000	0.9996	0.9971	0.9896	0.9727	0.9420	0.8939	0.8263	0.7396	0.6367	
32	0	5.0000	1.0000	0.9998	0.9988	0.9887	0.9747	0.9502	0.9115	0.8555	0.7700	
33	0	6.0000	1.0000	1.0000	0.9999	0.9987	0.9964	0.9915	0.9819	0.9643	0.9300	
34	0	7.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9993	0.9983	0.9961	0.9900	
35	0	8.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
36	0	9.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
37	0	10.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
38	0	11.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
39	0	12.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	

Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
 If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1-p) \geq n-k)$.

n	k	0.05	0.10	0.15	0.20	0.25	0.30	0.35	0.40	0.45	0.50	p
11	0	0.5688	0.3138	0.1673	0.0859	0.0422	0.0198	0.0088	0.0036	0.0014	0.0005	
12	0	0.5404	0.2824	0.1422	0.0687	0.0317	0.0138	0.0057	0.0022	0.0008	0.0002	
13	0	0.5040	0.2242	0.1087	0.0544	0.0274	0.0121	0.0055	0.0023	0.0009	0.0003	
14	0	0.4816	0.1659	0.0843	0.0443	0.0274	0.0158	0.0085	0.0042	0.0016	0.0003	
15	0	0.4552	0.1174	0.0574	0.0317	0.0197	0.0102	0.0053	0.0026	0.0011	0.0003	
16	0	0.4256	0.0963	0.0496	0.0274	0.0158	0.0085	0.0043	0.0021	0.0009	0.0003	
17	0	0.3971	0.0778	0.0389	0.0214	0.0121	0.0063	0.0033	0.0016	0.0007	0.0003	
18	0	0.3744	0.0681	0.0319	0.0197	0.0102	0.0053	0.0026	0.0011	0.0005	0.0003	
19	0	0.3532	0.0588	0.0287	0.0142	0.0071	0.0036	0.0018	0.0008	0.0004	0.0003	
20	0	0.3345	0.0496	0.0214	0.0102	0.0053	0.0026	0.0011	0.0005	0.0003	0.0002	
21	0	0.3125	0.0404	0.0171	0.0085	0.0042	0.0021	0.0010	0.0005	0.0003	0.0002	
22	0	0.2960	0.0319	0.0142	0.0063	0.0033	0.0016	0.0008	0.0004	0.0003	0.0002	
23	0	0.2823	0.0287	0.0114	0.0053	0.0026	0.0011	0.0005	0.0003	0.0002	0.0002	
24	0	0.2660	0.0214	0.0085	0.0042	0.0021	0.0010	0.0005	0.0003	0.0002	0.0002	
25	0	0.2500	0.0197	0.0071	0.0036	0.0016	0.0008	0.0004	0.0003	0.0002	0.0002	
26	0	0.2379	0.0171	0.0063	0.0033	0.0016	0.0008	0.0004	0.0003	0.0002	0.0002	
27	0	0.2279	0.0142	0.0053	0.0026	0.0011	0.0005	0.0003	0.0002	0.0002	0.0002	
28	0	0.2134	0.0114	0.0042	0.0021	0.0010	0.0005	0.0003	0.0002	0.0002	0.0002	
29	0	0.2025	0.0102	0.0042	0.0021	0.0010	0.0005	0.0003	0.0002	0.0002	0.0002	
30	0	0.1970	0.0091	0.0036	0.0016	0.0008	0.0004	0.0003	0.0002	0.0002	0.0002	
31	0	0.1918	0.0081	0.0033	0.0014	0.0007	0.0004	0.0003	0.0002	0.0002	0.0002	
32	0	0.1874	0.0071	0.0031	0.0013	0.0006	0.0004	0.0003	0.0002	0.0002	0.0002	
33	0	0.1836	0.0061	0.0029	0.0012	0.0005	0.0003	0.0003	0.0002	0.0002	0.0002	
34	0	0.1790	0.0051	0.0028	0.0011	0.0005	0.0003	0.0003	0.0002	0.0002	0.0002	
35	0	0.1746	0.0041	0.0027	0.0010	0.0005	0.0003	0.0003	0.0002	0.0002	0.0002	
36	0	0.1709	0.0031	0.0026	0.0010	0.0005	0.0003	0.0003	0.0002	0.0002	0.0002	
37	0	0.1674	0.0021	0.0025	0.0009	0.0005	0.0003	0.0003	0.0002	0.0002	0.0002	
38	0	0.1640	0.0011	0.0024	0.0008	0.0005	0.0003	0.0003	0.0002	0.0002	0.0002	
39	0	0.1606	0.0009	0.0023	0.0007	0.0005	0.0003	0.0003	0.0002	0.0002	0.0002	

Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1-p) \geq n-k)$.

n	k	0.05	0.10	0.15	0.20	0.25	p	0.30	0.35	0.40	0.45	0.50
14	0	0.4877	0.2288	0.1028	0.0440	0.0178	0.0068	0.0024	0.0008	0.0002	0.0001	
1	0.8470	0.5284	0.3567	0.1979	0.1010	0.0475	0.0205	0.0081	0.0009			
2	0.9699	0.8416	0.6479	0.4481	0.2811	0.1608	0.0839	0.0398	0.0170	0.0065		
3	0.9558	0.9559	0.8535	0.6982	0.5213	0.3552	0.2205	0.1243	0.0632	0.0287	0.0123	
4	0.9996	0.9908	0.9533	0.8702	0.7415	0.5842	0.4227	0.2793	0.1672	0.0898	0.0464	
5	1.0000	0.9985	0.9885	0.9561	0.8883	0.7805	0.6405	0.4859	0.3373	0.2120	0.1260	0.0596
6	1.0000	0.9998	0.9978	0.9884	0.9617	0.9067	0.8164	0.6925	0.5461	0.3953	0.2348	0.0245
7	1.0000	1.0000	0.9997	0.9976	0.9897	0.9685	0.9247	0.8499	0.7414	0.6047	0.4478	0.2902
8	1.0000	1.0000	1.0000	0.9996	0.9978	0.9917	0.9757	0.8811	0.7830	0.6405	0.4743	0.3145
9	1.0000	1.0000	1.0000	1.0000	0.9997	0.9983	0.9940	0.9825	0.9574	0.9102	0.8011	0.6626
10	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9989	0.9961	0.9886	0.9713	0.9177	0.6855
11	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9978	0.9935	0.9777	0.8338
12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9894	0.9283
13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9755
14	0	0.4633	0.2059	0.0874	0.0352	0.0134	0.0047	0.0016	0.0005	0.0001	0.0000	
1	0.8290	0.5490	0.3186	0.1671	0.0802	0.0353	0.0142	0.0052	0.0017	0.0005		
2	0.9638	0.8159	0.6042	0.3980	0.2361	0.1268	0.0617	0.0271	0.0107	0.0037		
3	0.9445	0.9444	0.8227	0.6482	0.4613	0.2969	0.1727	0.0905	0.0424	0.0176		
4	0.9994	0.9873	0.9383	0.8358	0.6865	0.5155	0.3519	0.2173	0.1204	0.0592		
5	0.9999	0.9978	0.9832	0.9389	0.8516	0.7216	0.5643	0.4032	0.2608	0.1509		
6	1.0000	0.9997	0.9964	0.9819	0.9434	0.8689	0.7548	0.6098	0.4522	0.3036		
7	1.0000	1.0000	0.9994	0.9958	0.9827	0.9500	0.8868	0.7869	0.6535	0.5000		
8	1.0000	1.0000	0.9999	0.9992	0.9958	0.9848	0.9578	0.9050	0.8182	0.6964		
9	1.0000	1.0000	1.0000	0.9990	0.9992	0.9963	0.9876	0.9662	0.9231	0.8491		
10	1.0000	1.0000	1.0000	0.9999	0.9993	0.9972	0.9907	0.9745	0.9408	0.8737		
11	1.0000	1.0000	1.0000	1.0000	0.9999	0.9995	0.9981	0.9937	0.9824	0.8807		
12	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9989	0.9963	0.9790		
13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9995	0.9939		
14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
15	0	0.4401	0.1853	0.0743	0.0281	0.0100	0.0033	0.0010	0.0003	0.0001	0.0000	
1	0.8108	0.5147	0.2839	0.1407	0.0635	0.0261	0.0098	0.0033	0.0010	0.0003		
2	0.9571	0.7892	0.5614	0.3518	0.1971	0.0994	0.0451	0.0183	0.0066	0.0021		
3	0.9360	0.9316	0.7899	0.5981	0.4050	0.2459	0.1339	0.0651	0.0281	0.0106		
4	0.9991	0.9830	0.9209	0.7982	0.6302	0.4499	0.2892	0.1666	0.0853	0.0384		
5	0.9999	0.9967	0.9765	0.9183	0.8103	0.6598	0.4900	0.3288	0.1976	0.1051		
6	1.0000	0.9995	0.9944	0.9733	0.9204	0.8247	0.6881	0.5272	0.3660	0.2272		
7	1.0000	0.9999	0.9930	0.9729	0.9256	0.8406	0.7161	0.5629	0.4018	0.2409		
8	1.0000	1.0000	0.9998	0.9985	0.9795	0.9743	0.9329	0.8579	0.7441	0.5982		
9	1.0000	1.0000	1.0000	0.9998	0.9984	0.9929	0.9771	0.9417	0.8759	0.7728		
10	1.0000	1.0000	1.0000	0.9997	0.9984	0.9938	0.9809	0.9514	0.8949	0.8287		
11	1.0000	1.0000	1.0000	1.0000	0.9997	0.9987	0.9951	0.9851	0.9616	0.9216		
12	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9991	0.9965	0.9894	0.9592		
13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9979	0.9979	0.9913		
14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9997		
15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		

Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1-p) \geq n-k)$.

n	k	0.05	0.10	0.15	0.20	0.25	p	0.30	0.35	0.40	0.45	0.50
17	0	0.4181	0.1668	0.0631	0.0225	0.0075	0.0023	0.0007	0.0002	0.0000	0.0000	
1	0.7922	0.4818	0.2525	0.1182	0.0501	0.0193	0.0067	0.0021	0.0006	0.0001		
2	0.9497	0.7618	0.5198	0.3096	0.1637	0.0774	0.0327	0.0123	0.0041	0.0012		
3	0.9912	0.9174	0.7556	0.5489	0.3530	0.2019	0.1028	0.0464	0.0184	0.0064		
4	0.9988	0.9779	0.9013	0.7582	0.5739	0.3887	0.2348	0.1260	0.0596	0.0245		
5	0.9999	0.9953	0.9681	0.8943	0.7653	0.5968	0.4197	0.2639	0.1471	0.0717		
6	1.0000	0.9992	0.9623	0.8929	0.7752	0.6188	0.4478	0.2902	0.1662			
7	1.0000	0.9999	0.9883	0.9598	0.8787	0.7872	0.6405	0.4743	0.3145			
8	1.0000	1.0000	0.9997	0.9974	0.9876	0.9597	0.9006	0.8011	0.6626	0.5000		
9	1.0000	1.0000	1.0000	0.9999	0.9999	0.9970	0.9894	0.9699	0.9283	0.7597		
10	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
11	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
16	0	0.3774	0.1351	0.0456	0.0144	0.0042	0.0011	0.0003	0.0001	0.0000		
1	0.7547	0.4203	0.1985	0.0829	0.0310	0.0104	0.0031	0.0008	0.0002	0.0000		
2	0.9335	0.7054	0.4413	0.2369	0.1113	0.0462	0.0170	0.0055	0.0015	0.0004		
3	0.9836	0.8850	0.6841	0.4551	0.2631	0.1332	0.0591	0.0230	0.0077	0.0022		
4	0.9980	0.9648	0.8556	0.6733	0.4654	0.2822	0.1500	0.0696	0.0280	0.0096		
5	0.9998	0.9914	0.9463	0.8369	0.6678	0.4739	0.2968	0.1629	0.0777	0.0318		
6	1.0000	0.9983	0.9837	0.9324	0.8251	0.6655	0.4812	0.3081	0.1727	0.0835		
7	1.0000	0.9997	0.9959	0.9767	0.9225	0.8180	0.6656	0.4878	0.3169	0.1796		
8	1.0000	1.0000	0.9992	0.9933	0.9713	0.9161	0.8145	0.6675	0.4940	0.3238		
9	1.0000	1.0000	0.9999	0.9974	0.9817	0.9077	0.8159	0.6710	0.4915	0.3238		
10	1.0000	1.0000	1.0000	0.9997	0.9977	0.9895	0.9653	0.9115	0.8159	0.6762		
11	1.0000	1.0000	1.0000	0.9995	0.9972	0.9886	0.9648	0.9129	0.8204	0.6822		
12	1.0000	1.0000	1.0000	0.9999	0.9994	0.9969	0.9884	0.9658	0.9165	0.7914		
13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
16	0	0.3774	0.1351	0.0456	0.0144	0.0042	0.0011	0.0003	0.0001	0.0000		

(9.5) Table for Poisson random variable $P(Po(\mu) \leq k)$.

k	μ									
k	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0
0	0.9048	0.8187	0.7408	0.6703	0.6065	0.5488	0.4966	0.4493	0.4066	0.3679
1	0.9953	0.9825	0.9631	0.9384	0.9098	0.8781	0.8442	0.8088	0.7725	0.7358
2	0.9998	0.9989	0.9964	0.9921	0.9856	0.9769	0.9659	0.9526	0.9371	0.9197
3	1.0000	0.9999	0.9997	0.9992	0.9982	0.9966	0.9942	0.9909	0.9865	0.9810
4	1.0000	1.0000	0.9999	0.9998	0.9996	0.9992	0.9986	0.9977	0.9963	0.9956
5	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9994
6	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
7	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
k	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0
0	0.3329	0.3012	0.2725	0.2466	0.2231	0.2019	0.1827	0.1653	0.1496	0.1353
1	0.6890	0.6626	0.6268	0.5918	0.5578	0.5249	0.4932	0.4628	0.4337	0.4060
2	0.904	0.8795	0.8571	0.8335	0.8088	0.7834	0.7572	0.7306	0.7037	0.6767
3	0.9743	0.9662	0.9569	0.9463	0.9344	0.9212	0.9068	0.8913	0.8747	0.8571
4	0.9946	0.9923	0.9893	0.9857	0.9814	0.9763	0.9704	0.9636	0.9559	0.9473
5	0.9990	0.9985	0.9978	0.9968	0.9955	0.9940	0.9920	0.9896	0.9834	0.9775
6	0.9999	0.9997	0.9996	0.9994	0.9991	0.9987	0.9981	0.9974	0.9966	0.9955
7	1.0000	1.0000	0.9999	0.9999	0.9998	0.9997	0.9996	0.9994	0.9992	0.9989
8	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
9	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
k	3.2	3.4	3.6	3.8	4.0	4.2	4.4	4.6	4.8	5.0
0	0.0408	0.0334	0.0273	0.0224	0.0183	0.0150	0.0123	0.0101	0.0082	0.0067
1	0.1712	0.1468	0.1257	0.1074	0.0916	0.0780	0.0663	0.0563	0.0477	0.0404
2	0.3799	0.3397	0.3027	0.2689	0.2381	0.2102	0.1851	0.1626	0.1425	0.1247
3	0.6025	0.5584	0.5152	0.4735	0.4335	0.3954	0.3594	0.3257	0.2942	0.2650
4	0.7806	0.7442	0.7064	0.6678	0.6288	0.5898	0.5512	0.5132	0.4763	0.4405
5	0.8946	0.8705	0.8441	0.8156	0.7851	0.7531	0.7199	0.6858	0.6510	0.6160
6	0.9554	0.9421	0.9267	0.9091	0.8893	0.8675	0.8436	0.8180	0.7908	0.7622
7	0.9832	0.9769	0.9692	0.9559	0.9489	0.9361	0.9214	0.9049	0.8867	0.8666
8	0.9943	0.9917	0.9883	0.9840	0.9786	0.9721	0.9642	0.9549	0.9442	0.9319
9	0.9995	0.9992	0.9987	0.9981	0.9972	0.9959	0.9943	0.9922	0.9896	0.9863
k	5.2	5.4	5.6	5.8	6.0	6.5	7.0	7.5	8.0	8.5
0	0.0055	0.0045	0.0037	0.0030	0.0025	0.0015	0.0009	0.0006	0.0003	0.0002
1	0.0342	0.0289	0.0244	0.0206	0.0174	0.0113	0.0073	0.0047	0.0030	0.0019
2	0.1088	0.0948	0.0824	0.0715	0.0620	0.0430	0.0296	0.0203	0.0138	0.0093
3	0.2381	0.2133	0.1906	0.1700	0.1512	0.1118	0.0818	0.0591	0.0424	0.0301
4	0.4061	0.3733	0.3422	0.3127	0.2851	0.2237	0.1730	0.1321	0.0966	0.0744
5	0.5809	0.5461	0.5119	0.4783	0.4457	0.3690	0.3007	0.2414	0.1912	0.1496
6	0.7324	0.7017	0.6703	0.6384	0.6063	0.5265	0.4497	0.3782	0.3134	0.2562
7	0.8449	0.8149	0.7970	0.7710	0.7440	0.6728	0.5987	0.5246	0.4530	0.3856
8	0.9181	0.9027	0.8857	0.8672	0.8472	0.7916	0.7291	0.6620	0.5925	0.5231
9	0.9603	0.9512	0.9409	0.9292	0.9161	0.8774	0.8305	0.7764	0.7166	0.6530
10	0.9823	0.9775	0.9718	0.9651	0.9574	0.9332	0.9015	0.8622	0.8159	0.7634
11	0.9904	0.9875	0.9841	0.9799	0.9661	0.9467	0.9208	0.8881	0.8487	0.8047
12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
k	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0
0	0.1225	0.1108	0.1003	0.0907	0.0821	0.0743	0.0672	0.0608	0.0550	0.0498
1	0.3796	0.3546	0.3309	0.3084	0.2873	0.2674	0.2487	0.2311	0.2146	0.1991
2	0.6496	0.6227	0.5960	0.5597	0.5348	0.5184	0.4936	0.4695	0.4460	0.4232
3	0.8386	0.8194	0.7787	0.7576	0.7360	0.7141	0.6919	0.6696	0.6474	0.6141
4	0.9379	0.9275	0.9162	0.9041	0.8912	0.8774	0.8477	0.8318	0.8153	0.7894
5	0.9796	0.9751	0.9700	0.9643	0.9580	0.9510	0.9433	0.9349	0.9258	0.9161
6	0.9941	0.9925	0.9884	0.9828	0.9794	0.9756	0.9713	0.9665	0.9623	0.9573
7	0.9985	0.9974	0.9967	0.9958	0.9947	0.9934	0.9919	0.9901	0.9881	0.9851
8	0.9997	0.9995	0.9994	0.9991	0.9989	0.9985	0.9981	0.9976	0.9969	0.9961
9	0.9999	0.9999	0.9999	0.9998	0.9997	0.9996	0.9995	0.9994	0.9993	0.9992
10	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
11	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

Table for Poisson random variable $P(Po(\mu) \leq k)$.

k	9.0	9.5	10.0	11.0	12.0	13.0	14.0	15.0	16.0	17.0	μ
0	0.0001	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
1	0.0012	0.0008	0.0005	0.0002	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
2	0.0062	0.0042	0.0028	0.0012	0.0005	0.0002	0.0001	0.0000	0.0000	0.0000	0.0000
3	0.0212	0.0149	0.0103	0.0049	0.0023	0.0011	0.0005	0.0002	0.0001	0.0000	0.0000
4	0.0550	0.0403	0.0293	0.0151	0.0076	0.0037	0.0018	0.0009	0.0004	0.0002	0.0000
5	0.1157	0.0885	0.0671	0.0375	0.0203	0.0107	0.0055	0.0028	0.0014	0.0007	0.0000
6	0.2068	0.1649	0.1301	0.0786	0.0458	0.0259	0.0142	0.0076	0.0040	0.0021	0.0000
7	0.3239	0.2687	0.2202	0.1432	0.0895	0.0540	0.0316	0.0180	0.0100	0.0054	0.0000
8	0.4557	0.3918	0.3328	0.2320	0.1550	0.0998	0.0621	0.0374	0.0220	0.0126	0.0000
9	0.5874	0.5218	0.4579	0.3405	0.2424	0.1658	0.1094	0.0699	0.0433	0.0261	0.0000
10	0.7060	0.6453	0.5830	0.4599	0.3472	0.2517	0.1757	0.1185	0.0774	0.0491	0.0000
11	0.8030	0.7520	0.6968	0.5793	0.4616	0.3532	0.2600	0.1848	0.1270	0.0847	0.0000
12	0.8758	0.8364	0.7916	0.6887	0.5760	0.4631	0.3585	0.2676	0.1931	0.1350	0.0000
13	0.9261	0.8981	0.8645	0.7813	0.6815	0.5730	0.4644	0.3632	0.2745	0.2009	0.0000
14	0.9585	0.9400	0.9165	0.8540	0.7720	0.6751	0.5704	0.4657	0.3675	0.2808	0.0000
15	0.9780	0.9665	0.9513	0.9074	0.8444	0.7636	0.6694	0.5681	0.4667	0.3715	0.0000
16	0.9889	0.9823	0.9730	0.9441	0.8987	0.8355	0.7559	0.6641	0.5660	0.4677	0.0000
17	0.9947	0.9911	0.9857	0.9678	0.9370	0.8905	0.8272	0.7489	0.6593	0.5640	0.0000
18	0.9976	0.9957	0.9928	0.9823	0.9626	0.9302	0.8826	0.8195	0.7423	0.6550	0.0000
19	0.9989	0.9980	0.9965	0.9907	0.9787	0.9573	0.9235	0.8752	0.8122	0.7363	0.0000
20	0.9996	0.9991	0.9984	0.9953	0.9884	0.9750	0.9521	0.9170	0.8682	0.8055	0.0000
21	0.9998	0.9996	0.9993	0.9977	0.9939	0.9859	0.9712	0.9469	0.9108	0.8615	0.0000
22	0.9999	0.9999	0.9997	0.9990	0.9970	0.9924	0.9833	0.9673	0.9418	0.9047	0.0000
23	1.0000	0.9999	0.9999	0.9995	0.9985	0.9960	0.9907	0.9805	0.9633	0.9367	0.0000
24	1.0000	1.0000	1.0000	0.9998	0.9993	0.9980	0.9950	0.9888	0.9777	0.9594	0.0000
25	1.0000	1.0000	1.0000	0.9999	0.9997	0.9974	0.9938	0.9869	0.9748	0.9548	0.0000
26	1.0000	1.0000	1.0000	1.0000	0.9999	0.9995	0.9987	0.9967	0.9925	0.9848	0.0000
27	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9994	0.9983	0.9959	0.9912	0.0000
28	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9991	0.9978	0.9950	0.9900	0.0000
29	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9996	0.9989	0.9973	0.9947	0.0000
30	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9994	0.9986	0.9976	0.0000
31	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9993	0.9980	0.0000
32	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9996	0.0000
33	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.0000
34	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.0000
35	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.0000