

Examiner: Xiangfeng Yang (Tel: 013 28 57 88).

Things allowed (Hjälpmedel): a calculator.

Scores rating (Betygsgränser): 8-11 points giving rate 3; 11.5-14.5 points giving rate 4; 15-18 points giving rate 5.

1 (3 points)

Sara is studying at LiU in \textcircled{S} -program. There are 30 students in \textcircled{S} -program: 10 female and 20 male. Now 5 students are randomly chosen for some mission.

(1.1) (1p) Find the probability that Sara will be chosen.

(1.2) (1p) Find the probability that there will be 5 female.

(1.3) (1p) Find the probability that there will be 3 female and 2 male.

Solution. (1.1)

$$P(\text{Sara}) = \frac{1 \cdot \binom{29}{4}}{\binom{30}{5}} = 0.16667.$$

(1.2)

$$P(5 \text{ female}) = \frac{\binom{10}{5}}{\binom{30}{5}} = 0.001768.$$

(1.3)

$$P(3 \text{ female and 2 male}) = \frac{\binom{10}{3} \cdot \binom{20}{2}}{\binom{30}{5}} = 0.15999.$$

□

2 (3 points)

A two dimensional random variable (X, Y) has a joint probability mass function as follows

$X \backslash Y$	6	8
1	0.12	$a = ?$
2	0.28	$b = ?$

The table tells that X can take values 1 and 2, and Y can take values 6 and 8.

(2.1) (1p) Assume that $a = 0.10$ and $b = 0.50$, find the conditional probability $P(X = 2 | Y = 6)$.

(2.2) (1p) Assume that $a = 0.10$ and $b = 0.50$, find the mean $E(X)$ of X .

(2.3) (1p) Find the values of a and b so that X and Y are independent.

Solution. (2.1)

$$P(X = 2 | Y = 6) = \frac{P(X = 2, Y = 6)}{P(Y = 6)} = \frac{0.28}{0.12 + 0.28} = \frac{0.28}{0.40} = 0.70.$$

(2.2) One gets

$$P(X = 1) = 0.12 + 0.10 = 0.22, \quad P(X = 2) = 0.28 + 0.50 = 0.78.$$

Therefore,

$$E(X) = 1 \cdot 0.22 + 2 \cdot 0.78 = 1.78.$$

(2.3) In order that X and Y are independent, we should have $p(x, y) = p_X(x) \cdot p_Y(y)$. Therefore,

$$0.12 = p(1, 6) = p_X(1) \cdot p_Y(y) = (0.12 + a) \cdot (0.12 + 0.28) = (0.12 + a) \cdot 0.4 \implies (0.12 + a) = 0.3 \implies a = 0.18.$$

Then

$$0.12 + 0.28 + a + b = 1 \implies b = 1 - 0.12 - 0.28 - a = 0.42.$$

□

3 (3 points)

Lars is very good at statistics, and he is now in a challenge to solve 100 problems in statistics. According to his past experience, the time X (in minutes) needed for Lars to solve one problem in statistics is a continuous random variable with a probability density function $f(x) = 2x$ for $0 < x < 1$.

(3.1) (1p) Find the mean $\mu = E(X)$ and variance $\sigma^2 = V(X)$ of X .

(3.2) (2p) Find the probability that Lars can solve all the 100 problems within 60 minutes.

Solution. (3.1)

$$\mu = E(X) = \int_0^1 x \cdot f(x) dx = \int_0^1 2x^2 dx = 2/3 = 0.6667,$$

$$\sigma^2 = V(X) = E(X^2) - \mu^2 = \int_0^1 x^2 \cdot f(x) dx - \mu^2 = \int_0^1 2x^3 dx - \mu^2 = 1/2 - 4/9 = 1/18 = 0.05556.$$

(3.2) Let X_1, X_2, \dots, X_{100} denote the times Lars spends on these 100 problems, it is then from CLT that

$$P(\text{Lars can solve all the 100 problems within 60 minutes}) = P(X_1 + X_2 + \dots + X_{100} \leq 60)$$

$$= P(\bar{X} \leq 0.6) = P\left(\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \leq \frac{0.6 - \mu}{\sigma/\sqrt{n}}\right)$$

$$= P(N(0, 1) \leq \frac{0.6 - 0.6667}{\sqrt{0.05556/\sqrt{100}}}) = P(N(0, 1) \leq -2.83) = \Phi(-2.83) = 1 - \Phi(2.83) = 1 - 0.9977 = 0.0023.$$

□

4 (3 points)

A population X is a continuous random variable with a probability density function $f(x) = 1/\theta$ for $0 < x \leq \theta$ where $\theta > 0$ is an unknown parameter. A sample $\{5, 3, 8, 2\}$ is taken from this population.

(4.1) (1p) Use the method of moments to find a point estimate $\hat{\theta}_{MM}$ of θ .

(4.2) (2p) Use the maximum-likelihood method to find a point estimate $\hat{\theta}_{ML}$ of θ .

Solution. (4.1) The mean $E(X)$ of the population can be computed as

$$E(X) = \int_0^\theta x f(x) dx = \int_0^\theta x/\theta dx = \theta/2.$$

It is then from the equation $E(X) = \bar{x}$ that

$$\theta/2 = \bar{x} \implies \hat{\theta}_{MM} = 2 \cdot \bar{x} = 2 \cdot \frac{5 + 3 + 8 + 2}{4} = 2 \cdot 4.5 = 9$$

(4.2) The likelihood function is

$$L(\theta) = f(x_1) \cdot f(x_2) \cdot \dots \cdot f(x_n) = 1/\theta^n, \quad \text{for } \theta \geq \max\{x_1, x_2, \dots, x_n\}.$$

The non-trivial domain is $\theta \geq \max\{x_1, x_2, \dots, x_n\}$ because for each function $f(x_i)$, the domain is $\theta \geq x_i$. It is from $L'(\theta) = -n/\theta^{n+1} < 0$ that the function $L(\theta)$ is decreasing for all $\theta \geq \max\{x_1, x_2, \dots, x_n\}$. This suggests that the maximum of $L(\theta)$ is reached when $\theta = \max\{x_1, x_2, \dots, x_n\}$. Therefore

$$\hat{\theta}_{ML} = \max\{x_1, x_2, \dots, x_n\} = \max\{5, 3, 8, 2\} = 8.$$

□

5 (3 points)

Suppose that the distribution X of weights (in grams) of all peanuts boxes produced by a factory is $X \sim N(\mu, 2^2)$. Now 36 peanuts boxes are randomly selected with sample mean $\bar{x} = 498$ g and sample standard deviation $s = 2.2$ g.

(5.1) (1.5p) Does this sample provide any evidence that the average weigh μ of all such peanuts boxes is less than 500 g? Answer this using appropriate one-sided 95% confidence interval.

(5.2) (1.5p) Does this sample provide any evidence that the average weigh μ of all such peanuts boxes is more than 490 g? Answer this using appropriate one-sided 95% confidence interval.

Solution. (5.1) The one-sided 95% confidence interval is

$$I_\mu = (-\infty, \bar{x} + z_\alpha \cdot \frac{\sigma}{\sqrt{n}}) = (-\infty, 498 + 1.64 \cdot \frac{2}{\sqrt{36}}) = (-\infty, 498.5467).$$

As $498.5467 < 500$, yes, the samples provide evidence showing that $\mu < 500$.

(5.2) The one-sided 95% confidence interval is

$$I_\mu = (\bar{x} - z_\alpha \cdot \frac{\sigma}{\sqrt{n}}, \infty) = (498 - 1.64 \cdot \frac{2}{\sqrt{36}}, \infty) = (497.4533, \infty).$$

As $497.4533 > 490$, yes, the samples provide evidence showing that $\mu > 490$. □

6 (3 points)

Assume that two populations $X \sim N(\mu_X, \sigma^2)$ and $Y \sim N(\mu_Y, \sigma^2)$ are independent. A sample from X gives: $n_1 = 10, \bar{x} = 5.2$ and $s_X = 1.6$. A sample from Y gives: $n_2 = 18, \bar{y} = 3$ and $s_Y = 1.2$.

(6.1) (1.5p) With a significance level 5%, test the hypotheses $H_0 : \mu_X - \mu_Y = 2$ against $H_a : \mu_X - \mu_Y > 2$.

(6.2) (1.5p) With a significance level 5%, test the hypotheses $H_0 : \sigma^2 = 1$ against $H_a : \sigma^2 > 1$.

Solution. (6.1) The test statistic can be chosen as

$$\frac{(\bar{X} - \bar{Y}) - (\mu_X - \mu_Y)}{s \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} \sim T(n_1 + n_2 - 2),$$

where $s^2 = \frac{(n_1-1)s_X^2 + (n_2-1)s_Y^2}{n_1+n_2-2} = 1.828$ (hence $s = \sqrt{s^2} = 1.352$). Therefore,

$$TS = \frac{(\bar{x} - \bar{y}) - 2}{s \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} = \frac{(5.2 - 3) - 2}{1.352 \sqrt{\frac{1}{10} + \frac{1}{18}}} = 0.3751.$$

The rejection region is

$$C = (t_\alpha(n_1 + n_2 - 2), \infty) = (t_{0.05}(10 + 18 - 2), \infty) = (1.71, \infty).$$

Since $TS \notin C$, we don't reject H_0 .

(6.2) The test statistic can be chosen as

$$\frac{(n_1 + n_2 - 2)S^2}{\sigma_0^2} \sim \chi^2(n_1 + n_2 - 2).$$

Therefore,

$$TS = \frac{(n_1 + n_2 - 2)s^2}{\sigma_0^2} = \frac{(10 + 18 - 2) \cdot 1.828}{1^2} = 47.528.$$

The rejection region is

$$C = (\chi_\alpha^2(n_1 + n_2 - 2), \infty) = (\chi_{0.05}^2(10 + 18 - 2), \infty) = (38.89, \infty).$$

Since $TS \in C$, we reject H_0 . □

1. Basic probability

- (1.1) Conditional probability $P(A|B) = \frac{P(A \cap B)}{P(B)}$.
 (1.2) Total probability $P(B) = \sum_{i=1}^k P(B|A_i)P(A_i)$ where $\{A_i\}$ are disjoint and $\cup_{i=1}^k A_i = S$.
 (1.3) Bayes' Theorem $P(A_i|B) = \frac{P(B|A_i)P(A_i)}{\sum_{i=1}^k P(B|A_i)P(A_i)}$ where $\{A_i\}$ are in (1.2).

2. Random variables (r.v.s)

- (2.1) Discrete r.v. X has a pmf $p(x) = P(X = x)$ satisfying $p(x) \geq 0$ and $\sum p(x_i) = 1$,

$$\begin{array}{c|cccc} X & x_1 & x_2 & \cdots & x_n & \cdots \\ \hline p(x) & p(x_1) & p(x_2) & \cdots & p(x_n) & \cdots \end{array}$$

- Expectation (or *Expected value* or *mean*) $\mu_X = E(X) = \sum x_i p(x_i)$;
 Variance $\sigma_X^2 = V(X) = E(X - \mu_X)^2 = E(X^2) - \mu_X^2 = \sum x_i^2 p(x_i) - (\sum x_i p(x_i))^2$.
 (2.2) Continuous r.v. X has a pdf $f(x)$ satisfying $f(x) \geq 0$ and $\int_{-\infty}^{\infty} f(x) dx = 1$,

$$P(a < X < b) = \int_a^b f(x) dx.$$

Expectation (or *Expected value* or *mean*) $\mu_X = E(X) = \int_{-\infty}^{\infty} x f(x) dx$;

Variance $\sigma_X^2 = V(X) = E(X - \mu_X)^2 = E(X^2) - \mu_X^2 = \int_{-\infty}^{\infty} x^2 f(x) dx - (\int_{-\infty}^{\infty} x f(x) dx)^2$.

- (2.3) Cumulative distribution function (cdf) of a r.v. X is $F(x) = P(X \leq x)$.

- (2.4) X and Y are r.v.s, a, b and c are scalars, then

$$E(aX + bY + c) = aE(X) + bE(Y) + c,$$

$$V(aX + bY + c) = a^2V(X) + b^2V(Y) + 2ab \operatorname{cov}(X, Y),$$

$$E(g(X, Y)) = \begin{cases} \sum_{i,j} g(x_i, y_j) \cdot p(x_i, y_j), & \text{for discrete } (X, Y), \\ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) \cdot f(x, y) dx dy, & \text{for continuous } (X, Y). \end{cases}$$

- (2.5) • Discrete r.v. (X, Y) has a joint pmf $p(x, y)$ satisfying $p(x, y) \geq 0$ and $\sum_{x_i} \sum_{y_j} p(x_i, y_j) = 1$.

The *marginal pmf* of X is $p_X(x) = \sum_y p(x, y)$;

The *marginal pmf* of Y is $p_Y(y) = \sum_x p(x, y)$;

X and Y are *independent* if $p(x, y) = p_X(x) \cdot p_Y(y)$.

- Continuous r.v. (X, Y) has a joint pdf $p(x, y)$ satisfying $f(x, y) \geq 0$ and $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) dx dy = 1$.

The *marginal pdf* of X is $f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy$;

The *marginal pdf* of Y is $f_Y(y) = \int_{-\infty}^{\infty} f(x, y) dx$;

X and Y are *independent* if $f(x, y) = f_X(x) \cdot f_Y(y)$.

3. Several special r.v.s

- (3.1) $X \sim \operatorname{Bin}(n, p)$ has a pmf $p(x) = P(X = x) = \binom{n}{x} \cdot p^x \cdot (1-p)^{n-x}$, $x = 0, 1, 2, \dots, n$.

$$E(X) = n \cdot p, \quad V(X) = n \cdot p \cdot (1-p).$$

- (3.2) $X \sim \operatorname{Po}(\lambda)$ has a pmf $p(x) = P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$, $x = 0, 1, 2, \dots$
 $E(X) = \lambda, \quad V(X) = \lambda$.

- (3.3) $X \sim \operatorname{Hypergeometric}$ has a pmf $p(x) = P(X = x) = \frac{\binom{M}{x} \binom{N-M}{n-x}}{\binom{N}{n}}$.

- (3.4) $X \sim \operatorname{Exp}(\lambda)$ has a pdf

$$f(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0, \\ 0, & \text{otherwise.} \end{cases}$$

- (3.5) $X \sim N(\mu, \sigma^2)$ has a pdf

$$f(x) = \frac{1}{\sigma \sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad -\infty < x < \infty.$$

- (3.6) $X \sim U(a, b)$ has a pdf

$$f(x) = \begin{cases} \frac{1}{b-a}, & a < x < b, \\ 0, & \text{otherwise.} \end{cases}$$

$$E(X) = \frac{a+b}{2}, \quad V(X) = \frac{(b-a)^2}{12}.$$

4. Central Limit Theorem (CLT)

Suppose that a population has mean $= \mu$ and variance $= \sigma^2$. A random sample $\{X_1, X_2, \dots, X_n\}$ from this population is given. Then for large $n \geq 30$,

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1). \tag{1}$$

- If the population is normal, then (1) holds for any n .
- Note that $\mu = E(\bar{X})$ and $(\sigma/\sqrt{n})^2 = V(\bar{X})$.

5. Several notations in statistics

- (5.1) Sample mean: $\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n} = \sum \bar{X}_i$; $\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \sum x_i$.

- (5.2) Sample variance:

$$S^2 = \frac{\sum (X_i - \bar{X})^2}{n-1} = \frac{1}{n-1} \left(\sum X_i^2 - \frac{(\sum X_i)^2}{n} \right); \quad s^2 = \frac{\sum (x_i - \bar{x})^2}{n-1} = \frac{1}{n-1} \left(\sum x_i^2 - \frac{(\sum x_i)^2}{n} \right).$$

- Capital letters \bar{X} and S^2 refer to the objects based on random sample (therefore they are in general r.v.s), while small letters \bar{x} and s^2 are the objects based on observations (so they are scalars).

- (5.3) A point estimator of θ obtained by Methods of Moments is denoted as $\hat{\theta}_{MM}$.

- (5.4) A point estimator of θ obtained by Maximum Likelihood method is denoted as $\hat{\theta}_{ML}$.

6. Confidence Interval (CI)

In this course, three types of confidence intervals are studied depending on the unknown population parameter(s): CI-1 (confidence intervals for population mean(s)), CI-2 (confidence intervals for population variance(s)), and CI-3 (confidence intervals for population proportion(s)).

CI-1: (1 - α) CI of a population mean μ

case 1.1 (any n) If population $X \sim N(\mu, \sigma^2)$ and σ^2 is known, then $\frac{\bar{X}-\mu}{\sigma/\sqrt{n}} \sim N(0, 1)$ and

$$I_\mu = (\bar{x} - z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}, \bar{x} + z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}) := \bar{x} \mp z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}.$$

case 1.2 (n ≥ 30) For any population X, it holds that $\frac{\bar{X}-\mu}{\sigma/\sqrt{n}} \sim N(0, 1)$ and

$$I_\mu = \bar{x} \mp z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \text{ or } I_\mu = \bar{x} \mp z_{\alpha/2} \cdot \frac{\hat{\sigma}}{\sqrt{n}}.$$

case 1.3 (any n) If population $X \sim N(\mu, \sigma^2)$ and σ^2 is unknown, then $\frac{\bar{X}-\mu}{S/\sqrt{n}} \sim T(n-1)$ and

$$I_\mu = \bar{x} \mp t_{\alpha/2}(n-1) \cdot \frac{s}{\sqrt{n}}.$$

CI-1': (1 - α) CI of the difference of two population means $\mu_X - \mu_Y$

case 1.1' (any n_1, n_2) If independent populations $X \sim N(\mu_X, \sigma_X^2)$, $Y \sim N(\mu_Y, \sigma_Y^2)$, and σ_X^2, σ_Y^2 are known, then

$$\frac{(\bar{X}-\bar{Y})-(\mu_X-\mu_Y)}{\sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}} \sim N(0, 1), \text{ and } I_{\mu_X-\mu_Y} = (\bar{x}-\bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}.$$

case 1.2' ($n_1, n_2 \geq 30$) For any independent populations X and Y, it holds that


$$\frac{(\bar{X}-\bar{Y})-(\mu_X-\mu_Y)}{\sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}} \sim N(0, 1) \text{ and}$$

$$I_{\mu_X-\mu_Y} = (\bar{x}-\bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}} \text{ or } I_{\mu_X-\mu_Y} = (\bar{x}-\bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{\sigma}_X^2}{n_1} + \frac{\hat{\sigma}_Y^2}{n_2}}.$$

case 1.3' (any n_1, n_2) If independent populations $X \sim N(\mu_X, \sigma_X^2)$, $Y \sim N(\mu_Y, \sigma_Y^2)$, where σ_X^2, σ_Y^2 are unknown but $\sigma_X^2 = \sigma_Y^2$, then

$$\frac{(\bar{X}-\bar{Y})-(\mu_X-\mu_Y)}{S\sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} \sim T(n_1+n_2-2), \text{ where } S^2 = \frac{(n_1-1)S_X^2 + (n_2-1)S_Y^2}{n_1+n_2-2}, \text{ and}$$

$$I_{\mu_X-\mu_Y} = (\bar{x}-\bar{y}) \mp t_{\alpha/2}(n_1+n_2-2) \cdot s\sqrt{\frac{1}{n_1} + \frac{1}{n_2}}.$$



CI-2: (1 - α) CI of population variance(s) σ^2


• If a population $X \sim N(\mu, \sigma^2)$ and σ^2 is unknown, then $\frac{(n-1)S^2}{\sigma^2} \sim \chi^2(n-1)$, and

$$I_{\sigma^2} = \left(\frac{(n-1)s^2}{\chi_{\alpha/2}^2(n-1)}, \frac{(n-1)s^2}{\chi_{1-\alpha/2}^2(n-1)} \right).$$

• If two independent populations $X \sim N(\mu_X, \sigma^2)$ and $Y \sim N(\mu_Y, \sigma^2)$, and σ^2 is unknown, then $\frac{(n_1+n_2-2)S^2}{\sigma^2} \sim \chi^2(n_1+n_2-2)$, and

$$I_{\sigma^2} = \left(\frac{(n_1+n_2-2)s^2}{\chi_{\alpha/2}^2(n_1+n_2-2)}, \frac{(n_1+n_2-2)s^2}{\chi_{1-\alpha/2}^2(n_1+n_2-2)} \right),$$

where $S^2 = \frac{(n_1-1)S_X^2 + (n_2-1)S_Y^2}{n_1+n_2-2}$.



CI-3: (1 - α) CI of population proportion(s)

• If a (large) population has an unknown proportion p, then $\frac{\hat{p}-p}{\sqrt{p(1-p)/n}} \sim N(0, 1)$ if $n\hat{p} \geq 10$ and $n(1-\hat{p}) \geq 10$ with $\hat{p} = x/n$, and $I_p = \hat{p} \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$.

• If two independent (large) populations have unknown proportions p_1 and p_2 , then

$$\frac{(\hat{p}_1-\hat{p}_2)-(p_1-p_2)}{\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{\hat{p}_2(1-\hat{p}_2)}{n_2}}} \sim N(0, 1)$$

if $n_i\hat{p}_i \geq 10$ and $n_i(1-\hat{p}_i) \geq 10$ for $i = 1, 2$, and $I_{p_1-p_2} = (\hat{p}_1-\hat{p}_2) \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{\hat{p}_2(1-\hat{p}_2)}{n_2}}$.

7. Hypothesis Test (HT)

	H_0 is true	H_0 is false and $\theta = \theta_1$
reject H_0	(type I error or significance level) α	(power) $h(\theta_1)$
don't reject H_0	$1 - \alpha$	(type II error) $\beta(\theta_1) = 1 - h(\theta_1)$

reject $H_0 \Leftrightarrow TS \in C \Leftrightarrow p\text{-value} < \alpha$

χ^2 tests for populations (non-parametric tests)

Suppose that for a random sample of a population X, the n elements of it are classified into k disjoint groups $A_i, 1 \leq i \leq k$. For each group $A_i, 1 \leq i \leq k$, suppose that there are $N_{i1}, 1 \leq i \leq k$ elements inside. Let $p_i = P(A_i)$ assuming a given distribution of X. Note that $p_1 + p_2 + \dots + p_k = 1$ and $N_1 + N_2 + \dots + N_k = n$. One wants to test the hypotheses

$$H_0 : P(A_i) = p_i, \quad 1 \leq i \leq k, \quad H_a : P(A_i) \neq p_i \text{ for some } 1 \leq i \leq k.$$

If n is large in the sense that $np_i \geq 5$ for all $1 \leq i \leq k$, then the test statistic is

$$\sum_{i=1}^k \frac{(N_i - np_i)^2}{np_i} \approx \chi^2(k-1).$$

Therefore the observation of the test statistic is

$$TS = \sum_{i=1}^k \frac{(n_i - np_i)^2}{np_i}, \text{ where } n_i \text{ is the observation of } N_i, 1 \leq i \leq k.$$

For the critical region C , one can take (note that if H_0 is true, then TS should be close to zero)

$$C = (\chi^2_{\alpha}(k-1), \infty).$$

The conclusion would be $TS \in C \iff H_0$ is rejected.

8. Linear and logistic regression

(Multiple) linear regression: $Y = \beta_0 + \beta_1 x_1 + \dots + \beta_k x_k + \varepsilon$, $\varepsilon \sim N(0, \sigma^2)$.

- Y : response variable (which is normal r.v.), $\{x_1, \dots, x_k\}$: predictors (which are scalars).
- sample: $\{(x_{11}, \dots, x_{1k}; y_1), (x_{21}, \dots, x_{2k}; y_2), \dots, (x_{n1}, \dots, x_{nk}; y_n)\}$.
- how to estimate $\beta_j \approx \hat{\beta}_j$: least square method, that is, to minimize $\sum_{i=1}^n (y_i - \hat{y}_i)^2$, where the estimated (multiple) linear regression line \hat{y} is

$$\hat{y} = \hat{\beta}_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k.$$

- $\frac{\hat{\beta}_j - \beta_j}{se(\hat{\beta}_j)} \sim T(n-k-1)$, this helps determine whether or not the real $\beta_j = 0$?
- $\sigma^2 \approx \frac{SSE}{n-k-1}$, this gives an estimation of the size of the error.
- $R^2 = \frac{SSR}{SSY}$, this gives how well the model is (if $R^2 \approx 1$, then the model fits the sample very well).
- How to test $\beta_1 = \dots = \beta_k = 0$? Use the random variable $\frac{SSR/k}{SSE/(n-k-1)} \sim F(k, n-k-1)$.

Logistic regression: Let Y can only take 0 or 1 with $P(Y=1) = p$ and $P(Y=0) = 1-p$.

$$E(Y) = p(x_1, \dots, x_k) = \frac{e^{\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k}}{1 + e^{\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k}}.$$

- Y : response variable (which is Bernoulli r.v. $P(Y=1) = p$ and $P(Y=0) = 1-p$, so $E(Y) = p$), $\{x_1, \dots, x_k\}$: predictors (which are scalars).
- sample: $\{(x_{11}, \dots, x_{1k}; y_1), (x_{21}, \dots, x_{2k}; y_2), \dots, (x_{n1}, \dots, x_{nk}; y_n)\}$.
- how to estimate $\beta_j \approx \hat{\beta}_j$: maximal likelihood method (maximize $\prod_{i=1}^n p(x_{i1}, \dots, x_{ik})^{y_i} (1 - p(x_{i1}, \dots, x_{ik}))^{1-y_i}$).
- $\frac{\hat{\beta}_j - \beta_j}{se(\hat{\beta}_j)} \approx N(0, 1)$ for large $n \geq 30$, this helps determine whether or not the real $\beta_j = 0$?
- Classification of a new object $Y(x_1, \dots, x_k)$ as 1 or 0 according

$$Y(x_1, \dots, x_k) = \begin{cases} 1, & \text{if } \hat{p}(x_1, \dots, x_k) \geq 0.5, \\ 0, & \text{if } \hat{p}(x_1, \dots, x_k) < 0.5, \end{cases}$$

where the estimated logit function $\hat{p}(x_1, \dots, x_k)$ is

$$\hat{p}(x_1, \dots, x_k) = \frac{e^{\hat{\beta}_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k}}{1 + e^{\hat{\beta}_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k}}.$$

9. Tables

(9.1) Table for $N(0, 1)$ standard normal random variable $\Phi(x) = P(N(0, 1) \leq x)$, $x \geq 0$.
There is an important relation $\Phi(-x) = 1 - \Phi(x)$, $x \geq 0$.

x	0	1	2	3	4	5	6	7	8	9
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9564	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9992	0.9993	0.9993	0.9993
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997	0.9997
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998	0.9998	0.9998
3.5	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998
3.6	0.9998	0.9998	0.9998	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.7	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.8	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.9	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
4.0	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

(9.2) Table for $T(f)$ random variable $F(x) = P(T(f) \leq x)$,
where f is a parameter called 'degrees of freedom'.

f	0.75	0.90	0.95	0.975	0.99	0.995	0.9975	0.9995
1	1.00	3.08	6.31	12.71	31.82	63.66	127.32	636.62
2	0.82	1.89	2.92	4.30	6.96	9.92	14.09	31.60
3	0.76	1.64	2.35	3.18	4.54	5.84	7.45	12.92
4	0.74	1.53	2.13	2.78	3.75	4.60	5.60	8.61
5	0.73	1.48	2.02	2.57	3.36	4.03	4.77	6.87
6	0.72	1.44	1.94	2.45	3.14	3.71	4.32	5.96
7	0.71	1.41	1.89	2.36	3.00	3.50	4.03	5.41
8	0.71	1.40	1.86	2.31	2.90	3.36	3.83	5.04
9	0.70	1.38	1.83	2.26	2.82	3.25	3.69	4.78
10	0.70	1.37	1.81	2.23	2.76	3.17	3.58	4.59
11	0.70	1.36	1.80	2.20	2.72	3.11	3.50	4.44
12	0.70	1.36	1.78	2.18	2.68	3.05	3.43	4.32
13	0.69	1.35	1.77	2.16	2.65	3.01	3.37	4.22
14	0.69	1.35	1.76	2.14	2.62	2.98	3.33	4.14
15	0.69	1.34	1.75	2.13	2.60	2.95	3.29	4.07
16	0.69	1.34	1.75	2.12	2.58	2.92	3.25	4.01
17	0.69	1.33	1.74	2.11	2.57	2.90	3.22	3.97
18	0.69	1.33	1.73	2.10	2.55	2.88	3.20	3.92
19	0.69	1.33	1.73	2.09	2.54	2.86	3.17	3.88
20	0.69	1.33	1.72	2.09	2.53	2.85	3.15	3.85
21	0.69	1.32	1.72	2.08	2.52	2.83	3.14	3.82
22	0.69	1.32	1.72	2.07	2.51	2.82	3.12	3.79
23	0.69	1.32	1.71	2.07	2.50	2.81	3.10	3.77
24	0.68	1.32	1.71	2.06	2.49	2.80	3.09	3.75
25	0.68	1.32	1.71	2.06	2.49	2.79	3.08	3.73
26	0.68	1.31	1.71	2.06	2.48	2.78	3.07	3.71
27	0.68	1.31	1.70	2.05	2.47	2.77	3.06	3.69
28	0.68	1.31	1.70	2.05	2.47	2.76	3.05	3.67
29	0.68	1.31	1.70	2.05	2.46	2.76	3.04	3.66
30	0.68	1.31	1.70	2.04	2.46	2.75	3.03	3.65
40	0.68	1.30	1.68	2.02	2.42	2.70	2.97	3.55
50	0.68	1.30	1.68	2.01	2.40	2.68	2.94	3.50
60	0.68	1.30	1.67	2.00	2.39	2.66	2.91	3.46
100	0.68	1.29	1.66	1.98	2.36	2.63	2.87	3.39
∞	0.67	1.28	1.65	1.96	2.33	2.58	2.81	3.29

(9.3) Table for $\chi^2(f)$ random variable $F(x) = P(\chi^2(f) \leq x)$, where f is a parameter.

f	0.0005	0.001	0.005	0.01	0.025	0.05	0.10	0.20	0.30	0.40	0.50
1	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.06	0.15	0.27	0.45
2	0.00	0.00	0.01	0.02	0.05	0.10	0.21	0.45	0.71	1.02	1.39
3	0.02	0.02	0.07	0.11	0.22	0.35	0.58	1.01	1.42	1.87	2.37
4	0.06	0.09	0.21	0.30	0.48	0.71	1.06	1.65	2.19	2.75	3.36
5	0.16	0.21	0.41	0.55	0.83	1.15	1.61	2.34	3.00	3.66	4.35
6	0.30	0.38	0.68	0.87	1.24	1.64	2.20	3.07	3.83	4.57	5.35
7	0.48	0.60	0.99	1.24	1.69	2.17	2.73	3.62	4.67	5.49	6.35
8	0.71	0.86	1.34	1.65	2.18	2.73	3.49	4.59	5.53	6.42	7.34
9	0.97	1.15	1.73	2.09	2.70	3.33	4.17	5.38	6.39	7.36	8.34
10	1.26	1.48	2.16	2.56	3.25	3.94	4.87	6.18	7.27	8.30	9.34
11	1.59	1.83	2.60	3.05	3.82	4.57	5.58	6.99	8.15	9.24	10.34
12	1.93	2.21	3.07	3.57	4.40	5.23	6.30	7.81	9.03	10.18	11.34
13	2.31	2.62	3.57	4.11	5.01	5.89	7.04	8.63	9.93	11.13	12.34
14	2.70	3.04	4.07	4.66	5.63	6.57	7.79	9.47	10.82	12.08	13.34
15	3.11	3.48	4.60	5.23	6.26	7.26	8.55	10.31	11.72	13.03	14.34
16	3.54	3.94	5.14	5.81	6.91	7.96	9.31	11.15	12.62	13.98	15.34
17	3.98	4.42	5.70	6.41	7.56	8.67	10.09	12.00	13.53	14.94	16.34
18	4.44	4.90	6.26	7.01	8.23	9.39	10.86	12.86	14.44	15.89	17.34
19	4.91	5.41	6.84	7.63	8.91	10.12	11.65	13.72	15.35	16.85	18.34
20	5.40	5.92	7.43	8.26	9.59	10.85	12.44	14.58	16.27	17.81	19.34
21	5.90	6.45	8.03	8.90	10.28	11.59	13.24	15.44	17.18	18.77	20.34
22	6.40	6.98	8.64	9.54	10.98	12.34	14.04	16.31	18.10	19.73	21.34
23	6.92	7.53	9.26	10.20	11.69	13.09	14.85	17.19	19.02	20.69	22.34
24	7.45	8.08	9.89	10.86	12.40	13.85	15.66	18.06	19.94	21.65	23.34
25	7.99	8.65	10.52	11.52	13.12	14.61	16.47	18.94	20.87	22.62	24.34
26	8.54	9.22	11.16	12.20	13.84	15.38	17.29	19.82	21.79	23.58	25.34
27	9.09	9.80	11.81	12.88	14.57	16.15	18.11	20.70	22.72	24.54	26.34
28	9.66	10.39	12.46	13.56	15.31	16.93	18.94	21.59	23.65	25.51	27.34
29	10.23	10.99	13.12	14.26	16.05	17.71	19.77	22.48	24.58	26.48	28.34
30	10.80	11.59	13.79	14.95	16.79	18.49	20.60	23.36	25.51	27.44	29.34
40	16.91	17.92	20.71	22.16	24.43	26.51	29.05	32.34	34.87	37.13	39.34
50	23.46	24.67	27.99	29.71	32.36	34.76	37.69	41.45	46.86	49.33	51.33
60	30.34	31.74	35.53	37.48	40.48	43.19	46.46	50.64	53.81	56.62	59.33
100	59.90	61.92	67.33	70.06	74.22	77.93	82.36	87.95	92.13	95.81	99.33

Table for $\chi^2(f)$ random variable $F(x) = P(\chi^2(f) \leq x)$, where f is a parameter.

f	0.60	0.70	0.80	0.90	0.95	0.975	0.99	0.995	0.999	0.9995
1	0.71	1.07	1.64	2.71	3.84	5.02	6.63	7.88	10.83	12.12
2	1.83	2.41	3.22	4.61	5.99	7.38	9.21	10.60	13.82	15.20
3	2.95	3.66	4.64	6.25	7.81	9.35	11.34	12.84	16.27	17.73
4	4.04	4.88	5.99	7.78	9.49	11.14	13.28	14.86	18.47	20.00
5	5.13	6.06	7.29	9.24	11.07	12.83	15.09	16.75	20.52	22.11
6	6.21	7.23	8.56	10.64	12.59	14.45	16.81	18.55	22.46	24.10
7	7.28	8.38	9.80	12.02	14.07	16.01	18.48	20.28	24.32	26.02
8	8.35	9.52	11.03	13.36	15.51	17.53	20.09	21.95	26.12	27.87
9	9.41	10.66	12.24	14.68	16.92	19.02	21.67	23.59	27.88	29.67
10	10.47	11.78	13.44	15.99	18.31	20.48	23.21	25.19	29.59	31.42
11	11.53	12.90	14.63	17.28	19.68	21.92	24.72	26.76	31.26	33.14
12	12.58	14.01	15.81	18.55	21.03	23.34	26.22	28.30	32.91	34.82
13	13.64	15.12	16.98	19.81	22.36	24.74	27.69	29.82	34.53	36.48
14	14.69	16.22	18.15	21.06	23.68	26.12	29.14	31.32	36.12	38.11
15	15.73	17.32	19.31	22.31	25.00	27.49	30.58	32.80	37.70	39.72
16	16.78	18.42	20.47	23.54	26.30	28.85	32.00	34.27	39.25	41.31
17	17.82	19.51	21.61	24.77	27.59	30.19	33.41	35.72	40.79	42.88
18	18.87	20.60	22.76	25.99	28.87	31.53	34.81	37.16	42.31	44.43
19	19.91	21.69	23.90	27.20	30.14	32.85	36.19	38.58	43.82	45.97
20	20.95	22.77	25.04	28.41	31.41	34.17	37.57	40.00	45.31	47.50
21	21.99	23.86	26.17	29.62	32.67	35.48	38.93	41.40	46.80	49.01
22	23.03	24.94	27.30	30.81	33.92	36.78	40.29	42.80	48.27	50.51
23	24.07	26.02	28.43	32.01	35.17	38.08	41.64	44.18	49.73	52.00
24	25.11	27.10	29.55	33.20	36.42	39.36	42.98	45.56	51.18	53.48
25	26.14	28.17	30.68	34.38	37.65	40.65	44.31	46.93	52.62	54.95
26	27.18	29.25	31.79	35.56	38.89	41.92	45.64	48.29	54.05	56.41
27	28.21	30.32	32.91	36.74	40.11	43.19	46.96	49.64	55.48	57.86
28	29.25	31.39	34.03	37.92	41.34	44.46	48.28	50.99	56.89	59.30
29	30.28	32.46	35.14	39.09	42.56	45.72	49.59	52.34	58.30	60.73
30	31.32	33.53	36.25	40.26	43.77	46.98	50.89	53.67	59.70	62.16
40	41.62	44.16	47.27	51.81	55.76	59.34	63.69	66.77	73.40	76.09
50	51.89	54.72	58.16	63.17	67.50	71.42	76.15	79.49	86.66	89.56
60	62.13	65.23	68.97	74.40	79.08	83.30	88.38	91.95	99.61	102.69
100	102.95	106.91	111.67	118.50	124.34	129.56	135.81	140.17	149.45	153.17

(9.4) Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
 If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1 - p) \geq n - k)$.

n	k	0.05	0.10	0.15	0.20	0.25	0.30	0.35	0.40	0.45	0.50
2	0	0.9025	0.8100	0.7225	0.6400	0.5625	0.4900	0.4225	0.3600	0.3025	0.2500
	1	0.9975	0.9900	0.9775	0.9600	0.9375	0.9100	0.8775	0.8400	0.7975	0.7500
3	0	0.8574	0.7290	0.6141	0.5120	0.4219	0.3430	0.2746	0.2160	0.1664	0.1250
	1	0.9928	0.9720	0.9392	0.8960	0.8438	0.7840	0.7183	0.6480	0.5747	0.5000
4	0	0.8145	0.6561	0.5220	0.4096	0.3164	0.2401	0.1785	0.1256	0.0915	0.0625
	1	0.9860	0.9477	0.8905	0.8192	0.7383	0.6517	0.5630	0.4735	0.3910	0.3125
5	0	0.7738	0.5905	0.4437	0.3277	0.2373	0.1681	0.1160	0.0778	0.0503	0.0313
	1	0.9774	0.9185	0.8352	0.7373	0.6328	0.5282	0.4284	0.3370	0.2562	0.1875
6	0	0.7351	0.5314	0.3771	0.2621	0.1780	0.1176	0.0754	0.0467	0.0277	0.0156
	1	0.9672	0.8847	0.7765	0.6554	0.5339	0.4202	0.3191	0.2333	0.1636	0.1094
7	0	0.6983	0.4783	0.3206	0.2097	0.1335	0.0824	0.0490	0.0280	0.0152	0.0078
	1	0.9566	0.8503	0.7166	0.5767	0.4449	0.3294	0.2338	0.1586	0.1024	0.0625
8	0	0.6634	0.4305	0.2725	0.1678	0.1001	0.0576	0.0319	0.0168	0.0084	0.0039
	1	0.9428	0.8131	0.6572	0.5033	0.3671	0.2553	0.1691	0.1064	0.0632	0.0352
9	0	0.6302	0.3874	0.2316	0.1342	0.0751	0.0404	0.0207	0.0101	0.0046	0.0020
	1	0.9288	0.7748	0.5995	0.4362	0.3003	0.1960	0.1211	0.0705	0.0385	0.0195

Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
 If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1 - p) \geq n - k)$.

n	k	0.05	0.10	0.15	0.20	0.25	0.30	0.35	0.40	0.45	0.50
10	0	0.5987	0.3487	0.1969	0.1074	0.0563	0.0282	0.0135	0.0060	0.0025	0.0010
	1	0.9139	0.7361	0.5443	0.3758	0.2440	0.1493	0.0860	0.0464	0.0233	0.0107
11	0	0.5688	0.3138	0.1673	0.0859	0.0422	0.0198	0.0088	0.0036	0.0014	0.0005
	1	0.8981	0.6974	0.4922	0.3221	0.1971	0.1130	0.0606	0.0302	0.0139	0.0059
12	0	0.5404	0.2824	0.1422	0.0687	0.0317	0.0138	0.0057	0.0022	0.0008	0.0002
	1	0.8816	0.6590	0.4435	0.2749	0.1584	0.0850	0.0424	0.0196	0.0083	0.0032

Table for Binomial random variable $P(\text{Bin}(n, p) \leq k)$ if $p \leq 0.5$.
 If $p > 0.5$, then $P(\text{Bin}(n, p) \leq k) = P(\text{Bin}(n, 1 - p) \geq n - k)$.

n	k	0.05	0.10	0.15	0.20	0.25	0.30	0.35	0.40	0.45	0.50
14	0	0.4877	0.2288	0.1028	0.0440	0.0178	0.0068	0.0024	0.0008	0.0002	0.0001
	1	0.8470	0.5846	0.3567	0.2075	0.1010	0.0475	0.0205	0.0081	0.0029	0.0009
	2	0.9699	0.8416	0.6479	0.4481	0.2811	0.1608	0.0839	0.0398	0.0170	0.0065
	3	0.9958	0.9559	0.8535	0.6982	0.5213	0.3552	0.2205	0.1243	0.0632	0.0287
	4	0.9996	0.9908	0.9533	0.8702	0.7415	0.5842	0.4227	0.2793	0.1672	0.0898
	5	1.0000	0.9985	0.9885	0.9561	0.8883	0.7805	0.6405	0.4859	0.3373	0.2120
	6	1.0000	0.9998	0.9978	0.9884	0.9617	0.9067	0.8164	0.6925	0.5461	0.3953
	7	1.0000	1.0000	0.9997	0.9976	0.9897	0.9685	0.9247	0.8499	0.7414	0.6047
	8	1.0000	1.0000	1.0000	0.9996	0.9978	0.9978	0.9917	0.9417	0.8811	0.7880
	9	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9983	0.9940	0.9825	0.9102
	10	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9998	0.9989	0.9961	0.9886
	11	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9994	0.9978	0.9935
	12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9991
	13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999
15	0	0.4633	0.2059	0.0874	0.0352	0.0134	0.0047	0.0016	0.0005	0.0001	0.0000
	1	0.8290	0.5490	0.3186	0.1671	0.0802	0.0353	0.0142	0.0052	0.0017	0.0005
	2	0.9638	0.8159	0.6042	0.3980	0.2361	0.1268	0.0617	0.0271	0.0107	0.0037
	3	0.9945	0.9444	0.8227	0.6482	0.4613	0.2969	0.1727	0.0905	0.0424	0.0176
	4	0.9994	0.9873	0.9383	0.8338	0.6865	0.5155	0.3519	0.2173	0.1204	0.0592
	5	0.9999	0.9978	0.9832	0.9389	0.8516	0.7216	0.5816	0.4303	0.2608	0.1509
	6	1.0000	0.9997	0.9964	0.9819	0.9434	0.8689	0.7548	0.6098	0.4522	0.3036
	7	1.0000	1.0000	0.9994	0.9958	0.9827	0.9500	0.8868	0.7869	0.6535	0.5000
	8	1.0000	1.0000	0.9999	0.9992	0.9958	0.9848	0.9578	0.9050	0.8182	0.6964
	9	1.0000	1.0000	1.0000	0.9999	0.9999	0.9992	0.9963	0.9921	0.9231	0.8491
	10	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9997	0.9972	0.9607	0.9408
	11	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9995	0.9981	0.9824
	12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9989	0.9963
	13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9995
	14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
	16	0	0.4401	0.1853	0.0743	0.0281	0.0100	0.0033	0.0010	0.0003	0.0001
1		0.8108	0.5147	0.2839	0.1407	0.0635	0.0261	0.0098	0.0033	0.0010	0.0003
2		0.9571	0.7892	0.5614	0.3518	0.1971	0.0994	0.0451	0.0183	0.0066	0.0021
3		0.9930	0.9316	0.7899	0.5981	0.4050	0.2459	0.1339	0.0651	0.0281	0.0106
4		0.9991	0.9830	0.9209	0.7982	0.6302	0.4499	0.2892	0.1666	0.0853	0.0384
5		0.9999	0.9967	0.9765	0.9183	0.8103	0.6598	0.4900	0.3288	0.1976	0.1051
6		1.0000	0.9995	0.9944	0.9733	0.9204	0.8247	0.6881	0.5272	0.3660	0.2272
7		1.0000	0.9999	0.9989	0.9930	0.9729	0.9236	0.8406	0.7161	0.5629	0.4018
8		1.0000	1.0000	0.9998	0.9985	0.9925	0.9743	0.9329	0.8577	0.7441	0.5982
9		1.0000	1.0000	1.0000	0.9998	0.9988	0.9929	0.9711	0.9417	0.8759	0.7728
10		1.0000	1.0000	1.0000	1.0000	0.9997	0.9984	0.9984	0.9987	0.9851	0.9616
11		1.0000	1.0000	1.0000	1.0000	1.0000	0.9997	0.9997	0.9991	0.9951	0.9849
12		1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9998	0.9991	0.9965	0.9894
13		1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9994	0.9994	0.9979
14		1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9997
15		1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

Table for Binomial random variable $P(\text{Bin}(n, p) \leq k)$ if $p \leq 0.5$.
 If $p > 0.5$, then $P(\text{Bin}(n, p) \leq k) = P(\text{Bin}(n, 1 - p) \geq n - k)$.

n	k	0.05	0.10	0.15	0.20	0.25	0.30	0.35	0.40	0.45	0.50
17	0	0.4181	0.1668	0.0631	0.0225	0.0075	0.0023	0.0007	0.0002	0.0000	0.0000
	1	0.7922	0.4818	0.2525	0.1182	0.0501	0.0193	0.0067	0.0021	0.0006	0.0001
	2	0.9497	0.7618	0.5198	0.3096	0.1637	0.0774	0.0327	0.0123	0.0041	0.0012
	3	0.9912	0.9174	0.7556	0.5489	0.3530	0.2019	0.1028	0.0464	0.0184	0.0064
	4	0.9988	0.9779	0.9013	0.7582	0.5739	0.3887	0.2348	0.1260	0.0596	0.0245
	5	0.9999	0.9953	0.9681	0.8943	0.7653	0.5668	0.4197	0.2639	0.1471	0.0717
	6	1.0000	0.9992	0.9922	0.9623	0.8929	0.7752	0.6188	0.4478	0.2902	0.1672
	7	1.0000	0.9999	0.9983	0.9891	0.9598	0.8954	0.8054	0.7872	0.6405	0.4743
	8	1.0000	1.0000	0.9997	0.9974	0.9876	0.9597	0.9006	0.8011	0.6626	0.5000
	9	1.0000	1.0000	1.0000	0.9995	0.9969	0.9969	0.9873	0.9617	0.9081	0.8166
	10	1.0000	1.0000	1.0000	0.9999	0.9994	0.9994	0.9980	0.9852	0.9174	0.8338
	11	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9993	0.9970	0.9894	0.9699
	12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9994	0.9914	0.9755
	13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9981	0.9936
	14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9988
	15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999
	18	0	0.3972	0.1501	0.0536	0.0180	0.0056	0.0016	0.0004	0.0001	0.0000
1		0.7735	0.4503	0.2241	0.0991	0.0395	0.0142	0.0046	0.0013	0.0003	0.0001
2		0.9419	0.7338	0.4797	0.2713	0.1353	0.0600	0.0236	0.0082	0.0025	0.0008
3		0.9891	0.9018	0.7202	0.5010	0.3057	0.1646	0.0783	0.0328	0.0120	0.0038
4		0.9985	0.9718	0.8794	0.7164	0.5187	0.3327	0.1886	0.0942	0.0411	0.0154
5		0.9998	0.9936	0.9351	0.8671	0.7175	0.5344	0.3550	0.2088	0.1077	0.0481
6		1.0000	0.9988	0.9882	0.9487	0.8610	0.7217	0.5491	0.3258	0.1189	0.0403
7		1.0000	0.9998	0.9973	0.9837	0.9431	0.8593	0.7283	0.5634	0.3915	0.2403
8		1.0000	1.0000	0.9995	0.9957	0.9807	0.9404	0.8609	0.7368	0.5778	0.4073
9		1.0000	1.0000	0.9999	0.9991	0.9946	0.9790	0.9403	0.8653	0.7473	0.5827
10		1.0000	1.0000	1.0000	0.9998	0.9988	0.9939	0.9788	0.9259	0.8720	0.7597
11		1.0000	1.0000	1.0000	0.9998	0.9998	0.9986	0.9937	0.9797	0.9463	0.8811
12		1.0000	1.0000	1.0000	1.0000	0.9997	0.9997	0.9986	0.9942	0.9817	0.9519
13		1.0000	1.0000	1.0000	1.0000	1.0000	0.9997	0.9997	0.9987	0.9951	0.9846
14		1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9998	0.9990	0.9962
15		1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9993
16		1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999
19	0	0.3774	0.1351	0.0456	0.0144	0.0042	0.0011	0.0003	0.0001	0.0000	0.0000
	1	0.7547	0.4203	0.1985	0.0829	0.0310	0.0104	0.0031	0.0008	0.0002	0.0000
	2	0.9335	0.7054	0.4413	0.2369	0.1113	0.0462	0.0170	0.0055	0.0015	0.0002
	3	0.9868	0.8850	0.6841	0.4551	0.2631	0.1332	0.0591	0.0230	0.0077	0.0024
	4	0.9980	0.9648	0.8556	0.6733	0.4654	0.2822	0.1500	0.0696	0.0280	0.0096
	5	0.9998	0.9914	0.9463	0.8369	0.6678	0.4739	0.2968	0.1629	0.0777	0.0318
	6	1.0000	0.9983	0.9837	0.9324	0.8251	0.6655	0.4812	0.3081	0.1727	0.0835
	7	1.0000	0.9997	0.9959	0.9767	0.9225	0.8180	0.6656	0.4878	0.3169	0.1796
	8	1.0000	1.0000	0.9992	0.9933	0.9713	0.9161	0.8145	0.6675	0.4940	0.3238

(9.5) Table for Poisson random variable $P(Po(\mu) \leq k)$.

k	μ									
	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0
0	0.9048	0.8187	0.7408	0.6703	0.6065	0.5488	0.4966	0.4493	0.4066	0.3679
1	0.9953	0.9825	0.9631	0.9384	0.9098	0.8781	0.8442	0.8088	0.7725	0.7358
2	0.9998	0.9989	0.9964	0.9921	0.9856	0.9769	0.9659	0.9536	0.9371	0.9197
3	1.0000	0.9999	0.9997	0.9992	0.9982	0.9966	0.9942	0.9909	0.9865	0.9810
4	1.0000	1.0000	1.0000	0.9999	0.9998	0.9996	0.9992	0.9986	0.9977	0.9963
5	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9997	0.9994
6	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999
7	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
k	μ									
	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0
0	0.3329	0.3012	0.2725	0.2466	0.2231	0.2019	0.1827	0.1653	0.1496	0.1353
1	0.6990	0.6626	0.6268	0.5918	0.5578	0.5249	0.4932	0.4628	0.4337	0.4060
2	0.9004	0.8795	0.8571	0.8335	0.8088	0.7834	0.7572	0.7306	0.7037	0.6767
3	0.9743	0.9662	0.9569	0.9463	0.9344	0.9212	0.9068	0.8913	0.8747	0.8571
4	0.9946	0.9923	0.9893	0.9857	0.9814	0.9763	0.9704	0.9636	0.9559	0.9473
5	0.9990	0.9985	0.9978	0.9968	0.9955	0.9940	0.9920	0.9896	0.9868	0.9834
6	0.9999	0.9997	0.9996	0.9994	0.9991	0.9987	0.9981	0.9974	0.9966	0.9955
7	1.0000	1.0000	0.9999	0.9999	0.9998	0.9997	0.9996	0.9994	0.9992	0.9989
8	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9999	0.9998
9	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
k	μ									
	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0
0	0.1225	0.1108	0.1003	0.0907	0.0821	0.0743	0.0672	0.0608	0.0550	0.0498
1	0.3796	0.3546	0.3309	0.3084	0.2873	0.2674	0.2487	0.2311	0.2146	0.1991
2	0.6496	0.6227	0.5960	0.5697	0.5438	0.5184	0.4936	0.4695	0.4460	0.4232
3	0.8386	0.8194	0.7993	0.7787	0.7576	0.7360	0.7141	0.6919	0.6696	0.6472
4	0.9379	0.9275	0.9162	0.9041	0.8912	0.8774	0.8629	0.8477	0.8318	0.8153
5	0.9796	0.9751	0.9700	0.9643	0.9580	0.9510	0.9433	0.9349	0.9258	0.9161
6	0.9941	0.9925	0.9906	0.9884	0.9858	0.9828	0.9794	0.9756	0.9713	0.9665
7	0.9985	0.9980	0.9974	0.9967	0.9958	0.9947	0.9934	0.9919	0.9901	0.9881
8	0.9997	0.9995	0.9994	0.9991	0.9989	0.9985	0.9981	0.9976	0.9969	0.9962
9	0.9999	0.9999	0.9999	0.9998	0.9997	0.9996	0.9995	0.9993	0.9991	0.9989
10	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9999	0.9998	0.9997	0.9996
11	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9999
12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

Table for Poisson random variable $P(Po(\mu) \leq k)$.

k	μ									
	3.2	3.4	3.6	3.8	4.0	4.2	4.4	4.6	4.8	5.0
0	0.0408	0.0334	0.0273	0.0224	0.0183	0.0150	0.0123	0.0101	0.0082	0.0067
1	0.1712	0.1468	0.1257	0.1074	0.0916	0.0780	0.0663	0.0563	0.0477	0.0404
2	0.3799	0.3397	0.3027	0.2689	0.2381	0.2102	0.1851	0.1626	0.1425	0.1247
3	0.6025	0.5584	0.5152	0.4735	0.4335	0.3954	0.3594	0.3257	0.2942	0.2650
4	0.7806	0.7442	0.7064	0.6678	0.6288	0.5898	0.5512	0.5132	0.4763	0.4405
5	0.8946	0.8705	0.8441	0.8156	0.7851	0.7531	0.7199	0.6858	0.6510	0.6160
6	0.9534	0.9421	0.9267	0.9091	0.8893	0.8675	0.8436	0.8180	0.7908	0.7622
7	0.9832	0.9769	0.9682	0.9599	0.9509	0.9419	0.9321	0.9214	0.9099	0.8966
8	0.9943	0.9917	0.9883	0.9840	0.9786	0.9721	0.9642	0.9549	0.9442	0.9319
9	0.9982	0.9973	0.9960	0.9942	0.9919	0.9889	0.9851	0.9805	0.9749	0.9682
10	0.9995	0.9992	0.9987	0.9981	0.9972	0.9959	0.9943	0.9922	0.9896	0.9863
11	0.9999	0.9998	0.9996	0.9994	0.9991	0.9986	0.9980	0.9971	0.9960	0.9945
12	1.0000	0.9999	0.9999	0.9999	0.9998	0.9997	0.9996	0.9993	0.9990	0.9986
13	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9997	0.9995	0.9993
14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9999	0.9998
15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999
k	μ									
	5.2	5.4	5.6	5.8	6.0	6.5	7.0	7.5	8.0	8.5
0	0.0055	0.0045	0.0037	0.0030	0.0025	0.0015	0.0009	0.0006	0.0003	0.0002
1	0.0342	0.0289	0.0244	0.0206	0.0174	0.0113	0.0073	0.0047	0.0030	0.0019
2	0.1088	0.0948	0.0824	0.0715	0.0620	0.0430	0.0286	0.0203	0.0138	0.0093
3	0.2381	0.2133	0.1906	0.1700	0.1512	0.1118	0.0818	0.0591	0.0424	0.0301
4	0.4061	0.3733	0.3422	0.3127	0.2851	0.2237	0.1730	0.1321	0.0996	0.0744
5	0.5809	0.5461	0.5119	0.4783	0.4457	0.3690	0.3007	0.2414	0.1912	0.1496
6	0.7324	0.7017	0.6703	0.6384	0.6063	0.5265	0.4497	0.3782	0.3134	0.2562
7	0.8449	0.8217	0.7970	0.7710	0.7440	0.6728	0.5987	0.5246	0.4530	0.3856
8	0.9181	0.9027	0.8857	0.8672	0.8472	0.7916	0.7291	0.6620	0.5925	0.5231
9	0.9603	0.9512	0.9409	0.9292	0.9161	0.8774	0.8305	0.7764	0.7166	0.6530
10	0.9823	0.9775	0.9718	0.9651	0.9574	0.9332	0.9015	0.8622	0.8159	0.7634
11	0.9927	0.9904	0.9875	0.9841	0.9799	0.9661	0.9467	0.9208	0.8881	0.8487
12	0.9972	0.9962	0.9949	0.9932	0.9912	0.9840	0.9730	0.9573	0.9362	0.9091
13	0.9990	0.9986	0.9980	0.9973	0.9964	0.9929	0.9872	0.9784	0.9658	0.9486
14	0.9999	0.9995	0.9993	0.9990	0.9986	0.9970	0.9943	0.9897	0.9827	0.9726
15	0.9999	0.9998	0.9998	0.9996	0.9995	0.9988	0.9976	0.9954	0.9918	0.9862
16	1.0000	0.9999	0.9999	0.9999	0.9998	0.9996	0.9990	0.9980	0.9963	0.9934
17	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9996	0.9992	0.9984	0.9970
18	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9997	0.9993	0.9987
19	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9995
20	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998
21	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999
22	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

Table for Poisson random variable $P(Po(\mu) \leq k)$.

k	μ														
	9.0	9.5	10.0	11.0	12.0	13.0	14.0	15.0	16.0	17.0					
0	0.0001	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000					
1	0.0012	0.0008	0.0005	0.0002	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000					
2	0.0062	0.0042	0.0028	0.0012	0.0005	0.0002	0.0001	0.0000	0.0000	0.0000					
3	0.0212	0.0149	0.0103	0.0049	0.0023	0.0011	0.0005	0.0002	0.0001	0.0000					
4	0.0550	0.0403	0.0293	0.0151	0.0076	0.0037	0.0018	0.0009	0.0004	0.0002					
5	0.1157	0.0885	0.0671	0.0375	0.0203	0.0107	0.0055	0.0028	0.0014	0.0007					
6	0.2068	0.1649	0.1301	0.0786	0.0458	0.0259	0.0142	0.0076	0.0040	0.0021					
7	0.3239	0.2687	0.2202	0.1432	0.0895	0.0540	0.0316	0.0180	0.0100	0.0054					
8	0.4557	0.3918	0.3328	0.2320	0.1550	0.0998	0.0621	0.0374	0.0220	0.0126					
9	0.5874	0.5218	0.4579	0.3405	0.2424	0.1658	0.1094	0.0699	0.0433	0.0261					
10	0.7060	0.6453	0.5830	0.4599	0.3472	0.2517	0.1757	0.1185	0.0774	0.0491					
11	0.8030	0.7520	0.6968	0.5793	0.4616	0.3532	0.2600	0.1848	0.1270	0.0847					
12	0.8758	0.8364	0.7916	0.6887	0.5760	0.4631	0.3585	0.2676	0.1931	0.1350					
13	0.9261	0.8981	0.8645	0.7813	0.6815	0.5730	0.4644	0.3632	0.2745	0.2009					
14	0.9585	0.9400	0.9165	0.8540	0.7720	0.6751	0.5704	0.4657	0.3675	0.2808					
15	0.9780	0.9665	0.9513	0.9074	0.8444	0.7636	0.6694	0.5681	0.4667	0.3715					
16	0.9889	0.9823	0.9730	0.9441	0.8987	0.8355	0.7559	0.6641	0.5660	0.4677					
17	0.9947	0.9911	0.9857	0.9678	0.9370	0.8905	0.8272	0.7489	0.6593	0.5640					
18	0.9976	0.9957	0.9928	0.9823	0.9626	0.9302	0.8826	0.8195	0.7423	0.6550					
19	0.9989	0.9980	0.9965	0.9907	0.9787	0.9573	0.9235	0.8752	0.8122	0.7363					
20	0.9996	0.9991	0.9984	0.9953	0.9884	0.9750	0.9521	0.9170	0.8682	0.8055					
21	0.9998	0.9996	0.9993	0.9977	0.9939	0.9859	0.9712	0.9469	0.9108	0.8615					
22	0.9999	0.9999	0.9997	0.9990	0.9970	0.9924	0.9833	0.9673	0.9418	0.9047					
23	1.0000	0.9999	0.9999	0.9995	0.9985	0.9960	0.9907	0.9805	0.9633	0.9367					
24	1.0000	1.0000	1.0000	0.9998	0.9993	0.9980	0.9950	0.9888	0.9777	0.9594					
25	1.0000	1.0000	1.0000	0.9999	0.9997	0.9990	0.9974	0.9938	0.9869	0.9748					
26	1.0000	1.0000	1.0000	1.0000	0.9999	0.9995	0.9987	0.9967	0.9925	0.9848					
27	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9994	0.9983	0.9959	0.9912					
28	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9991	0.9978	0.9950					
29	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9996	0.9994	0.9986					
30	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9993					
31	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9996					
32	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9996					
33	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9999					
34	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999					
35	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000					