

Examiner: Xiangfeng Yang (Tel: 013-285788).

Things allowed (Hjälpmedel): a calculator.

Scores rating (Betygsgränser): 8-11 points giving rate 3; 11.5-14.5 points giving rate 4; 15-18 points giving rate 5.

1 (3 points)

In a box, there are 12 white balls and 8 black balls. One randomly takes out 6 balls one by one, and observes the colors.

(1.1) (1p) If this is done without replacement (namely, once a ball is taken out and observed, it will be put outside of the box), then what is the probability that exactly 2 black balls are observed?

(1.2) (2p) If this is done with replacement (namely, after a ball is taken out and observed, it will be put back into the box), then what is the probability that exactly 2 black balls are observed?

Solution. (1.1) This is exactly hypergeometric random variable, namely,

$$P(\text{exactly 2 black}) = \frac{\binom{8}{2} \cdot \binom{12}{4}}{\binom{20}{6}} = 0.3576.$$

(1.2) Let X be the number of black balls in these 6 chosen balls, then $X \sim \text{Bin}(6, 8/(12+8)) = \text{Bin}(6, 0.4)$.

$$P(\text{exactly 2 black}) = P(X = 2) = P(X \leq 2) - P(X \leq 1) = (\text{Binomial table}) = 0.5443 - 0.2333 = 0.311.$$

□

2 (3 points)

The distribution X of the numbers of cars of all households in Linköping is as follows

X	0	1	2	3
$p(x)$	20%	50%	25%	5%

The above tables says that each randomly chosen household in Linköping may have 0, 1, 2 or 3 cars with the corresponding probabilities. Now one randomly chooses 200 households in Linköping, what is the probability that the average number of cars of these 200 households is less than or equal to 1?

Solution. Let X_i denote the number of cars of the i -th household for $i = 1, 2, \dots, 200$, then we want to find the probability $P(\bar{X} \leq 1)$ where $\bar{X} = \frac{X_1 + X_2 + \dots + X_{200}}{200}$. To achieve this, one needs to find individual (population) mean μ and variance σ^2 as follows (as a discrete random variable give by the above table)

$$\mu = 0 \cdot 20\% + 1 \cdot 50\% + 2 \cdot 25\% + 3 \cdot 5\% = 1.15, \quad \sigma^2 = [0^2 \cdot 20\% + 1^2 \cdot 50\% + 2^2 \cdot 25\% + 3^2 \cdot 5\%] - 1.15^2 = 1.95 - 1.3225 = 0.6275.$$

Then it is from CLT that

$$\begin{aligned} P(\bar{X} \leq 1) &= P\left(\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \leq \frac{1 - \mu}{\sigma/\sqrt{n}}\right) = P\left(N(0, 1) \leq \frac{1 - 1.15}{\sqrt{0.6275}/\sqrt{200}}\right) = P(N(0, 1) \leq -2.68) \\ &= \Phi(-2.68) = 1 - \Phi(2.68) = 1 - 0.9963 = 0.0037. \end{aligned}$$

□

3 (3 points)

Two fair dice are tossed one by one independently. Let X_1 be the upper side number of the first die, X_2 be the upper side number of the second die and X_3 be the sum of the upper side numbers of the two dice.

(3.1) (1p) Are the two events $\{X_1 = 4\}$ and $\{X_3 = 6\}$ independent? Why?

(3.2) (1p) Are the two events $\{X_1 = 4\}$ and $\{X_3 = 7\}$ independent? Why?

(3.3) (1p) Find the conditional probability $P(X_3 = 5 \mid X_2 = 2)$.

Solution. (3.1) Not independent! It is from the fact that X_1 and X_2 are independent (since two dice are independent)

$$\begin{aligned}P(X_1 = 4, X_3 = 6) &= P(X_1 = 4, X_2 = 2) = P(X_1 = 4) \cdot P(X_2 = 2) = \frac{1}{6} \cdot \frac{1}{6} = \frac{1}{36}, \\P(X_1 = 4) &= \frac{1}{6}, \\P(X_3 = 6) &= P((1, 5), (5, 1), (2, 4), (4, 2), (3, 3)) = \frac{5}{36}.\end{aligned}$$

It is clear that $P(X_1 = 4, X_3 = 6) \neq P(X_1 = 4) \cdot P(X_3 = 6)$.

(3.2) Independent!

$$\begin{aligned}P(X_1 = 4, X_3 = 7) &= P(X_1 = 4, X_2 = 3) = P(X_1 = 4) \cdot P(X_2 = 3) = \frac{1}{6} \cdot \frac{1}{6} = \frac{1}{36}, \\P(X_1 = 4) &= \frac{1}{6}, \\P(X_3 = 7) &= P((1, 6), (6, 1), (2, 5), (5, 2), (3, 4), (4, 3)) = \frac{6}{36} = \frac{1}{6}.\end{aligned}$$

It is clear that $P(X_1 = 4, X_3 = 7) = P(X_1 = 4) \cdot P(X_3 = 7)$.

(3.3)

$$P(X_3 = 5 \mid X_2 = 2) = \frac{P(X_3 = 5, X_2 = 2)}{P(X_2 = 2)} = \frac{P(X_1 = 3, X_2 = 2)}{P(X_2 = 2)} = \frac{P(X_1 = 3) \cdot P(X_2 = 2)}{P(X_2 = 2)} = P(X_1 = 3) = \frac{1}{6}.$$

□

4 (3 points)

A population $X \sim \text{Bin}(3, p)$ is a Binomial random variable with an unknown parameter p . A sample $\{2, 1, 1, 0\}$ is taken from this population.

(4.1) (1p) Use the Method of Moments to find a point estimate \hat{p}_{MM} of p .

(4.2) (2p) Use the Maximum Likelihood Method to find a point estimate \hat{p}_{ML} of p .

Solution. (4.1) It is from the method of moments that $E(X) = \bar{x}$ where $E(X) = 3 \cdot p$, which implies that

$$3 \cdot p = \bar{x} \Rightarrow \hat{p}_{MM} = \bar{x}/3 = 1/3 = 0.3333.$$

(4.2) The likelihood function is

$$\begin{aligned}L(p) &= p(x_1) \cdot \dots \cdot p(x_4) = \left[\binom{3}{x_1} p^{x_1} \cdot (1-p)^{3-x_1} \right] \cdot \dots \cdot \left[\binom{3}{x_4} p^{x_4} \cdot (1-p)^{3-x_4} \right] \\&= \left[\binom{3}{x_1} \dots \binom{3}{x_4} \right] \cdot p^{x_1 + \dots + x_4} \cdot (1-p)^{12 - (x_1 + \dots + x_4)}.\end{aligned}$$

The log likelihood function is $\ln L(p) = \ln \left[\binom{3}{x_1} \dots \binom{3}{x_4} \right] + (x_1 + \dots + x_4) \ln p + (12 - (x_1 + \dots + x_4)) \ln(1-p)$.

Taking derivative yields

$$0 = \ln' L(p) = \frac{x_1 + \dots + x_4}{p} - \frac{12 - (x_1 + \dots + x_4)}{1-p} \Rightarrow \hat{p}_{ML} = \bar{x}/3 = 1/3 = 0.3333.$$

□

5 (3 points)

There are two equally popular restaurants in Linköping (marked as A and B). “Equally popular” means that the preference of each restaurant is $\begin{bmatrix} A & B \\ 50\% & 50\% \end{bmatrix}$. This month the restaurant A has employed a new chef, and the aim is to increase the restaurant’s preference. Now a sample of 100 people have been asked for their preference with the following results $\begin{bmatrix} A & B \\ 53\% & 47\% \end{bmatrix}$. Based on the sample, has the new chef really changed the preference of the restaurants? Answer this question using a χ^2 -test with a level $\alpha = 5\%$.

Solution. Here we want to test the hypothesis:

$$H_0 : P(A) = 50\%, P(B) = 50\% \text{ vs. } H_a : \text{some is not true in } H_0$$

The observed value of the test statistic under H_0 is

$$TS = \sum_{i=1}^2 \frac{(n_i - np_i)^2}{np_i} = \frac{(53 - 100 \cdot 50\%)^2}{100 \cdot 50\%} + \frac{(47 - 100 \cdot 50\%)^2}{100 \cdot 50\%} = \frac{18}{50} = 0.36.$$

The rejection region is

$$C = (\chi_{\alpha}^2(k-1), \infty) = (\chi_{0.05}^2(1), \infty) = (3.84, \infty).$$

As $TS \notin C$, H_0 is not rejected (the sample does not provide any evidence that the new chef has really changed the preference). □

6 (3 points)

A professor at LiU claims that the heights of students at LiU differ a lot. To conduct a statistical study on this, let us assume that the distribution X of the heights of students at LiU is normal $X \sim N(\mu, \sigma^2)$ (with μ denoting the average height and σ^2 the variation of the heights). A sample is taken from X with sample size $n = 20$, sample mean $\bar{x} = 178$ and sample variance $s^2 = 15.6$. Does the sample provide any evidence that $\sigma^2 > 10$ with a confidence 95%? Answer this question by using an appropriate one-sided 95% confidence interval (Do NOT use hypothesis test).

Solution. Because here we want to check whether or not $\sigma^2 > 10$, an one-sided 95% confidence interval $I_{\sigma^2} = (a, \infty)$ should be constructed (and then one needs to compare a with 10). To this end, one-sided 95% confidence interval is

$$I_{\sigma^2} = \left(\frac{(n-1)s^2}{\chi_{\alpha}^2(n-1)}, \infty \right) = \left(\frac{(20-1)15.6}{\chi_{0.05}^2(20-1)}, \infty \right) = \left(\frac{296.4}{30.14}, \infty \right) = (9.834, \infty).$$

As $9.834 \not> 10$, the sample does NOT provide any evidence that $\sigma^2 > 10$. □

1. Basic probability

- (1.1) Conditional probability $P(A|B) = \frac{P(A \cap B)}{P(B)}$.
 (1.2) Total probability $P(B) = \sum_{i=1}^k P(B|A_i)P(A_i)$ where $\{A_i\}$ are disjoint and $\cup_{i=1}^k A_i = S$.
 (1.3) Bayes' Theorem $P(A_i|B) = \frac{P(B|A_i)P(A_i)}{\sum_{i=1}^k P(B|A_i)P(A_i)}$ where $\{A_i\}$ are in (1.2).

2. Random variables (r.v.s)

- (2.1) Discrete r.v. X has a pmf $p(x) = P(X = x)$ satisfying $p(x) \geq 0$ and $\sum p(x_i) = 1$,

$$\begin{array}{c|cccc} X & x_1 & x_2 & \cdots & x_n & \cdots \\ \hline p(x) & p(x_1) & p(x_2) & \cdots & p(x_n) & \cdots \end{array}$$

- Expectation (or *Expected value* or *mean*) $\mu_X = E(X) = \sum x_i p(x_i)$;
 Variance $\sigma_X^2 = V(X) = E(X - \mu_X)^2 = E(X^2) - \mu_X^2 = \sum x_i^2 p(x_i) - (\sum x_i p(x_i))^2$.
 (2.2) Continuous r.v. X has a pdf $f(x)$ satisfying $f(x) \geq 0$ and $\int_{-\infty}^{\infty} f(x) dx = 1$,

$$P(a < X < b) = \int_a^b f(x) dx.$$

Expectation (or *Expected value* or *mean*) $\mu_X = E(X) = \int_{-\infty}^{\infty} x f(x) dx$;

Variance $\sigma_X^2 = V(X) = E(X - \mu_X)^2 = E(X^2) - \mu_X^2 = \int_{-\infty}^{\infty} x^2 f(x) dx - (\int_{-\infty}^{\infty} x f(x) dx)^2$.

- (2.3) Cumulative distribution function (cdf) of a r.v. X is $F(x) = P(X \leq x)$.

- (2.4) X and Y are r.v.s, a, b and c are scalars, then

$$E(aX + bY + c) = aE(X) + bE(Y) + c,$$

$$V(aX + bY + c) = a^2 V(X) + b^2 V(Y) + 2ab \operatorname{cov}(X, Y),$$

$$E(g(X, Y)) = \begin{cases} \sum_{i,j} g(x_i, y_j) \cdot p(x_i, y_j), & \text{for discrete } (X, Y), \\ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) \cdot f(x, y) dx dy, & \text{for continuous } (X, Y). \end{cases}$$

- (2.5) • Discrete r.v. (X, Y) has a joint pmf $p(x, y)$ satisfying $p(x, y) \geq 0$ and $\sum_{x_i} \sum_{y_j} p(x_i, y_j) = 1$.

The *marginal pmf* of X is $p_X(x) = \sum_y p(x, y)$;

The *marginal pmf* of Y is $p_Y(y) = \sum_x p(x, y)$;

X and Y are *independent* if $p(x, y) = p_X(x) \cdot p_Y(y)$.

- Continuous r.v. (X, Y) has a joint pdf $p(x, y)$ satisfying $f(x, y) \geq 0$ and $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) dx dy = 1$.

The *marginal pdf* of X is $f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy$;

The *marginal pdf* of Y is $f_Y(y) = \int_{-\infty}^{\infty} f(x, y) dx$;

X and Y are *independent* if $f(x, y) = f_X(x) \cdot f_Y(y)$.

3. Several special r.v.s

- (3.1) $X \sim \operatorname{Bin}(n, p)$ has a pmf $p(x) = P(X = x) = \binom{n}{x} \cdot p^x \cdot (1-p)^{n-x}$, $x = 0, 1, 2, \dots, n$.

$$E(X) = n \cdot p, \quad V(X) = n \cdot p \cdot (1-p).$$

- (3.2) $X \sim \operatorname{Po}(\lambda)$ has a pmf $p(x) = P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$, $x = 0, 1, 2, \dots$
 $E(X) = \lambda, \quad V(X) = \lambda$.

- (3.3) $X \sim \operatorname{Hypergeometric}$ has a pmf $p(x) = P(X = x) = \frac{\binom{M}{x} \binom{N-M}{n-x}}{\binom{N}{n}}$.

- (3.4) $X \sim \operatorname{Exp}(\lambda)$ has a pdf

$$f(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0, \\ 0, & \text{otherwise.} \end{cases}$$

- (3.5) $X \sim N(\mu, \sigma^2)$ has a pdf

$$f(x) = \frac{1}{\sigma \sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad -\infty < x < \infty.$$

- (3.6) $X \sim U(a, b)$ has a pdf

$$f(x) = \begin{cases} \frac{1}{b-a}, & a < x < b, \\ 0, & \text{otherwise.} \end{cases}$$

$$E(X) = \frac{a+b}{2}, \quad V(X) = \frac{(b-a)^2}{12}.$$

4. Central Limit Theorem (CLT)

Suppose that a population has mean $= \mu$ and variance $= \sigma^2$. A random sample $\{X_1, X_2, \dots, X_n\}$ from this population is given. Then for large $n \geq 30$,

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1). \tag{1}$$

- If the population is normal, then (1) holds for any n .
- Note that $\mu = E(\bar{X})$ and $(\sigma/\sqrt{n})^2 = V(\bar{X})$.

5. Several notations in statistics

- (5.1) Sample mean: $\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n} = \sum \bar{X}_i$; $\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{\sum x_i}{n}$.

- (5.2) Sample variance:

$$S^2 = \frac{\sum (X_i - \bar{X})^2}{n-1} = \frac{1}{n-1} \left(\sum X_i^2 - \frac{(\sum X_i)^2}{n} \right); \quad s^2 = \frac{\sum (x_i - \bar{x})^2}{n-1} = \frac{1}{n-1} \left(\sum x_i^2 - \frac{(\sum x_i)^2}{n} \right).$$

- Capital letters \bar{X} and S^2 refer to the objects based on random sample (therefore they are in general r.v.s), while small letters \bar{x} and s^2 are the objects based on observations (so they are scalars).

- (5.3) A point estimator of θ obtained by Methods of Moments is denoted as $\hat{\theta}_{MM}$.

- (5.4) A point estimator of θ obtained by Maximum Likelihood method is denoted as $\hat{\theta}_{ML}$.

6. Confidence Interval (CI)

In this course, three types of confidence intervals are studied depending on the unknown population parameter(s): CI-1 (confidence intervals for population mean(s)), CI-2 (confidence intervals for population variance(s)), and CI-3 (confidence intervals for population proportion(s)).

CI-1: (1 - α) CI of a population mean μ

case 1.1 (any n) If population $X \sim N(\mu, \sigma^2)$ and σ^2 is known, then $\frac{\bar{X}-\mu}{\sigma/\sqrt{n}} \sim N(0, 1)$ and

$$I_\mu = (\bar{x} - z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}, \bar{x} + z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}) := \bar{x} \mp z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}.$$

case 1.2 (n ≥ 30) For any population X , it holds that $\frac{\bar{X}-\mu}{\sigma/\sqrt{n}} \sim N(0, 1)$ and

$$I_\mu = \bar{x} \mp z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \text{ or } I_\mu = \bar{x} \mp z_{\alpha/2} \cdot \frac{\hat{\sigma}}{\sqrt{n}}.$$

case 1.3 (any n) If population $X \sim N(\mu, \sigma^2)$ and σ^2 is unknown, then $\frac{\bar{X}-\mu}{S/\sqrt{n}} \sim T(n-1)$ and

$$I_\mu = \bar{x} \mp t_{\alpha/2}(n-1) \cdot \frac{s}{\sqrt{n}}.$$

CI-1': (1 - α) CI of the difference of two population means $\mu_X - \mu_Y$

case 1.1' (any n_1, n_2) If independent populations $X \sim N(\mu_X, \sigma_X^2)$, $Y \sim N(\mu_Y, \sigma_Y^2)$, and σ_X^2, σ_Y^2 are known, then

$$\frac{(\bar{X}-\bar{Y})-(\mu_X-\mu_Y)}{\sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}} \sim N(0, 1), \text{ and } I_{\mu_X-\mu_Y} = (\bar{x}-\bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}.$$

case 1.2' ($n_1, n_2 \geq 30$) For any independent populations X and Y , it holds that



$$\frac{(\bar{X}-\bar{Y})-(\mu_X-\mu_Y)}{\sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}} \sim N(0, 1) \text{ and}$$

$$I_{\mu_X-\mu_Y} = (\bar{x}-\bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}} \text{ or } I_{\mu_X-\mu_Y} = (\bar{x}-\bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{\sigma}_X^2}{n_1} + \frac{\hat{\sigma}_Y^2}{n_2}}.$$

case 1.3' (any n_1, n_2) If independent populations $X \sim N(\mu_X, \sigma_X^2)$, $Y \sim N(\mu_Y, \sigma_Y^2)$, where σ_X^2, σ_Y^2 are unknown but $\sigma_X^2 = \sigma_Y^2$, then

$$\frac{(\bar{X}-\bar{Y})-(\mu_X-\mu_Y)}{S\sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} \sim T(n_1+n_2-2), \text{ where } S^2 = \frac{(n_1-1)S_X^2 + (n_2-1)S_Y^2}{n_1+n_2-2}, \text{ and}$$

$$I_{\mu_X-\mu_Y} = (\bar{x}-\bar{y}) \mp t_{\alpha/2}(n_1+n_2-2) \cdot s\sqrt{\frac{1}{n_1} + \frac{1}{n_2}}.$$



CI-2: (1 - α) CI of population variance(s) σ^2


• If a population $X \sim N(\mu, \sigma^2)$ and σ^2 is unknown, then $\frac{(n-1)S^2}{\sigma^2} \sim \chi^2(n-1)$, and

$$I_{\sigma^2} = \left(\frac{(n-1)s^2}{\chi_{\alpha/2}^2(n-1)}, \frac{(n-1)s^2}{\chi_{1-\alpha/2}^2(n-1)} \right).$$

• If two independent populations $X \sim N(\mu_X, \sigma^2)$ and $Y \sim N(\mu_Y, \sigma^2)$, and σ^2 is unknown, then $\frac{(n_1+n_2-2)S^2}{\sigma^2} \sim \chi^2(n_1+n_2-2)$, and

$$I_{\sigma^2} = \left(\frac{(n_1+n_2-2)s^2}{\chi_{\alpha/2}^2(n_1+n_2-2)}, \frac{(n_1+n_2-2)s^2}{\chi_{1-\alpha/2}^2(n_1+n_2-2)} \right),$$

where $S^2 = \frac{(n_1-1)S_X^2 + (n_2-1)S_Y^2}{n_1+n_2-2}$.



CI-3: (1 - α) CI of population proportion(s)

• If a (large) population has an unknown proportion p , then $\frac{\hat{p}-p}{\sqrt{p(1-p)/n}} \sim N(0, 1)$ if $n\hat{p} \geq 10$ and $n(1-\hat{p}) \geq 10$ with $\hat{p} = x/n$, and $I_p = \hat{p} \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$.

• If two independent (large) populations have unknown proportions p_1 and p_2 , then

$$\frac{(\hat{p}_1-\hat{p}_2)-(p_1-p_2)}{\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{\hat{p}_2(1-\hat{p}_2)}{n_2}}} \sim N(0, 1)$$

if $n_i\hat{p}_i \geq 10$ and $n_i(1-\hat{p}_i) \geq 10$ for $i = 1, 2$, and $I_{p_1-p_2} = (\hat{p}_1-\hat{p}_2) \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{\hat{p}_2(1-\hat{p}_2)}{n_2}}$.

7. Hypothesis Test (HT)

	H_0 is true	H_0 is false and $\theta = \theta_1$
reject H_0	(type I error or significance level) α	(power) $h(\theta_1)$
don't reject H_0	$1 - \alpha$	(type II error) $\beta(\theta_1) = 1 - h(\theta_1)$

reject $H_0 \Leftrightarrow TS \in C \Leftrightarrow p\text{-value} < \alpha$

χ^2 tests for populations (non-parametric tests)

Suppose that for a random sample of a population X , the n elements of it are classified into k disjoint groups $A_i, 1 \leq i \leq k$. For each group $A_i, 1 \leq i \leq k$, suppose that there are $N_i, 1 \leq i \leq k$ elements inside. Let $p_i = P(A_i)$ assuming a given distribution of X . Note that $p_1 + p_2 + \dots + p_k = 1$ and $N_1 + N_2 + \dots + N_k = n$. One wants to test the hypotheses

$$H_0 : P(A_i) = p_i, \quad 1 \leq i \leq k, \quad H_a : P(A_i) \neq p_i \text{ for some } 1 \leq i \leq k.$$

If n is large in the sense that $np_i \geq 5$ for all $1 \leq i \leq k$, then the test statistic is

$$\sum_{i=1}^k \frac{(N_i - np_i)^2}{np_i} \approx \chi^2(k-1).$$

Therefore the observation of the test statistic is

$$TS = \sum_{i=1}^k \frac{(n_i - np_i)^2}{np_i}, \text{ where } n_i \text{ is the observation of } N_i, 1 \leq i \leq k.$$

For the critical region C , one can take (note that if H_0 is true, then TS should be close to zero)

$$C = (\chi^2_{\alpha}(k-1), \infty).$$

The conclusion would be $TS \in C \iff H_0$ is rejected.

8. Linear and logistic regression

(Multiple) linear regression: $Y = \beta_0 + \beta_1 x_1 + \dots + \beta_k x_k + \varepsilon$, $\varepsilon \sim N(0, \sigma^2)$.

- Y : response variable (which is normal r.v.), $\{x_1, \dots, x_k\}$: predictors (which are scalars).
- sample: $\{(x_{11}, \dots, x_{1k}; y_1), (x_{21}, \dots, x_{2k}; y_2), \dots, (x_{n1}, \dots, x_{nk}; y_n)\}$.
- how to estimate $\beta_j \approx \hat{\beta}_j$: least square method, that is, to minimize $\sum_{i=1}^n (y_i - \hat{y}_i)^2$, where the estimated (multiple) linear regression line \hat{y} is

$$\hat{y} = \hat{\beta}_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k.$$

- $\frac{\hat{\beta}_j - \beta_j}{se(\hat{\beta}_j)} \sim T(n-k-1)$, this helps determine whether or not the real $\beta_j = 0$?
- $\sigma^2 \approx \frac{SSE}{n-k-1}$, this gives an estimation of the size of the error.
- $R^2 = \frac{SSR}{SSY}$, this gives how well the model is (if $R^2 \approx 1$, then the model fits the sample very well).
- How to test $\beta_1 = \dots = \beta_k = 0$? Use the random variable $\frac{SSR/k}{SSE/(n-k-1)} \sim F(k, n-k-1)$.

Logistic regression: Let Y can only take 0 or 1 with $P(Y=1) = p$ and $P(Y=0) = 1-p$.

$$E(Y) = p(x_1, \dots, x_k) = \frac{e^{\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k}}{1 + e^{\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k}}.$$

- Y : response variable (which is Bernoulli r.v. $P(Y=1) = p$ and $P(Y=0) = 1-p$, so $E(Y) = p$), $\{x_1, \dots, x_k\}$: predictors (which are scalars).
- sample: $\{(x_{11}, \dots, x_{1k}; y_1), (x_{21}, \dots, x_{2k}; y_2), \dots, (x_{n1}, \dots, x_{nk}; y_n)\}$.
- how to estimate $\beta_j \approx \hat{\beta}_j$: maximal likelihood method (maximize $\prod_{i=1}^n p(x_{i1}, \dots, x_{ik})^{y_i} (1 - p(x_{i1}, \dots, x_{ik}))^{1-y_i}$).
- $\frac{\hat{\beta}_j - \beta_j}{se(\hat{\beta}_j)} \approx N(0, 1)$ for large $n \geq 30$, this helps determine whether or not the real $\beta_j = 0$?
- Classification of a new object $Y(x_1, \dots, x_k)$ as 1 or 0 according

$$Y(x_1, \dots, x_k) = \begin{cases} 1, & \text{if } \hat{p}(x_1, \dots, x_k) \geq 0.5, \\ 0, & \text{if } \hat{p}(x_1, \dots, x_k) < 0.5, \end{cases}$$

where the estimated logit function $\hat{p}(x_1, \dots, x_k)$ is

$$\hat{p}(x_1, \dots, x_k) = \frac{e^{\hat{\beta}_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k}}{1 + e^{\hat{\beta}_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k}}.$$

9. Tables

(9.1) Table for $N(0, 1)$ standard normal random variable $\Phi(x) = P(N(0, 1) \leq x)$, $x \geq 0$.
There is an important relation $\Phi(-x) = 1 - \Phi(x)$, $x \geq 0$.

x	0	1	2	3	4	5	6	7	8	9
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9564	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9992	0.9992	0.9993	0.9993
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997	0.9997
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998	0.9998
3.5	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998
3.6	0.9998	0.9998	0.9998	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.7	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.8	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.9	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
4.0	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

(9.2) Table for $T(f)$ random variable $F(x) = P(T(f) \leq x)$,
where f is a parameter called 'degrees of freedom'.

f	0.75	0.90	0.95	0.975	0.99	0.995	0.9975	0.9995
1	1.00	3.08	6.31	12.71	31.82	63.66	127.32	636.62
2	0.82	1.89	2.92	4.30	6.96	9.92	14.09	31.60
3	0.76	1.64	2.35	3.18	4.54	5.84	7.45	12.92
4	0.74	1.53	2.13	2.78	3.75	4.60	5.60	8.61
5	0.73	1.48	2.02	2.57	3.36	4.03	4.77	6.87
6	0.72	1.44	1.94	2.45	3.14	3.71	4.32	5.96
7	0.71	1.41	1.89	2.36	3.00	3.50	4.03	5.41
8	0.71	1.40	1.86	2.31	2.90	3.36	3.83	5.04
9	0.70	1.38	1.83	2.26	2.82	3.25	3.69	4.78
10	0.70	1.37	1.81	2.23	2.76	3.17	3.58	4.59
11	0.70	1.36	1.80	2.20	2.72	3.11	3.50	4.44
12	0.70	1.36	1.78	2.18	2.68	3.05	3.43	4.32
13	0.69	1.35	1.77	2.16	2.65	3.01	3.37	4.22
14	0.69	1.35	1.76	2.14	2.62	2.98	3.33	4.14
15	0.69	1.34	1.75	2.13	2.60	2.95	3.29	4.07
16	0.69	1.34	1.75	2.12	2.58	2.92	3.25	4.01
17	0.69	1.33	1.74	2.11	2.57	2.90	3.22	3.97
18	0.69	1.33	1.73	2.10	2.55	2.88	3.20	3.92
19	0.69	1.33	1.73	2.09	2.54	2.86	3.17	3.88
20	0.69	1.33	1.72	2.09	2.53	2.85	3.15	3.85
21	0.69	1.32	1.72	2.08	2.52	2.83	3.14	3.82
22	0.69	1.32	1.72	2.07	2.51	2.82	3.12	3.79
23	0.69	1.32	1.71	2.07	2.50	2.81	3.10	3.77
24	0.68	1.32	1.71	2.06	2.49	2.80	3.09	3.75
25	0.68	1.32	1.71	2.06	2.49	2.79	3.08	3.73
26	0.68	1.31	1.71	2.06	2.48	2.78	3.07	3.71
27	0.68	1.31	1.70	2.05	2.47	2.77	3.06	3.69
28	0.68	1.31	1.70	2.05	2.47	2.76	3.05	3.67
29	0.68	1.31	1.70	2.05	2.46	2.76	3.04	3.66
30	0.68	1.31	1.70	2.04	2.46	2.75	3.03	3.65
40	0.68	1.30	1.68	2.02	2.42	2.70	2.97	3.55
50	0.68	1.30	1.68	2.01	2.40	2.68	2.94	3.50
60	0.68	1.30	1.67	2.00	2.39	2.66	2.91	3.46
100	0.68	1.29	1.66	1.98	2.36	2.63	2.87	3.39
∞	0.67	1.28	1.65	1.96	2.33	2.58	2.81	3.29

(9.3) Table for $\chi^2(f)$ random variable $F(x) = P(\chi^2(f) \leq x)$, where f is a parameter.

f	0.0005	0.001	0.005	0.01	0.025	0.05	0.10	0.20	0.30	0.40	0.50
1	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.06	0.15	0.27	0.45
2	0.00	0.00	0.01	0.02	0.05	0.10	0.21	0.45	0.71	1.02	1.39
3	0.02	0.02	0.07	0.11	0.22	0.35	0.58	1.01	1.42	1.87	2.37
4	0.06	0.09	0.21	0.30	0.48	0.71	1.06	1.65	2.19	2.75	3.36
5	0.16	0.21	0.41	0.55	0.83	1.15	1.61	2.34	3.00	3.66	4.35
6	0.30	0.38	0.68	0.87	1.24	1.64	2.20	3.07	3.83	4.57	5.35
7	0.48	0.60	0.99	1.24	1.69	2.17	2.73	3.62	4.67	5.49	6.35
8	0.71	0.86	1.34	1.65	2.18	2.73	3.49	4.59	5.53	6.42	7.34
9	0.97	1.15	1.73	2.09	2.70	3.33	4.17	5.38	6.39	7.36	8.34
10	1.26	1.48	2.16	2.56	3.25	3.94	4.87	6.18	7.27	8.30	9.34
11	1.59	1.83	2.60	3.05	3.82	4.57	5.58	6.99	8.15	9.24	10.34
12	1.93	2.21	3.07	3.57	4.40	5.23	6.30	7.81	9.03	10.18	11.34
13	2.31	2.62	3.57	4.11	5.01	5.89	7.04	8.63	9.93	11.13	12.34
14	2.70	3.04	4.07	4.66	5.63	6.57	7.79	9.47	10.82	12.08	13.34
15	3.11	3.48	4.60	5.23	6.26	7.26	8.55	10.31	11.72	13.03	14.34
16	3.54	3.94	5.14	5.81	6.91	7.96	9.31	11.15	12.62	13.98	15.34
17	3.98	4.42	5.70	6.41	7.56	8.67	10.09	12.00	13.53	14.94	16.34
18	4.44	4.90	6.26	7.01	8.23	9.39	10.86	12.86	14.44	15.89	17.34
19	4.91	5.41	6.84	7.63	8.91	10.12	11.65	13.72	15.35	16.85	18.34
20	5.40	5.92	7.43	8.26	9.59	10.85	12.44	14.58	16.27	17.81	19.34
21	5.90	6.45	8.03	8.90	10.28	11.59	13.24	15.44	17.18	18.77	20.34
22	6.40	6.98	8.64	9.54	10.98	12.34	14.04	16.31	18.10	19.73	21.34
23	6.92	7.53	9.26	10.20	11.69	13.09	14.85	17.19	19.02	20.69	22.34
24	7.45	8.08	9.89	10.86	12.40	13.85	15.66	18.06	19.94	21.65	23.34
25	7.99	8.65	10.52	11.52	13.12	14.61	16.47	18.94	20.87	22.62	24.34
26	8.54	9.22	11.16	12.20	13.84	15.38	17.29	19.82	21.79	23.58	25.34
27	9.09	9.80	11.81	12.88	14.57	16.15	18.11	20.70	22.72	24.54	26.34
28	9.66	10.39	12.46	13.56	15.31	16.93	18.94	21.59	23.65	25.51	27.34
29	10.23	10.99	13.12	14.26	16.05	17.71	19.77	22.48	24.58	26.48	28.34
30	10.80	11.59	13.79	14.95	16.79	18.49	20.60	23.36	25.51	27.44	29.34
40	16.91	17.92	20.71	22.16	24.43	26.51	29.05	32.34	34.87	37.13	39.34
50	23.46	24.67	27.99	29.71	32.36	34.76	37.69	41.45	46.86	49.33	51.33
60	30.34	31.74	35.53	37.48	40.48	43.19	46.46	50.64	53.81	56.62	59.33
100	59.90	61.92	67.33	70.06	74.22	77.93	82.36	87.95	92.13	95.81	99.33

Table for $\chi^2(f)$ random variable $F(x) = P(\chi^2(f) \leq x)$, where f is a parameter.

f	0.60	0.70	0.80	0.90	0.95	0.975	0.99	0.995	0.999	0.9995
1	0.71	1.07	1.64	2.71	3.84	5.02	6.63	7.88	10.83	12.12
2	1.83	2.41	3.22	4.61	5.99	7.38	9.21	10.60	13.82	15.20
3	2.95	3.66	4.64	6.25	7.81	9.35	11.34	12.84	16.27	17.73
4	4.04	4.88	5.99	7.78	9.49	11.14	13.28	14.86	18.47	20.00
5	5.13	6.06	7.29	9.24	11.07	12.83	15.09	16.75	20.52	22.11
6	6.21	7.23	8.56	10.64	12.59	14.45	16.81	18.55	22.46	24.10
7	7.28	8.38	9.80	12.02	14.07	16.01	18.48	20.28	24.32	26.02
8	8.35	9.52	11.03	13.36	15.51	17.53	20.09	21.95	26.12	27.87
9	9.41	10.66	12.24	14.68	16.92	19.02	21.67	23.59	27.88	29.67
10	10.47	11.78	13.44	15.99	18.31	20.48	23.21	25.19	29.59	31.42
11	11.53	12.90	14.63	17.28	19.68	21.92	24.72	26.76	31.26	33.14
12	12.58	14.01	15.81	18.55	21.03	23.34	26.22	28.30	32.91	34.82
13	13.64	15.12	16.98	19.81	22.36	24.74	27.69	29.82	34.53	36.48
14	14.69	16.22	18.15	21.06	23.68	26.12	29.14	31.32	36.12	38.11
15	15.73	17.32	19.31	22.31	25.00	27.49	30.58	32.80	37.70	39.72
16	16.78	18.42	20.47	23.54	26.30	28.85	32.00	34.27	39.25	41.31
17	17.82	19.51	21.61	24.77	27.59	30.19	33.41	35.72	40.79	42.88
18	18.87	20.60	22.76	25.99	28.87	31.53	34.81	37.16	42.31	44.43
19	19.91	21.69	23.90	27.20	30.14	32.85	36.19	38.58	43.82	45.97
20	20.95	22.77	25.04	28.41	31.41	34.17	37.57	40.00	45.31	47.50
21	21.99	23.86	26.17	29.62	32.67	35.48	38.93	41.40	46.80	49.01
22	23.03	24.94	27.30	30.81	33.92	36.78	40.29	42.80	48.27	50.51
23	24.07	26.02	28.43	32.01	35.17	38.08	41.64	44.18	49.73	52.00
24	25.11	27.10	29.55	33.20	36.42	39.36	42.98	45.56	51.18	53.48
25	26.14	28.17	30.68	34.38	37.65	40.65	44.31	46.93	52.62	54.95
26	27.18	29.25	31.79	35.56	38.89	41.92	45.64	48.29	54.05	56.41
27	28.21	30.32	32.91	36.74	40.11	43.19	46.96	49.64	55.48	57.86
28	29.25	31.39	34.03	37.92	41.34	44.46	48.28	50.99	56.89	59.30
29	30.28	32.46	35.14	39.09	42.56	45.72	49.59	52.34	58.30	60.73
30	31.32	33.53	36.25	40.26	43.77	46.98	50.89	53.67	59.70	62.16
40	41.62	44.16	47.27	51.81	55.76	59.34	63.69	66.77	73.40	76.09
50	51.89	54.72	58.16	63.17	67.50	71.42	76.15	79.49	86.66	89.56
60	62.13	65.23	68.97	74.40	79.08	83.30	88.38	91.95	99.61	102.69
100	102.95	106.91	111.67	118.50	124.34	129.56	135.81	140.17	149.45	153.17

(9.4) Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
 If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1 - p) \geq n - k)$.

n	k	0.05	0.10	0.15	0.20	0.25	0.30	0.35	0.40	0.45	0.50
2	0	0.9025	0.8100	0.7225	0.6400	0.5625	0.4900	0.4225	0.3600	0.3025	0.2500
	1	0.9975	0.9900	0.9775	0.9600	0.9375	0.9100	0.8775	0.8400	0.7975	0.7500
3	0	0.8574	0.7290	0.6141	0.5120	0.4219	0.3430	0.2746	0.2160	0.1664	0.1250
	1	0.9928	0.9720	0.9392	0.8960	0.8438	0.7840	0.7183	0.6480	0.5747	0.5000
4	0	0.8145	0.6561	0.5220	0.4096	0.3164	0.2401	0.1785	0.1256	0.0915	0.0625
	1	0.9860	0.9477	0.8905	0.8192	0.7383	0.6517	0.5630	0.4735	0.3910	0.3125
5	0	0.7738	0.5905	0.4437	0.3277	0.2373	0.1681	0.1160	0.0778	0.0503	0.0313
	1	0.9774	0.9185	0.8352	0.7373	0.6328	0.5282	0.4284	0.3370	0.2562	0.1875
6	0	0.7351	0.5314	0.3771	0.2621	0.1780	0.1176	0.0754	0.0467	0.0277	0.0156
	1	0.9672	0.8847	0.7765	0.6554	0.5339	0.4202	0.3191	0.2333	0.1636	0.1094
7	0	0.6983	0.4783	0.3206	0.2097	0.1335	0.0824	0.0490	0.0280	0.0152	0.0078
	1	0.9566	0.8503	0.7166	0.5767	0.4449	0.3294	0.2338	0.1586	0.1024	0.0625
8	0	0.6634	0.4305	0.2725	0.1678	0.1001	0.0576	0.0319	0.0168	0.0084	0.0039
	1	0.9428	0.8131	0.6572	0.5033	0.3671	0.2553	0.1691	0.1064	0.0632	0.0352
9	0	0.6302	0.3874	0.2316	0.1342	0.0751	0.0404	0.0207	0.0101	0.0046	0.0020
	1	0.9288	0.7748	0.5995	0.4362	0.3003	0.1960	0.1211	0.0705	0.0385	0.0195

Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
 If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1 - p) \geq n - k)$.

n	k	0.05	0.10	0.15	0.20	0.25	0.30	0.35	0.40	0.45	0.50
10	0	0.5987	0.3487	0.1969	0.1074	0.0563	0.0282	0.0135	0.0060	0.0025	0.0010
	1	0.9139	0.7361	0.5443	0.3758	0.2440	0.1493	0.0860	0.0464	0.0233	0.0107
11	0	0.5688	0.3138	0.1673	0.0859	0.0422	0.0198	0.0088	0.0036	0.0014	0.0005
	1	0.8981	0.6974	0.4922	0.3221	0.1971	0.1130	0.0606	0.0302	0.0139	0.0059
12	0	0.5404	0.2824	0.1422	0.0687	0.0317	0.0138	0.0057	0.0022	0.0008	0.0002
	1	0.8816	0.6590	0.4435	0.2749	0.1584	0.0850	0.0424	0.0196	0.0083	0.0032

(9.5) Table for Poisson random variable $P(Po(\mu) \leq k)$.

k	μ									
	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0
0	0.9048	0.8187	0.7408	0.6703	0.6065	0.5488	0.4966	0.4493	0.4066	0.3679
1	0.9953	0.9825	0.9631	0.9384	0.9098	0.8781	0.8442	0.8088	0.7725	0.7358
2	0.9998	0.9989	0.9964	0.9921	0.9856	0.9769	0.9659	0.9526	0.9371	0.9197
3	1.0000	0.9999	0.9997	0.9992	0.9982	0.9966	0.9942	0.9909	0.9865	0.9810
4	1.0000	1.0000	1.0000	0.9999	0.9998	0.9996	0.9992	0.9986	0.9977	0.9963
5	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9997	0.9994
6	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999
7	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
k	μ									
	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0
0	0.3329	0.3012	0.2725	0.2466	0.2231	0.2019	0.1827	0.1653	0.1496	0.1353
1	0.6990	0.6626	0.6268	0.5918	0.5578	0.5249	0.4932	0.4628	0.4337	0.4060
2	0.9004	0.8795	0.8571	0.8335	0.8088	0.7834	0.7572	0.7306	0.7037	0.6767
3	0.9743	0.9662	0.9569	0.9463	0.9344	0.9212	0.9068	0.8913	0.8747	0.8571
4	0.9946	0.9923	0.9893	0.9857	0.9814	0.9763	0.9704	0.9636	0.9559	0.9473
5	0.9990	0.9985	0.9978	0.9968	0.9955	0.9940	0.9920	0.9896	0.9868	0.9834
6	0.9999	0.9997	0.9996	0.9994	0.9991	0.9987	0.9981	0.9974	0.9966	0.9955
7	1.0000	1.0000	0.9999	0.9999	0.9998	0.9997	0.9996	0.9994	0.9992	0.9989
8	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9999	0.9998
9	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
k	μ									
	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0
0	0.1225	0.1108	0.1003	0.0907	0.0821	0.0743	0.0672	0.0608	0.0550	0.0498
1	0.3796	0.3546	0.3309	0.3084	0.2873	0.2674	0.2487	0.2311	0.2146	0.1991
2	0.6496	0.6227	0.5960	0.5697	0.5438	0.5184	0.4936	0.4695	0.4460	0.4232
3	0.8386	0.8194	0.7993	0.7787	0.7576	0.7360	0.7141	0.6919	0.6696	0.6472
4	0.9379	0.9275	0.9162	0.9041	0.8912	0.8774	0.8629	0.8477	0.8318	0.8153
5	0.9796	0.9751	0.9700	0.9643	0.9580	0.9510	0.9433	0.9349	0.9258	0.9161
6	0.9941	0.9925	0.9906	0.9884	0.9858	0.9828	0.9794	0.9756	0.9713	0.9665
7	0.9985	0.9980	0.9974	0.9967	0.9958	0.9947	0.9934	0.9919	0.9901	0.9881
8	0.9997	0.9995	0.9994	0.9991	0.9989	0.9985	0.9981	0.9976	0.9969	0.9962
9	0.9999	0.9999	0.9999	0.9998	0.9997	0.9996	0.9995	0.9993	0.9991	0.9989
10	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9999	0.9998	0.9997	0.9996
11	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9999
12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

Table for Poisson random variable $P(Po(\mu) \leq k)$.

k	μ									
	3.2	3.4	3.6	3.8	4.0	4.2	4.4	4.6	4.8	5.0
0	0.0408	0.0334	0.0273	0.0224	0.0183	0.0150	0.0123	0.0101	0.0082	0.0067
1	0.1712	0.1468	0.1257	0.1074	0.0916	0.0780	0.0663	0.0563	0.0477	0.0404
2	0.3799	0.3397	0.3027	0.2689	0.2381	0.2127	0.1851	0.1626	0.1425	0.1247
3	0.6025	0.5584	0.5152	0.4735	0.4335	0.3954	0.3594	0.3257	0.2942	0.2650
4	0.7806	0.7442	0.7064	0.6678	0.6288	0.5898	0.5512	0.5132	0.4763	0.4405
5	0.8946	0.8705	0.8441	0.8156	0.7851	0.7531	0.7199	0.6858	0.6510	0.6160
6	0.9534	0.9421	0.9267	0.9091	0.8893	0.8675	0.8436	0.8180	0.7908	0.7622
7	0.9832	0.9769	0.9682	0.9599	0.9509	0.9419	0.9319	0.9214	0.9049	0.8866
8	0.9943	0.9917	0.9883	0.9840	0.9786	0.9721	0.9642	0.9549	0.9442	0.9319
9	0.9982	0.9973	0.9960	0.9942	0.9919	0.9889	0.9851	0.9805	0.9749	0.9682
10	0.9995	0.9992	0.9987	0.9981	0.9972	0.9959	0.9943	0.9922	0.9896	0.9863
11	0.9999	0.9998	0.9996	0.9994	0.9991	0.9986	0.9980	0.9971	0.9960	0.9945
12	1.0000	0.9999	0.9999	0.9999	0.9998	0.9997	0.9996	0.9993	0.9990	0.9986
13	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9996	0.9993	0.9990	0.9986
14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9997	0.9995
15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9998
k	μ									
	5.2	5.4	5.6	5.8	6.0	6.5	7.0	7.5	8.0	8.5
0	0.0055	0.0045	0.0037	0.0030	0.0025	0.0015	0.0009	0.0006	0.0003	0.0002
1	0.0342	0.0289	0.0244	0.0206	0.0174	0.0113	0.0073	0.0047	0.0030	0.0019
2	0.1088	0.0948	0.0824	0.0715	0.0620	0.0430	0.0286	0.0203	0.0138	0.0093
3	0.2381	0.2133	0.1906	0.1700	0.1512	0.1118	0.0818	0.0591	0.0424	0.0301
4	0.4061	0.3733	0.3422	0.3127	0.2851	0.2237	0.1730	0.1321	0.0996	0.0744
5	0.5809	0.5461	0.5119	0.4783	0.4457	0.3690	0.3007	0.2414	0.1912	0.1496
6	0.7324	0.7017	0.6703	0.6384	0.6063	0.5265	0.4497	0.3782	0.3134	0.2562
7	0.8449	0.8217	0.7970	0.7710	0.7440	0.6728	0.5987	0.5246	0.4530	0.3856
8	0.9181	0.9027	0.8857	0.8672	0.8472	0.7916	0.7291	0.6620	0.5925	0.5231
9	0.9603	0.9512	0.9409	0.9292	0.9161	0.8774	0.8305	0.7764	0.7166	0.6530
10	0.9823	0.9775	0.9718	0.9651	0.9574	0.9332	0.9015	0.8622	0.8159	0.7634
11	0.9927	0.9904	0.9875	0.9841	0.9799	0.9661	0.9467	0.9208	0.8881	0.8487
12	0.9972	0.9962	0.9949	0.9932	0.9912	0.9840	0.9730	0.9573	0.9362	0.9091
13	0.9990	0.9986	0.9980	0.9973	0.9964	0.9929	0.9872	0.9784	0.9658	0.9486
14	0.9999	0.9995	0.9993	0.9990	0.9986	0.9970	0.9943	0.9897	0.9827	0.9726
15	0.9999	0.9998	0.9998	0.9996	0.9995	0.9988	0.9976	0.9954	0.9918	0.9862
16	1.0000	0.9999	0.9999	0.9999	0.9998	0.9996	0.9990	0.9980	0.9963	0.9934
17	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9996	0.9992	0.9984	0.9970
18	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9997	0.9993	0.9987
19	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9995
20	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9998
21	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999
22	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

Table for Poisson random variable $P(Po(\mu) \leq k)$.

k	μ														
	9.0	9.5	10.0	11.0	12.0	13.0	14.0	15.0	16.0	17.0					
0	0.0001	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000					
1	0.0012	0.0008	0.0005	0.0002	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000					
2	0.0062	0.0042	0.0028	0.0012	0.0005	0.0002	0.0001	0.0000	0.0000	0.0000					
3	0.0212	0.0149	0.0103	0.0049	0.0023	0.0011	0.0005	0.0002	0.0001	0.0000					
4	0.0550	0.0403	0.0293	0.0151	0.0076	0.0037	0.0018	0.0009	0.0004	0.0002					
5	0.1157	0.0885	0.0671	0.0375	0.0203	0.0107	0.0055	0.0028	0.0014	0.0007					
6	0.2068	0.1649	0.1301	0.0786	0.0458	0.0259	0.0142	0.0076	0.0040	0.0021					
7	0.3239	0.2687	0.2202	0.1432	0.0895	0.0540	0.0316	0.0180	0.0100	0.0054					
8	0.4557	0.3918	0.3328	0.2320	0.1550	0.0998	0.0621	0.0374	0.0220	0.0126					
9	0.5874	0.5218	0.4579	0.3405	0.2424	0.1658	0.1094	0.0699	0.0433	0.0261					
10	0.7060	0.6453	0.5830	0.4599	0.3472	0.2517	0.1757	0.1185	0.0774	0.0491					
11	0.8030	0.7520	0.6968	0.5793	0.4616	0.3532	0.2600	0.1848	0.1270	0.0847					
12	0.8758	0.8364	0.7916	0.6887	0.5760	0.4631	0.3585	0.2676	0.1931	0.1350					
13	0.9261	0.8981	0.8645	0.7813	0.6815	0.5730	0.4644	0.3632	0.2745	0.2009					
14	0.9585	0.9400	0.9165	0.8540	0.7720	0.6751	0.5704	0.4657	0.3675	0.2808					
15	0.9780	0.9665	0.9513	0.9074	0.8444	0.7636	0.6694	0.5681	0.4667	0.3715					
16	0.9889	0.9823	0.9730	0.9441	0.8987	0.8355	0.7559	0.6641	0.5660	0.4677					
17	0.9947	0.9911	0.9857	0.9678	0.9370	0.8905	0.8272	0.7489	0.6593	0.5640					
18	0.9976	0.9957	0.9928	0.9823	0.9626	0.9302	0.8826	0.8195	0.7423	0.6550					
19	0.9989	0.9980	0.9965	0.9907	0.9787	0.9573	0.9235	0.8752	0.8122	0.7363					
20	0.9996	0.9991	0.9984	0.9953	0.9884	0.9750	0.9521	0.9170	0.8682	0.8055					
21	0.9998	0.9996	0.9993	0.9977	0.9939	0.9859	0.9712	0.9469	0.9108	0.8615					
22	0.9999	0.9999	0.9997	0.9990	0.9970	0.9924	0.9833	0.9673	0.9418	0.9047					
23	1.0000	0.9999	0.9999	0.9995	0.9985	0.9960	0.9907	0.9805	0.9633	0.9367					
24	1.0000	1.0000	1.0000	0.9998	0.9993	0.9980	0.9950	0.9888	0.9777	0.9594					
25	1.0000	1.0000	1.0000	0.9999	0.9997	0.9990	0.9974	0.9938	0.9869	0.9748					
26	1.0000	1.0000	1.0000	1.0000	0.9999	0.9995	0.9987	0.9967	0.9925	0.9848					
27	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9994	0.9983	0.9959	0.9912					
28	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9991	0.9978	0.9950					
29	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9996	0.9994	0.9986					
30	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9993					
31	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9996					
32	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9996					
33	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9999	0.9999					
34	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999					
35	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000					