

Exam in Transform Theory

TATA57/TEN1 2021-10-22

You are permitted to use:

- Transformteori - Sammanfattning, Formler & Lexikon (from MAI);
- Table of Formulæ (by Johan Thim). This one is included at the end of the exam.
- A calculator (with cleared memories).

Swedish version of the exam comes after the English version. You may answer in Swedish or English.

Each problem is worth 5 points (for a total of 35 points). No half-points will be awarded in the grading. A solution is deemed good if the question is awarded at least 3 points out of 5. For grade n , you need at least n good answers.

ERASMUS students will have their grades marked according to the scale: A = grade 5, B = grade 4, C = grade 3.

Grading (sufficient limit): 15 points for grade 3. Your solutions need to be complete, well motivated, carefully written and concluded with a clear answer. Be very careful with motivations since these are a huge part of your solutions. Make sure to point out that conditions in theorems you are using hold. Assumptions you make need to be explicit. The exercises are not necessarily in order of difficulty.

Solutions can be found on the homepage a couple of hours after the finished exam.

1. For each of these statements, prove or disprove the claims. For each correct answer, you get 1 p and for each incorrect answer, you get 0 p.

- (a) The Fourier transform of a function $f \in G$ is always bounded.
- (b) The function $U(z) = \cos(|z|)$, $|z| > 1$, is the Z transform of a bounded sequence $u[n]$, $n = 0, 1, 2, \dots$
- (c) The function $u(t) = 3e^{t^2}$ is exponentially bounded, that is, there exists $C > 0$ and $k > 0$ such that $|u(t)| \leq Ce^{kt}$ for $t \geq 0$.
- (d) The 2π -periodic function defined by $f(x) = \arctan(|x|)$ when $|x| \leq \pi$ has a Fourier series that converges for all $x \in \mathbf{R}$.
- (e) The 2π -periodic function defined by $y(x) = |x|^3$ when $|x| \leq \pi$ solves some differential equation $y''(x) + ay(x + v) = 0$, $x \in \mathbf{R}$, where $a \neq 0$ och $v \in \mathbf{R}$.

2. Calculate $\sum_{k=0}^m (k^2 + 3k)2^k$ for $m = 0, 1, 2, \dots$

3. Let

$$u(x) = \begin{cases} 1, & |x| < \frac{\pi}{2}, \\ 0, & \frac{\pi}{2} \leq |x| \leq \pi, \end{cases}$$

and extend u periodically. Find the Fourier series for $u(x)$ and show what the Fourier series converges to for $x \in \mathbf{R}$. Also find the value of $\sum_{m=0}^{\infty} \frac{1}{(2m+1)^2}$.

4. Find all continuous exponentially bounded solutions to

$$y'(t) - 2y(t) = \begin{cases} 4t, & 0 \leq t < 1, \\ 4(3-t), & t \geq 1, \end{cases}$$

where $0 \leq t < \infty$ and $y(0) = 0$. Is your solution of class C^1 ? In what sense does your solution solve the equation when $t = 1$?

5. Calculate the integral $\int_0^{\infty} \frac{\cos(tx)}{4+x^2} dx$ for $t \in \mathbf{R}$.

6. Find the continuous function f that has the Fourier coefficients $c_n = e^{-n}$ if $n \geq 0$ and $c_n = 0$ if $n < 0$. Note that you also have to prove that your suggested function f actually has these Fourier coefficients.

7. Construct a continuous function $u \in G(\mathbf{R})$ such that $\mathcal{F}u \notin L^1(\mathbf{R})$.

Svensk översättning av uppgifterna

1. För varje påstående, bevisa eller motbevisa påståendet. Du får 1 p för varje rätt och 0 p för varje fel.

- (a) Fouriertransformen av en funktion $f \in G$ är alltid begränsad.
- (b) Funktionen $U(z) = \cos(|z|)$, $|z| > 1$, är Z transformen av en begränsad följd $u[n]$, $n = 0, 1, 2, \dots$
- (c) Funktionen $u(t) = 3e^{t^2}$ är exponentiellt begränsad, det vill säga det existerar $C > 0$ och $k > 0$ så att $|u(t)| \leq Ce^{kt}$ för $t \geq 0$.
- (d) Den 2π -periodiska funktionen definierad av $f(x) = \arctan(|x|)$ då $|x| \leq \pi$ har en Fourierserie som konvergerar för alla $x \in \mathbf{R}$.
- (e) Den 2π -periodiska funktionen definierad av $y(x) = |x|^3$ då $|x| \leq \pi$ löser någon differentialekvation $y''(x) + ay(x+v) = 0$, $x \in \mathbf{R}$, där $a \neq 0$ och $v \in \mathbf{R}$.

2. Beräkna $\sum_{k=0}^m (k^2 + 3k)2^k$ för $m = 0, 1, 2, \dots$

3. Låt

$$u(x) = \begin{cases} 1, & |x| < \frac{\pi}{2}, \\ 0, & \frac{\pi}{2} \leq |x| \leq \pi, \end{cases}$$

och utvidga u periodiskt. Hitta Fourierserien för $u(x)$ och visa vad Fourierserien konvergerar till för $x \in \mathbf{R}$. Beräkna även serien $\sum_{m=0}^{\infty} \frac{1}{(2m+1)^2}$.

4. Hitta alla kontinuerliga exponentiellt begränsade lösningar till

$$y'(t) - 2y(t) = \begin{cases} 4t, & 0 \leq t < 1, \\ 4(3-t), & t \geq 1, \end{cases}$$

där $0 \leq t < \infty$ och $y(0) = 0$. Är din lösning av klassen C^1 ? I vilken mening löser din lösning ekvationen då $t = 1$?

5. Beräkna integralen $\int_0^{\infty} \frac{\cos(tx)}{4+x^2} dx$ för $t \in \mathbf{R}$.

6. Hitta den kontinuerliga funktion f som har Fourierkoefficienterna $c_n = e^{-n}$ om $n \geq 0$ och $c_n = 0$ om $n < 0$. Notera att du också måste visa att din föreslagna funktion f faktiskt har dessa Fourierkoefficienter.

7. Konstruera en kontinuerlig funktion $u \in G(\mathbf{R})$ så att $\mathcal{F}u \notin L^1(\mathbf{R})$.

Table of formulæ

Johan Thim (johan.thim@liu.se)

May 25, 2021

1 Notation and Definitions

- \mathbf{R} is the set of all real numbers.
- \mathbf{Q} is the set of all rational numbers.
- \mathbf{C} is the set of all complex numbers.
- \mathbf{Z} is the set of all integers.
- $\mathbf{N} = \{0, 1, 2, \dots\}$ is the set of all natural numbers (including 0).

For $z = x + iy \in \mathbf{C}$, $x, y \in \mathbf{R}$,

$$\operatorname{Re} z = x, \quad \operatorname{Im} z = y, \quad |z| = \sqrt{x^2 + y^2}.$$

1.1 Continuity and Differentiability

- One-sided limits:
- One-sided derivatives:
- $C(I)$: The set of all continuous functions on a set I .
- $C^m(I)$: The set of all continuously differentiable (up to order m) functions on a set I .

A function $u: I \rightarrow \mathbf{C}$ on an interval I is called piecewise continuous if...

- I is finite and there are a finite number of points such that u is continuous everywhere on I except for at these points. Moreover, if $c \in I$ is a point where u is discontinuous, the limits
- $\lim_{I \ni x \rightarrow c^-} u(x)$ and $\lim_{I \ni x \rightarrow c^+} u(x)$ if points on the boundary of I .
- I is infinite and there are a finite number of exception points (as in the finite case) in each finite sub-interval of I .

1.2 Function Spaces

A normed linear space is a linear space V endowed with a norm $\|\cdot\|: V \rightarrow [0, \infty[$ such that

- (i) $\|u\| \geq 0$
 - (ii) $\|\alpha u\| = |\alpha| \|u\|$, $\alpha \in \mathbf{C}$
 - (iii) $\|u + v\| \leq \|u\| + \|v\|$.
- An inner product $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbf{C}$ on a vector space V satisfies
- (i) $\langle u, v \rangle = \overline{\langle v, u \rangle}$
 - (ii) $\langle u + v, w \rangle = \langle u, w \rangle + \langle v, w \rangle$
 - (iii) $\langle \alpha u, v \rangle = \alpha \langle u, v \rangle$
 - (iv) $\langle u, u \rangle \geq 0$
 - (v) $\langle u, u \rangle = 0 \Leftrightarrow u = 0$.

In an inner product space, we use $\|u\| = \sqrt{\langle u, u \rangle}$ as the norm.

Sequence Spaces

The sequence spaces l^p , $1 \leq p \leq \infty$, consists of sequences $(x_0, x_1, x_2, x_3, \dots)$, $x_i \in \mathbf{C}$, such that the norm

$$\|x\|_p = \left(\sum_{k=0}^{\infty} |x_k|^p \right)^{1/p} < \infty, \quad 1 \leq p < \infty,$$

or

$$\|x\|_{l^\infty} = \sup_{k \geq 0} |x_k| < \infty, \quad p = \infty.$$

Sometimes we write $l^p(\mathbf{N})$. The spaces $l^p(\mathbf{Z})$ are defined analogously. Only l^2 is an inner product space with

$$\langle x, y \rangle = \sum_{k=0}^{\infty} x_k \overline{y_k}, \quad x, y \in l^2.$$

Lebesgue Spaces (integrable functions)

The space $L^1(a, b)$ of absolutely integrable functions $u:]a, b[\rightarrow \mathbf{C}$ with norm

$$\|f\|_{L^1(a,b)} = \int_a^b |f(x)| dx < \infty.$$

The space $L^2(a, b)$ consists of all “square integrable” functions with the norm

$$\|f\|_{L^2(a,b)} = \left(\int_a^b |f(x)|^2 dx \right)^{1/2} < \infty,$$

which is an inner product space with

$$\langle f, g \rangle = \int_a^b f(x) \overline{g(x)} dx.$$

The space $L^\infty(a, b)$ of bounded functions with norm

$$\|f\|_{L^\infty(a,b)} = \sup_{a \leq x \leq b} |f(x)| < \infty.$$

Note that $a = -\infty$ and/or $b = \infty$ is allowed (so we might write $L^p(\mathbf{R})$). Sometimes we write $\|f\|_p$ instead of $\|f\|_{L^p(a,b)}$.

Spaces of Piecewise Functions

- $E[a, b]$ (or E): The linear space of all piecewise continuous functions on an interval $[a, b]$.
- $E^+[a, b]$ (or E^+): The linear space of those $u \in E[a, b]$ such that $D^-u(x)$ exists for $a < x \leq b$ and that $D^+u(x)$ exists for $a \leq x < b$.
- $G(\mathbf{R})$ (or G): The linear space of all piecewise continuous functions on \mathbf{R} that are absolutely integrable on \mathbf{R} .

1.3 Special Functions

- **Heaviside function:**
- **Signum function:**

$$H(x) = \begin{cases} 0, & x < 0, \\ 1, & x \geq 0. \end{cases}$$

$$\operatorname{sgn}(x) = \begin{cases} -1, & x < 0, \\ 0, & x = 0, \\ 1, & x > 0. \end{cases}$$

Discrete Functions

- **Discrete Heaviside function:**
- **Discrete impulse function:**
- **Binomial coefficient functions:**

$$\binom{n}{k} = \begin{cases} \frac{n!}{(n-k)!k!}, & k = 0, 1, 2, \dots, \\ 0, & k > n. \end{cases}$$

Convolutions (on \mathbf{R})

The convolution $u * v : \mathbf{R} \rightarrow \mathbf{C}$ of two functions $u : \mathbf{R} \rightarrow \mathbf{C}$ and $v : \mathbf{R} \rightarrow \mathbf{C}$ is defined by

$$(u * v)(x) = \int_{-\infty}^{\infty} u(t)v(x-t)dt, \quad x \in \mathbf{R},$$

whenever this integral exists. If $u, v \in L^1(\mathbf{R})$, then $u * v \in L^1(\mathbf{R})$. If one function is also bounded, then $u * v$ is continuous and bounded. Suppose that $u, v, w \in G(\mathbf{R})$ and one function in each convolution is bounded. Then the convolution has the following properties.

- Associative: $((u * v) * w)(x) = (u * (v * w))(x)$.
- Distributive: $((u + v) * w)(x) = (u * w)(x) + (v * w)(x)$.
- Commutative: $(u * v)(x) = (v * u)(x)$.

Convolutions (on \mathbf{Z})

- For $u, v : \mathbf{Z} \rightarrow \mathbf{C}$, the **discrete convolution** $u * v$ is

$$(u * v)[n] = \sum_{k=-\infty}^{\infty} u[k]v[n-k], \quad n \in \mathbf{Z},$$

whenever this series exists.

- For $u, v : \mathbf{N} \rightarrow \mathbf{C}$, the **unilateral (or one-sided) discrete convolution** $u * v$ is

$$(u * v)[n] = \sum_{k=0}^n u[k]v[n-k], \quad n = 0, 1, 2, \dots$$

1.4 Inequalities

- **The Cauchy-Schwarz inequality:** If $u, v \in V$ and V is an inner product space, then
- **Bessel's inequality:** Let V be an inner product space, let $v \in V$ and let $\{e_1, e_2, \dots\}$ be an ON system in V . Then

$$|\langle u, v \rangle| \leq \|u\|\|v\|.$$

$$\sum_{k=1}^{\infty} |\langle v, e_k \rangle|^2 \leq \|v\|^2.$$

This implies the **Riemann-Lebesgue lemma** for inner product spaces:

$$\lim_{n \rightarrow \infty} \langle v, e_n \rangle = 0.$$

- **The triangle inequality:** In a normed space V ,
- **Young's inequality** ($r = p = q = 1$):

$$\|u\| - \|v\| \leq \|u + v\| \leq \|u\| + \|v\|.$$

$$\|u * v\|_{L^1(\mathbf{R})} \leq \|u\|_{L^1(\mathbf{R})}\|v\|_{L^1(\mathbf{R})}.$$

and

$$\|u * v\|_{L^1(\mathbf{Z})} \leq \|u\|_{L^1(\mathbf{Z})}\|v\|_{L^1(\mathbf{Z})}.$$

1.5 Convergence of Sequences

Let u_1, u_2, \dots be a sequence in a normed space V . We say that $u_n \rightarrow u$ for some $u \in V$ if $\|u_n - u\| \rightarrow 0$ as $n \rightarrow \infty$. This is called **strong convergence** or **convergence in norm**.

Convergence of Functions

- **Pointwise convergence:** We say that $u_k \rightarrow u$ pointwise on I as $k \rightarrow \infty$ if

$$\lim_{k \rightarrow \infty} u_k(x) = u(x)$$

for every $x \in I$. We often refer to u as the *limiting function*.

- **Uniform convergence:** We say that $u_k \rightarrow u$ uniformly on $[a, b]$ as $k \rightarrow \infty$ if

$$\lim_{k \rightarrow \infty} \|u_k - u\|_{L^\infty(a,b)} = 0.$$

Weierstrass' M-test: If $I \subset \mathbf{R}$ and M_k , $k = 1, 2, \dots$, are constants such that $|u_k(x)| \leq M_k$ for $x \in I$, then

$$\sum_{k=1}^{\infty} M_k < \infty \Rightarrow \sum_{k=1}^{\infty} u_k(x) \text{ converges uniformly on } I.$$

If:

- u_0, u_1, u_2, \dots are continuous functions on $[a, b]$
- and $u(x) = \sum_{k=0}^{\infty} u_k(x)$ is uniformly convergent for $x \in [a, b]$,

then

- the series u is a continuous function on $[a, b]$,
- we can exchange the order of summation and integration:

$$\int_c^d u(x) dx = \int_c^d \left(\sum_{k=0}^{\infty} u_k(x) \right) dx = \sum_{k=0}^{\infty} \int_c^d u_k(x) dx, \quad \text{for } a \leq c < d \leq b,$$

- and if in addition $\sum_{k=0}^{\infty} u'_k(x)$ converges uniformly on $[a, b]$, then

$$u'(x) = \frac{d}{dx} \left(\sum_{k=0}^{\infty} u_k(x) dx \right) = \sum_{k=0}^{\infty} \frac{d}{dx} u_k(x) = \sum_{k=0}^{\infty} u'_k(x), \quad x \in [a, b].$$

1.6 Integration Theory

The **principal value** integral is defined by

$$\text{P. V. } \int_{-\infty}^{\infty} u(x) dx = \lim_{R \rightarrow \infty} \int_{-R}^R u(x) dx.$$

- If $F(x) = \int_{-\infty}^x f(x, y) dy$ exists for every $x \in I$ and

$$\sup_{x \in I} \left| \int_{-R}^R f(x, y) dy - F(x) \right| \rightarrow 0, \quad \text{as } R \rightarrow \infty,$$

then we call the integral defining $F(x)$ uniformly convergent on I .

- **Dominated convergence:**

If:

- $f: \mathbf{R}^2 \rightarrow \mathbf{C}$,
- $F(x) = \int_{-\infty}^{\infty} f(x, y) dy$ exists for all x ,
- there is a $g \in L^1(\mathbf{R})$ such that $|f(x, y)| \leq g(y)$ for all $x, y \in \mathbf{R}$,

then $\int_{-\infty}^{\infty} f(x, y) dy$ converges uniformly on \mathbf{R} .

- **Continuity:** If $f: \mathbf{R}^2 \rightarrow \mathbf{C}$ is continuous on $[c, d] \times [a, R]$. Then

$$- F_R(x) = \int_a^R f(x, y) dy \text{ is continuous on } [c, d]$$

- and if in addition f is continuous on $[c, d] \times [a, \infty[$ and $F(x) = \int_a^{\infty} f(x, y) dy$ converges uniformly (on $[c, d]$), then F is continuous.

- **Order of integration:** If $f: \mathbf{R}^2 \rightarrow \mathbf{C}$ is continuous on $[c, d] \times [a, \infty[$ and $F(x)$ converges uniformly (on $[c, d]$), then

$$\int_c^d \left(\int_a^{\infty} f(x, y) dy \right) dx = \int_a^{\infty} \left(\int_c^d f(x, y) dx \right) dy.$$

- Note that we can let $a = -\infty$ in the previous theorems by exchanging $[a, R]$ by $[-R, R]$ and consider the principal values.

- **Leibniz's rule:** If

- $f: \mathbf{R}^2 \rightarrow \mathbf{C}$ and $f'_x(x, y)$ exist and are continuous,
- $F(x) = \int_{-\infty}^{\infty} f(x, y) dy$ is convergent for every x ,
- and $\int_{-\infty}^{\infty} f'_x(x, y) dy$ is uniformly convergent,

then

$$F'(x) = \frac{d}{dx} \int_{-\infty}^{\infty} f(x, y) dy = \int_{-\infty}^{\infty} f'_x(x, y) dy.$$

2 Fourier Series

For $u \in L^1(-\pi, \pi)$:

$$a_k = \frac{1}{\pi} \int_{-\pi}^{\pi} u(x) \cos kx dx \quad \text{and} \quad b_k = \frac{1}{\pi} \int_{-\pi}^{\pi} u(x) \sin kx dx \quad \text{or} \quad c_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} u(x) e^{-ikx} dx$$

are the Fourier coefficients (real or complex) for u . The series

$$S(x) = \frac{a_0}{2} + \sum_{k=1}^{\infty} (a_k \cos kx + b_k \sin kx) = \sum_{k=-\infty}^{\infty} c_k e^{ikx}$$

is called the **Fourier series** of the function u (real or complex). We write $u(x) \sim S(x)$. Note that:

- if u is even, then $b_k = 0$ for $k = 1, 2, 3, \dots$;
- if u is odd, then $a_k = 0$ for $k = 1, 2, 3, \dots$.

If u is a T -periodic function, we define $\Omega = \frac{2\pi}{T}$. The real Fourier series of u is then given by

$$u(x) \sim \frac{a_0}{2} + \sum_{k=1}^{\infty} (a_k \cos k\Omega x + b_k \sin k\Omega x),$$

where

$$a_k = \frac{2}{T} \int_{-T/2}^{T/2} u(x) \cos k\Omega x \, dx \quad \text{and} \quad b_k = \frac{2}{T} \int_{-T/2}^{T/2} u(x) \sin k\Omega x \, dx.$$

The complex Fourier series is given by

$$u(x) \sim \sum_{k=-\infty}^{\infty} c_k e^{ik\Omega x}, \quad \text{where } c_k = \frac{1}{T} \int_{-T/2}^{T/2} u(x) e^{-ik\Omega x} \, dx.$$

Sometimes we denote $c_k = \hat{u}[k]$.

2.1 Parseval's identity

- **Parseval's identity:**

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} |u(x)|^2 \, dx = \sum_{k=-\infty}^{\infty} |c_k|^2 \quad \text{or} \quad \frac{1}{\pi} \int_{-\pi}^{\pi} |u(x)|^2 \, dx = \frac{|a_0|^2}{2} + \sum_{k=1}^{\infty} (|a_k|^2 + |b_k|^2),$$

where $u(x) \sim \sum_{k=-\infty}^{\infty} c_k e^{ikx}$ (or the real counterpart).

- **Parseval's generalized identity:**

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} u(x) \overline{v(x)} \, dx = \sum_{k=-\infty}^{\infty} c_k \overline{d_k},$$

where $u(x) \sim \sum_{k=-\infty}^{\infty} c_k e^{ikx}$ and $v(x) \sim \sum_{k=-\infty}^{\infty} d_k e^{ikx}$.

2.2 Convergence

Kernels

- The **Dirichlet kernel:** $D_n(x) = \sum_{k=-n}^n e^{ikx}$, $x \in \mathbf{R}$, $n = 1, 2, 3, \dots$
- The **Fejér kernel:** $F_n(x) = \frac{1}{n+1} \sum_{l=0}^n \sum_{k=-l}^l e^{ikx} = \sum_{k=-n}^n \left(1 - \frac{|k|}{n+1}\right) e^{ikx}$, $n = 0, 1, 2, \dots$

2.3 Convergence Results

- If $u \in L^1(-\pi, \pi)$, then u has a Fourier series.
- Let $u \in E'$. Then

$$S_n(x) = \sum_{k=-n}^n c_k e^{ikx} \rightarrow \frac{u(x^+) + u(x^-)}{2}, \quad x \in [-\pi, \pi].$$

- If $u \in E$ and $D^\pm u(x)$ exists, then

$$\lim_{n \rightarrow \infty} S_n(x) = \frac{u(x^+) + u(x^-)}{2}.$$

- If $\sum_{k=-\infty}^{\infty} |c_k| < \infty$, then $\sum_{k=-\infty}^{\infty} c_k e^{ikx}$ converges uniformly.
- If $u \in E$, then the Fejér means $\overline{S}_n(x) \rightarrow \frac{u(x^+) + u(x^-)}{2}$.
- If $u, v \in E$ and $\hat{u}[k] = \hat{v}[k]$, $k \in \mathbf{Z}$, then $u(x) = v(x)$ whenever u and v are continuous at x .
- If $u \in E$, u is continuous and $u(-\pi) = u(\pi)$, then $S_n(x)$ converges uniformly to $u(x)$.
- If $u' \in E$ and u is continuous on $[a, b] \subset]-\pi, \pi[$, then $S_n(x)$ converges uniformly on $[a, b]$.
- If $u \in E$ is continuous and $u(-\pi) = u(\pi)$, then $\overline{S}_n(x)$ converges uniformly to $u(x)$.

The statement $u' \in E$ does not mean that $u'(x)$ exists everywhere, but that there exists a $v \in E$ such that $v(x) = u'(x)$ when $u'(x)$ exists and that u' exists everywhere except at a finite number of points in $[a, b]$.

2.4 General Fourier Series

- For a given ON system, the complex numbers $\langle v, e_k \rangle$, $k = 1, 2, \dots$, are called the **generalized Fourier coefficients** of v .
- If $W = \{e_1, e_2, \dots\}$ is an ON system in V , then W is closed if and only if **Parseval's identity** holds:

$$\sum_{k=1}^{\infty} |\langle v, e_k \rangle|^2 = \|v\|^2, \quad v \in V;$$

or if $a_k = \langle u, e_k \rangle$ and $b_k = \langle v, e_k \rangle$, then

$$\langle u, v \rangle = \sum_{k=1}^{\infty} a_k \overline{b_k}.$$

2.5 Rules for Fourier Coefficients

Let $u, v \in E$ be periodic with period $T > 0$ and define $\Omega = 2\pi/T$.

Table 1: Rules for Fourier Coefficients

Function	Fourier coefficient	Notes
$c_1 u(x) + c_2 v(x)$	$c_1 \widehat{u}[n] + c_2 \widehat{v}[n]$	
$(u * v)_T(x)$	$\widehat{u}[n] \widehat{v}[n]$	periodic convolution [†]
$u(x)v(x)$	$(\widehat{u} * \widehat{v})[n]$	
$e^{im\Omega x} u(x)$	$\widehat{u}[n - m]$	$m \in \mathbf{Z}$
$u(x - a)$	$e^{-in\Omega a} \widehat{u}[n]$	$a \in \mathbf{R}$
$u(ax)$	$\widehat{u}[n]$	period T/a , $a > 0$
$u(-x)$	$\widehat{u}[-n]$	
$\overline{u(x)}$	$\overline{\widehat{u}[-n]}$	
$u'(x)$	$in\Omega \widehat{u}[n]$	
$u^{(k)}(x)$	$(i\Omega n)^k \widehat{u}[n]$	$k = 1, 2, \dots$

[†] $(u * v)_T(x) = \frac{1}{T} \int_0^T u(x-t)v(t) dt$.

3 The Fourier Transform

The Fourier transform of a function $u: \mathbf{R} \rightarrow \mathbf{C}$ given by

$$U(\omega) = \widehat{u}(\omega) = \mathcal{F}u(\omega) = \int_{-\infty}^{\infty} u(x)e^{-i\omega x} dx, \quad \omega \in \mathbf{R},$$

when this integral exists.

- If $u \in L^1(\mathbf{R})$ then $\mathcal{F}u(\omega)$ exists for all $\omega \in \mathbf{R}$ and $\|\mathcal{F}u\|_{\infty} \leq \|u\|_{L^1(\mathbf{R})}$.
- For $u \in G$, the Fourier transform $\mathcal{F}u$ is uniformly continuous on \mathbf{R} .
- The Riemann-Lebesgue lemma: For $u \in G$ we have $\mathcal{F}u(\omega) \rightarrow 0$ as $|\omega| \rightarrow \infty$.

3.1 Convergence

Kernels

- The Dirichlet kernel (on \mathbf{R}):

$$D_R(x) = \frac{\sin(Rx)}{\pi x}, \quad x \neq 0,$$

and $D_R(0) = R/\pi$.

- The Fejér kernel (on \mathbf{R}):

$$F_M(t) = \frac{1}{2\pi} \int_{-M}^M \left(1 - \frac{|\omega|}{M}\right) e^{i\omega t} d\omega = \frac{1 - \cos Mx}{\pi M x^2} = \frac{M}{2\pi} \left(\frac{\sin(Mx/2)}{Mx/2}\right)^2,$$

where the last two equalities assumes that $x \neq 0$.

Inversion

- If $u \in G(\mathbf{R})$ and $D^{\pm}u(x)$ exists, then

$$\lim_{R \rightarrow \infty} \frac{1}{2\pi} \int_{-R}^R \mathcal{F}u(\omega) e^{i\omega x} d\omega = \frac{u(x^+) + u(x^-)}{2}.$$

- If $u \in G(\mathbf{R})$, then

$$\lim_{R \rightarrow \infty} \frac{1}{2\pi} \int_{-R}^R \mathcal{F}u(\omega) \left(1 - \frac{|\omega|}{R}\right) e^{i\omega x} d\omega = \frac{u(x^+) + u(x^-)}{2}.$$

- If $u \in G(\mathbf{R})$, then

$$\lim_{R \rightarrow \infty} \frac{1}{2\pi} \int_{-R}^R \mathcal{F}u(\omega) e^{i\omega x} d\omega = \frac{u(x^+) + u(x^-)}{2},$$

whenever the limit exists.

- **Uniqueness:** If $u, v \in G(\mathbf{R})$ and $\mathcal{F}u(\omega) = \mathcal{F}v(\omega)$ for every $\omega \in \mathbf{R}$, then $u(x) = v(x)$ for every $x \in \mathbf{R}$ where both u and v are continuous.

3.2 Special Rules

- If $u, U \in G(\mathbf{R})$ and $U(\omega) = \mathcal{F}u(\omega)$, then

$$\mathcal{F}^{-1}(U)(x) = \frac{1}{2\pi} \mathcal{F}((\mathcal{F}u)(-\omega))(x) \quad \text{and} \quad \mathcal{F}(\mathcal{F}u(\omega))(x) = 2\pi u(-x),$$

for every x where u is continuous and $D^{\pm}u(x)$ exist.

- If $u, v \in G(\mathbf{R})$ such that $uv, \mathcal{F}u, \mathcal{F}v \in G(\mathbf{R})$, then

$$\mathcal{F}(uv)(\omega) = \frac{1}{2\pi} (\mathcal{F}u * \mathcal{F}v)(\omega).$$

3.3 Plancherel's Theorem

- If $u \in G(\mathbf{R}) \cap L^2(\mathbf{R})$, then $\mathcal{F}u \in L^2(\mathbf{R})$.
- Parseval's formula: If $u, v \in G(\mathbf{R}) \cap L^2(\mathbf{R})$, then

$$\int_{-\infty}^{\infty} u(x) \overline{v(x)} dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} \mathcal{F}u(\omega) \overline{\mathcal{F}v(\omega)} d\omega.$$

3.4 Rules for the Fourier Transform

Let $u, v \in G(\mathbf{R})$ with $U(\omega) = \mathcal{F}u(\omega)$ and $V(\omega) = \mathcal{F}v(\omega)$.

3.5 Fourier Transforms

Table 3: Fourier transforms

Function	Fourier transform	Notes
e^{-ax^2}	$\sqrt{\frac{\pi}{a}} e^{-\omega^2/4a}$	$a > 0$
$e^{-a x }$	$\frac{2a}{a^2 + \omega^2}$	$a > 0$
$\operatorname{sgn}(x)e^{-a x }$	$\frac{-2i\omega}{a^2 + \omega^2}$	$a > 0$
$H(x)e^{-ax}$	$\frac{1}{a + i\omega}$	$\operatorname{Re} a > 0$
$H(-x)e^{ax}$	$\frac{1}{a - i\omega}$	$\operatorname{Re} a > 0$
$\frac{1}{a^2 + x^2}$	$\frac{\pi e^{-a \omega }}{a}$	$a > 0$
$H(x+a) - H(x-a)$	$\frac{2 \sin a\omega}{\omega}$	$a > 0$
$\operatorname{sgn}(x)(H(x+a) - H(x-a))$	$\frac{2(1 - \cos a\omega)}{i\omega}$	$a > 0$
$(a - x)(H(x+a) - H(x-a))$	$\frac{2(1 - \cos a\omega)}{\omega^2}$	$a > 0$
$\frac{1 - \cos at}{t^2}$	$\pi(a - \omega)(H(\omega+a) - H(\omega-a))$	$a > 0$

Table 2: Rules for the Fourier transform

Function	Fourier transform	Notes
$c_1 u(x) + c_2 v(x)$	$c_1 U(\omega) + c_2 V(\omega)$	
$(u * v)(x)$	$U(\omega)V(\omega)$	
$e^{iax}u(x)$	$U(\omega - a)$	$a \in \mathbf{R}$
$u(x) \cos ax$	$\frac{U(\omega - a) + U(\omega + a)}{2}$	$a \in \mathbf{R}$
$u(x) \sin ax$	$\frac{U(\omega - a) - U(\omega + a)}{2i}$	$a \in \mathbf{R}$
$u(x - x_0)$	$e^{-ix_0\omega}U(\omega)$	$x_0 \in \mathbf{R}$
$u(ax)$	$\frac{1}{ a }U\left(\frac{\omega}{a}\right)$	$a \in \mathbf{R}, a \neq 0$
$\overline{u(x)}$	$\overline{U(-\omega)}$	
$u'(x)$	$i\omega U(\omega)$	$u \in C(\mathbf{R}), u' \in G$
$u^{(k)}(x)$	$(i\omega)^k U(\omega)$	$u^{(k)} \in G(\mathbf{R})$
$x^m u(x)$	$i^m U^{(m)}(\omega)$	$x^m u(x) \in G, m = 1, 2, 3, \dots$

4 The (unilateral) Laplace Transform

The Laplace transform of $u: [0, \infty[\rightarrow \mathbf{C}$ is given by

$$\mathcal{L}u(s) = \int_0^\infty u(t)e^{-st} dt,$$

for those $s = \sigma + i\omega \in \mathbf{C}$, $\sigma, \omega \in \mathbf{R}$, where this integral is convergent.

- **Exponential growth:** A piecewise continuous $u: [0, \infty[$ is of exponential growth (of order a) if there exists constants $a > 0$ and $K > 0$ such that $|u(t)| \leq Ke^{at}$ for $t \geq 0$. The set of all such functions will be denoted by X_a .
- **Existence of $\mathcal{L}u(s)$:** If $u \in X_a$ for some $a > 0$, then the Laplace transform $\mathcal{L}u(s)$ exists (at least) for $\operatorname{Re} s > a$.
- $\mathcal{L}u(s) \rightarrow 0$ as $\mathbf{R} \ni s \rightarrow \infty$.
- $\mathcal{L}u(s)$ converges uniformly for $\operatorname{Re} s > a$.
- $\mathcal{L}u(s)$ is analytic for $\operatorname{Re} s > a$.
- **Periodicity:** If there exists $T > 0$ such that $u(t+T) = u(t)$ for every $t \geq 0$, then

$$\mathcal{L}u(s) = \frac{1}{1 - e^{-sT}} \int_0^T u(\tau)e^{-s\tau} d\tau.$$

4.1 Inversion

- If $u \in X_a$ has right- and lefthand limits at a point $t > 0$, then

$$\lim_{L \rightarrow \infty} \frac{1}{2\pi} \int_{-L}^L \mathcal{L}u(\sigma + i\omega) e^{\sigma t} e^{i\omega t} d\omega = \frac{u(t^+) + u(t^-)}{2},$$

where the vertical line $\text{Re } z = \sigma$ is contained in the region of convergence of $\mathcal{L}u(s)$

- If $u, v \in X_a$ and $\mathcal{L}u(s) = \mathcal{L}v(s)$ on some vertical line $\text{Re } s = \sigma$, then $u(t) = v(t)$ for all t where u and v are continuous.

4.2 Limit Theorems

- **Final value theorem:**
If $u: [0, \infty[\rightarrow \mathbf{C}$ is bounded and $\lim_{t \rightarrow \infty} u(t) = A$, then $A = \lim_{s \rightarrow 0^+} s \mathcal{L}u(s)$.
- **Initial value theorem:**
If $u: [0, \infty[\rightarrow \mathbf{C}$ belongs to X_b and $\lim_{t \rightarrow 0^+} u(t) = a$, then $a = \lim_{s \rightarrow \infty} s \mathcal{L}u(s)$.

4.3 Rules for the Laplace Transform

Let $U(s) = \mathcal{L}u(t)$, $\sigma > \sigma_u$ and $V(s) = \mathcal{L}v(t)$, $\sigma > \sigma_v$.

Table 4: Rules for Laplace transforms

Function	Unilateral Laplace transform	Region of convergence
$c_1 u(t) + c_2 v(t)$	$c_1 U(s) + c_2 V(s)$	$\sigma > \max\{\sigma_u, \sigma_v\}$
$(u * v)(t)$	$U(s)V(s)$	unilateral conv. [†] ; $\sigma > \max\{\sigma_u, \sigma_v\}$
$e^{at}u(t)$	$U(s - a)$	$\sigma > \sigma_u + \text{Re } a$
$u(t - t_0)H(t - t_0)$	$e^{-ts}U(s)$	$\sigma > \sigma_u, t_0 > 0$
$u(at)$	$\frac{1}{a}U\left(\frac{s}{a}\right)$	$\sigma > a\sigma_u, a > 0$
$\overline{u(t)}$	$\overline{U(\overline{s})}$	$\sigma > \sigma_u$
$u'(t)$	$sU(s) - u(0)$	$\sigma > \sigma_u$
$u^{(n)}(t)$	$s^n U(s) - s^{n-1}u(0) - \dots$ $\dots - s u^{(n-2)}(0) - u^{(n-1)}(0)$	$\sigma > \max\{\sigma_u, \sigma_{u'}, \dots, \sigma_{u^{(n-1)}}\}$
$\int_0^t u(\tau) d\tau$	$\frac{U(s)}{s}$	$\sigma > \max\{\sigma_u, 0\}$
$t^m u(t)$	$(-1)^m U^{(m)}(s)$	$\sigma > \sigma_u$

[†] $(u * v)(t) = \int_0^t u(\tau)v(t - \tau) d\tau$.

4.4 Laplace Transforms

Table 5: Laplace transforms

Function	Unilateral Laplace transform	Region of convergence
$H(t) = 1$	$\frac{1}{s}$	$\sigma > 0$
t	$\frac{1}{s^2}$	$\sigma > 0$
t^m	$\frac{m!}{s^{m+1}}$	$\sigma > 0$
t^a	$\frac{\Gamma(a+1)}{s^{a+1}}$	$m = 1, 2, 3, \dots$ $\sigma > 0$
e^{at}	$\frac{1}{s - a}$	$a > 0$ $\sigma > \text{Re } a$
$t^m e^{at}$	$\frac{m!}{(s - a)^{m+1}}$	$\sigma > \text{Re } a$ $m = 1, 2, 3, \dots$
$\cos at$	$\frac{s}{s^2 + a^2}$	$\sigma > \text{Im } a $
$t \cos at$	$\frac{s^2 - a^2}{(s^2 + a^2)^2}$	$\sigma > \text{Im } a $
$\sin at$	$\frac{a}{s^2 + a^2}$	$\sigma > \text{Im } a $
$t \sin at$	$\frac{2as}{(s^2 + a^2)^2}$	$\sigma > \text{Im } a $
$\frac{\sin at}{t}$	$\arctan\left(\frac{a}{s}\right)$	$\sigma > \text{Im } a $
$\cosh at$	$\frac{s}{s^2 - a^2}$	$\sigma > \text{Re } a $
$\sinh at$	$\frac{a}{s^2 - a^2}$	$\sigma > \text{Re } a $
$J_0(at)$	$\frac{1}{\sqrt{a^2 + s^2}}$	$\sigma > \text{Im } a $

5 The (unilateral) Z Transform

The **Z transform** of a sequence $u[k]$, $k = 0, 1, 2, \dots$, is defined by

$$\mathcal{Z}(u)(z) = \sum_{k=0}^{\infty} u[k] z^{-k},$$

for those $z = x + iy \in \mathbf{C}$, $x, y \in \mathbf{R}$, where this series is absolutely convergent.

5.2 Z Transforms

- **Existence of $\mathcal{Z}u(z)$:** For a sequence $u[k]$, $k = 0, 1, 2, \dots$, the Z transform $\mathcal{Z}u(z)$ has a region of convergence R such that $\mathcal{Z}u(z)$ is absolutely (uniformly) convergent for $|z| > R$ and divergent for $|z| < R$. It is possible that $R = 0$ or $R = \infty$.

- **Inversion.** If $U(z) = \mathcal{Z}u(z)$, then

$$u[k] = \frac{1}{2\pi i} \oint_{\gamma} z^{k-1} U(z) dz, \quad k = 0, 1, 2, \dots,$$

where γ is a closed curve (inside $|z| > R_u$) counterclockwise around the origin.

- **Uniqueness:** If $\mathcal{Z}u(z) = \mathcal{Z}v(z)$ for all $|z| > R$ for some $R > 0$, then $u[k] = v[k]$ for $k = 0, 1, 2, \dots$
- **Initial value theorem:** If there's an $R > 0$ such that $\mathcal{Z}u(z)$ exists for $|z| > R$, then

$$\lim_{|z| \rightarrow \infty} \mathcal{Z}u(z) = u[0].$$

5.1 Rules for the Z Transform

Let $U(z) = \mathcal{Z}(u[k])(z)$, $|z| > R_u$ and $V(z) = \mathcal{Z}(v[k])(z)$, $|z| > R_v$.

Table 6: Rules for Z transforms

Function	Unilateral Z transform	Region of convergence
$c_1 u[k] + c_2 v[k]$	$c_1 U(z) + c_2 V(z)$	$ z > \max\{R_u, R_v\}$
$(u * v)[k]$	$U(z)V(z)$	unilateral conv. [†] ; $ z > \max\{R_u, R_v\}$
$a^k u[k]$	$U\left(\frac{z}{a}\right)$	$ z > a R_u, a \in \mathbf{C} \setminus \{0\}$
$u[k-m]H[k-m]$	$z^{-m}U(z)$	$ z > R_u, m = 1, 2, 3, \dots$
$u[k-m]$	$z^{-m}U(z) + z^{-m+1}u[-1] + \dots$ $\dots + z^{-1}u[-m+1] + u[-m]$	$ z > R_u, m = 1, 2, 3, \dots$
$u[k+m]$	$z^m U(z) - z^m u[0] + \dots$ $\dots - z^2 u[m-2] - zu[m-1]$	$ z > R_u, m = 1, 2, 3, \dots$
$\overline{u[k]}$	$\overline{U(\bar{z})}$	$ z > R_u$
$\sum_{l=0}^k u[l]$	$\frac{z}{z-1} U(z)$	$ z > \max\{R_u, 1\}$
$k^m u[k]$	$\left(-z \frac{d}{dz}\right)^m U(z)$	$ z > R_u$

[†] $(u * v)[k] = \sum_{l=0}^k u[l]v[k-l]$.

Table 7: Z transforms

Function	Unilateral Z transform	Region of convergence
$\delta[k]$	1	$z \in \mathbf{C}$
$\delta[k-m]$	z^{-m}	$ z > 0, m = 1, 2, \dots$
$H[k]$	$\frac{z}{z-1}$	$ z > 1$
k	$\frac{z}{(z-1)^2}$	$ z > 1$
a^k	$\frac{z}{z-a}$	$ z > a $
ka^k	$\frac{az}{(z-a)^2}$	$ z > a $
$k^2 a^k$	$\frac{az^2 + a^2 z}{(z-a)^3}$	$ z > a $
$k^3 a^k$	$\frac{az^3 + 4a^2 z^2 + a^3 z}{(z-a)^4}$	$ z > a $
$(k+1)a^k$	$\frac{z^2}{(z-a)^2}$	$ z > a $
$\binom{k+m}{m} a^k$	$\frac{z^{m+1}}{(z-a)^{m+1}}$	$ z > a , m = 2, 3, \dots$
$\binom{k}{m} a^k$	$\frac{a^m z}{(z-a)^{m+1}}$	$ z > a , m = 2, 3, \dots$
$\binom{k+n}{m} a^k$	$\frac{a^{m-n} z^{n+1}}{(z-a)^{m+1}}$	$ z > a , m = 2, 3, \dots$ $n = 1, \dots, m-1$
$\cos k\alpha$	$\frac{z^2 - z \cos \alpha}{z^2 - 2z \cos \alpha + 1}$	$ z > 1$
$\sin k\alpha$	$\frac{z \sin \alpha}{z^2 - 2z \cos \alpha + 1}$	$ z > 1$
$k \cos k\alpha$	$\frac{z^3 \cos \alpha - 2z^2 + z \cos \alpha}{(z^2 - 2z \cos \alpha + 1)^2}$	$ z > 1$
$k \sin k\alpha$	$\frac{z^3 \sin \alpha - z \sin \alpha}{(z^2 - 2z \cos \alpha + 1)^2}$	$ z > 1$
$\frac{a^k}{k!}$	$e^{a/z}$	$ z > 0$
$\frac{1}{k} H[k-1]$	$\ln \frac{z}{z-1}$	$ z > 1$
$\binom{n}{k} a^k b^{n-k}$	$\frac{(bz+a)^n}{z^n}$	$ z > 0, n = 1, 2, 3, \dots$