

ODEs and Dynamical Systems (TATA71)

Course programme, fall 2018

General information

Ordinary Differential Equations and Dynamical Systems (TATA71) is an optional course for Mat2, Y4, M4, EMM4. It is given in the second half of the fall semester (period ht2). All information is publicly available on the course webpage courses.mai.liu.se/GU/TATA71/. Lisam is currently not used in this course.

Literature

D. K. Arrowsmith & C. M. Place, *Dynamical Systems: Differential Equations, Maps, and Chaotic Behaviour* Chapman and Hall/CRC (1992), ISBN 9780412390807.

Prerequisites

Basic courses in **single-variable** and **multi-variable calculus**, plus **linear algebra**. (An honours course in **real analysis** may be helpful for understanding certain subtle details, but that's optional.)

Teaching

There will be 11 lectures and 11 lessons (problem-solving classes). See the [lecture plan](#) below.

As always, it is a good idea to read (or at least skim) the relevant sections in the textbook before each lecture, and also to try out a few of the problems before each lesson, since there will usually not be time enough to do all of them in the classroom.

Examination

The examination consists of two parts:

- UPG1. Homework assignments, worth 2 hp (= 2 ECTS credits).

Some of the exercises for the lessons are assigned as homework problems (underlined in the programme), to be handed in continually during the course. These problems are only graded pass/fail, and if you fail a problem, you simply hand in a corrected version later. It is allowed to discuss the problems with the teacher and with your fellow students, but please write the solution in your own words, don't just copy someone else's solution! Handwritten solutions are fine, and you can write them in English or in Swedish.

Since this is a small course with few participants, there is probably no need to set any strict deadlines, but the idea is that the homework problems are a way of giving you feedback, so the sooner you hand them in, the more useful they will be to you. And the intent is also that all problems should be completed before the Christmas break.

- TEN1. A written test (5 hours), worth 4 hp.

The test contains 6 problems, each of which is graded as pass (3 or 2 points) or fail (1 or 0 points). The total grade for the course is determined by the grade for the written test, which in turn is determined as follows: for grade 3/4/5 (respectively), you need 3/4/5 passed problems and in addition at least 8/11/14 points in total.

What's this course about?

As the name of the course suggests, we will study **ODEs** and **dynamical systems**.

- **ODE** is a standard abbreviation for **ordinary differential equation**, where the function that we seek depends on *one* variable. So an ODE is just a good old differential equation like those which you have already seen in your single-variable calculus course. We will also encounter *systems* of ODEs, involving several unknown functions at once, but each function will still only depend on one variable.

In contrast, a **partial differential equation (PDE)** is a differential equation where one seeks a function depending on *several* variables, so that *partial derivatives* come into play.

- The idea of a **dynamical system** is rather broad, and it is hard to give a precise mathematical definition which would cover every possible use of the phrase. But it refers to a “system” (whatever that is) which changes in a deterministic way as time passes, and which is “memoryless” in the sense that (at any given instant) the future of the system is uniquely determined by the present state alone; the past is irrelevant.

We will usually assume when talking about dynamical systems that the laws governing the evolution don't change with time. That is, if we start the system in a given state T units of time from now, we will get the same evolution as if we start it in that state right now (except that everything will be delayed by T time units of course). If this needs to be emphasized, one uses the phrase **autonomous dynamical system** – a system which “runs on its own”, in contrast to **non-autonomous** dynamical systems where there may be some external time-dependent factors which influence the evolution of the system.

The state of the system is represented mathematically by an element of a set called the **state space**, typically \mathbf{R}^n , or maybe some subset of \mathbf{R}^n (like a cylinder, a sphere, or a torus). So one pictures the evolution of the system as the motion of a point in phase space.

- A **discrete-time dynamical system** is one where things happen at distinct time steps. We can use integers to label the time steps, so that the system is in the state $x_n \in S$ at time $n \in \mathbf{Z}$, where S is the state space. Then the evolution of the system is simply specified by some function $f: S \rightarrow S$:

$$x_{n+1} = f(x_n), \quad n \in \mathbf{Z}.$$

(The system is autonomous since the function f is the same for all n .)

Discrete-time dynamical systems are a very important part of the general theory of dynamical systems, but will not be encountered very much in this course.

- In a **continuous-time dynamical system**, time passes smoothly, so we use real numbers to describe time, and talk about the system being in the state $x(t)$ at time $t \in \mathbf{R}$. In this case, if the state space is \mathbf{R}^n for simplicity, the evolution is determined by a system of first-order ODEs for the state $\mathbf{x}(t) = (x_1(t), \dots, x_n(t))$:

$$\begin{aligned} dx_1/dt &= f_1(x_1, \dots, x_n), \\ dx_2/dt &= f_2(x_1, \dots, x_n), \\ &\vdots \\ dx_n/dt &= f_n(x_1, \dots, x_n), \end{aligned}$$

or simply $d\mathbf{x}/dt = \mathbf{f}(\mathbf{x})$ for short, with $\mathbf{x} \in \mathbf{R}^n$ and $\mathbf{f}: \mathbf{R}^n \rightarrow \mathbf{R}^n$. (The system is autonomous since the function \mathbf{f} doesn't depend on t .)

In order for this to really define a dynamical system, we must impose suitable conditions on \mathbf{f} which will guarantee **existence** and **uniqueness** of the solution to the ODEs for a given initial condition, so that the future is uniquely determined by the present state.

Plan for lectures and lessons

Lecture 1. Basics of first-order ODEs	4
Lesson 1	6
Lecture 2. Phase portraits for two-dimensional systems	7
Lesson 2	8
Lecture 3. Two-dimensional linear systems	8
Lesson 3	9
Lecture 4. More about linear systems	10
Lesson 4	11
Lecture 5. Nonlinear systems, linearization at an equilibrium point	12
Lesson 5	13
Lecture 6. Stability theorems	14
Lesson 6	19
Lecture 7. Limit sets	20
Lesson 7	23
Lecture 8. Some applications	24
Lesson 8	25
Lecture 9. More about existence and uniqueness	25
Lesson 9	31
Lecture 10. Linear equations with non-constant coefficients	33
Lesson 10	39
Lecture 11. Outlook: Poincaré maps, attractors, chaotic systems	40
Lesson 11	40

Lecture 1. Basics of first-order ODEs

(Arrowsmith & Place, sections 1.1, 1.2. And your old calculus textbook, if needed.)

This lecture introduces some basic ideas:

- **Existence and uniqueness theorems** for first-order ODEs $\frac{dx}{dt} = X(t, x)$.

Proposition 1.1.1 is usually called **Peano's existence theorem**.

Proposition 1.1.2 is a slightly simplified version of the **Picard–Lindelöf theorem**. (The assumption “ $\partial X/\partial x$ exists and is continuous” is stronger than necessary; we’ll study this more thoroughly later in the course.)

- How to find **explicit solutions** in simple cases.
(This is mostly a question of remembering the methods you have learned in previous courses. See summary below.)
- How to sketch the **solution curves** directly from the ODE.
- How to draw the **phase portrait** for a one-dimensional dynamical system.

A comment about notation

You might be used to ODEs looking something like this:

$$y''(x) + 5y'(x) + 4y(x) = \cos x \quad (\text{or simply } y'' + 5y' + 4y = \cos x),$$

where the independent variable is called x , and $y = y(x)$ is the function that we seek. But since this is a course about ODEs with a “dynamical systems perspective”, we will instead call the independent variable t , for “time”, and use names like $x(t)$ or $y(t)$ for the sought functions. So the same ODE now instead looks as follows:

$$x''(t) + 5x'(t) + 4x(t) = \cos t \quad \text{or} \quad \ddot{x}(t) + 5\dot{x}(t) + 4x(t) = \cos t.$$

(It is common to use **dots** instead of primes to denote **derivatives with respect to time**.)

Two very fundamental examples

- **Exponential growth/decay:**

$$x'(t) = r x(t).$$

This is a linear equation, and we can solve it in several ways: integrating factor, characteristic polynomial, separation of variables. Either way, the solution with initial condition $x(0) = x_0$ is

$$x(t) = x_0 e^{rt}.$$

This will be encountered again and again in this course, and you will be expected to *instantly* recognize this equation and know its solution. Phase portrait (if $r > 0$): “ $\leftarrow 0 \rightarrow$ ”.

- The **logistic equation** with growth rate r and carrying capacity K :

$$x'(t) = r x(t) \left(1 - \frac{x(t)}{K}\right).$$

This nonlinear equation is often solved via separation of variables (followed by integration using partial fractions), but an easier way is to use the substitution $x(t) = 1/y(t)$, since this is a Bernoulli equation. Solution, with $x(0) = x_0$:

$$x(t) = \frac{K x_0}{x_0 + (K - x_0) e^{-rt}} = \frac{K x_0 e^{rt}}{K + (e^{rt} - 1) x_0}.$$

Here you don’t need to memorize the solution formula, but you should know roughly what the *graph* of the solution $x(t)$ looks like for different values of x_0 . (In particular, $x(t) = 0$ and $x(t) = K$ are constant solutions.) Phase portrait (if $r > 0$ and $K > 0$): “ $\leftarrow 0 \rightarrow K \leftarrow$ ”.

Summary of some exact solution methods

- **Linear first order equations** $x' + ax = b$, where the coefficients a and b may be functions of the time variable t :

$$x'(t) + a(t)x(t) = b(t).$$

How to solve: Find an antiderivative of $a(t)$; call it $A(t)$. Then multiply both sides of the ODE by the **integrating factor** $e^{A(t)}$, and use the product rule for derivatives (backwards). This gives

$$(e^{A(t)} x(t))' = e^{A(t)} b(t),$$

which can now be integrated.

- **Separable equations** $f(x) x' = g(t)$.

What this means is that we seek $x(t)$ such that

$$f(x(t)) x'(t) = g(t).$$

Integrating this with respect to t , using the chain rule (backwards), we immediately obtain the solution in the implicit form

$$F(x(t)) = G(t) + C,$$

where $F(x)$ is some antiderivative of $f(x)$ and $G(t)$ is some antiderivative of $g(t)$. Usually this is remembered via the trick of writing $x' = dx/dt$ and “separating the variables” by “multiplying by dt ”, and then attaching integral signs:

$$f(x) \frac{dx}{dt} = g(t) \quad \Leftrightarrow \quad \int f(x) dx = \int g(t) dt.$$

It's sometimes convenient to use *definite* integrals instead, particularly if we want to find a solution satisfying a given **initial condition** $x(t_0) = x_0$:

$$\int_{x_0}^{x(t)} f(\xi) d\xi = \int_{t_0}^t g(\tau) d\tau,$$

or in other words

$$F(x(t)) - F(x_0) = G(t) - G(t_0).$$

As an important special case of separable equations we have **one-dimensional dynamical systems** $x' = a(x)$, which can always be solved by separation of variables:

$$x' = a(x) \quad \Leftrightarrow \quad x(t) = x^* \quad \text{where } a(x^*) = 0$$

$$\text{or} \quad \int \frac{dx}{a(x)} = \int dt = t + C.$$

Warning! This method is full of pitfalls! Don't forget the case $a(x) = 0$; there is a constant solution $x(t) = x^*$ for each zero x^* of the function $a(x)$. And one should also be *very* careful with handling logarithms and absolute values correctly when integrating and when simplifying the solution. And in principle the constant of integration may be different in the different intervals into which the real line is divided by the zeros of $a(x)$. So if the ODE can be solved by some other method, it may be wise to try that method first.

- **Linear equations of arbitrary order:**

$$x^{(n)}(t) + a_{n-1}(t)x^{(n-1)}(t) + \dots + a_2(t)x''(t) + a_1(t)x'(t) + a_0(t)x(t) = b(t).$$

The general solution of such an equation has the structure

$$x(t) = x_{\text{hom}}(t) + x_{\text{part}}(t)$$

where $x_{\text{part}}(t)$ is a particular solution and $x_{\text{hom}}(t)$ is the general solution of the corresponding homogeneous equation which has 0 instead of $b(t)$ on the right-hand side.

Solving higher-order equations with **time-dependent coefficients** is usually a rather hopeless task, except that one may try to find solutions in the form of **power series**.

But it's fairly easy in the special case of **constant coefficients** $a_k(t) = c_k$:

$$x^{(n)}(t) + c_{n-1}x^{(n-1)}(t) + \dots + c_2x''(t) + c_1x'(t) + c_0x(t) = b(t).$$

Then $x_{\text{hom}}(t)$ can be found by looking at the roots of the **characteristic polynomial**

$$p(r) = r^n + c_{n-1}r^{n-1} + \dots + c_2r^2 + c_1r + c_0.$$

(If the roots are distinct and real, it's easy. Repeated and/or complex roots are a bit harder.) To find $x_{\text{part}}(t)$ one usually makes a suitable *Ansatz*. See your calculus book for details about all this.

- If we start with $x'(t) + a(t)x(t) = b(t)$, which we know that we can solve using an integrating factor, and let $x(t) = 1/y(t)^k$ (for some $k \neq 0$), we get $x'(t) = -ky'(t)/y(t)^{k+1}$, and consequently $y(t)$ satisfies the ODE

$$-k \frac{y'(t)}{y(t)^{k+1}} + a(t) \frac{1}{y(t)^k} = b(t),$$

that is,

$$y'(t) - \frac{a(t)}{k} y(t) = \frac{b(t)}{k} y(t)^{k+1}$$

The conclusion is that if we encounter an equation of this form, a so-called **Bernoulli equation**

$$y'(t) + p(t)y(t) = q(t)y(t)^n,$$

then we can reduce it to a linear equation by letting $x(t) = 1/y(t)^{n-1}$.

- Later in this course we will thoroughly study **linear first order systems**

$$\mathbf{x}'(t) = A(t)\mathbf{x}(t),$$

where $\mathbf{x}(t) = (x_1(t), \dots, x_n(t))^T$ is a column vector of functions and $A(t)$ is matrix of size $n \times n$. But you may already have seen in your linear algebra course how to solve such a system in the case where A is a **constant** matrix which happens to be **diagonalizable**: set $\mathbf{x} = M\mathbf{y}$, where the columns of M are linearly independent eigenvectors of A , and you get a decoupled system $\mathbf{y}'(t) = D\mathbf{y}(t)$, where D is a diagonal matrix with the eigenvalues of A along the diagonal.

Lesson 1

The problems labelled A1, A2, etc., can be found in the section **Additional problems** just below. The remaining problems (1.1, 1.2, etc.) are from the **course textbook** (Arrowsmith & Place), where they are located at the end of each chapter. Problems marked with an **asterisk** might be a bit more challenging.

- Solving ODEs using calculus techniques: 1.1, 1.2, 1.4, [A1](#), [A2](#).
- Sketching solution curves directly from the ODE: 1.11.
- Drawing phase portraits: 1.12, 1.13.
- Phase portraits for parameter-dependent ODEs: 1.14, 1.17*.
- Recovering the ODE from the family of solution curves: [A3](#).

Additional problems for lesson 1

A1 Find the general solution of the following constant-coefficient second-order ODEs:

(a) $\ddot{x} + 6\dot{x} + 8x = t + 2e^{-2t}$ [Answer: $x(t) = Ae^{-2t} + Be^{-4t} + (4t - 3)/32 + te^{-2t}$.]

(b) $\ddot{x} + 6\dot{x} + 9x = 0$ [Answer: $x(t) = (At + B)e^{-3t}$.]

(c) $\ddot{x} + 6\dot{x} + 10x = 2e^{-3t} \cos t$ [Answer: $x(t) = e^{-3t}(A \cos t + B \sin t) + te^{-3t} \sin t$.]

A2 The *Airy equation* $\ddot{x}(t) = tx(t)$ is a second-order ODE with non-constant coefficients. Find the solution which satisfies $x(0) = 1$ and $\dot{x}(0) = 0$, in the form of a power series $x(t) = \sum_{k=0}^{\infty} a_k t^k$.

[Answer: $x(t) = 1 + \frac{t^3}{2 \cdot 3} + \frac{t^6}{2 \cdot 3 \cdot 5 \cdot 6} + \frac{t^9}{2 \cdot 3 \cdot 5 \cdot 6 \cdot 8 \cdot 9} + \dots$]

A3 A first-order ODE (usually) has a general solution consisting of a one-parameter family of curves. Here's an opposite problem: Find an ODE $\dot{x} = X(t, x)$ such that the one-parameter family of curves $t^2 + x^2 = 2Ct$ are solution curves.

(Hint: Differentiate $t^2 + x(t)^2 = 2Ct$ with respect to t , and eliminate the parameter C from the two equations.)

[Answer: $\dot{x} = (x^2 - t^2)/2tx$.]

Lecture 2. Phase portraits for two-dimensional systems

(Arrowsmith & Place, sections 1.3, 1.4, 1.5.)

Now we turn to **two-dimensional** dynamical systems

$$\dot{x}_1 = X_1(x_1, x_2), \quad \dot{x}_2 = X_2(x_1, x_2),$$

where we assume that the functions X_1 and X_2 are nice enough to guarantee (local) existence and uniqueness of solutions. In vector notation: $\dot{\mathbf{x}}(t) = \mathbf{X}(\mathbf{x}(t))$, or simply $\dot{\mathbf{x}} = \mathbf{X}(\mathbf{x})$. The function \mathbf{X} should be thought of as a **vector field** in phase space: a vector $\mathbf{X}(\mathbf{x})$ is prescribed at each point \mathbf{x} . The solution curves for the system are curves $\mathbf{x} = \mathbf{x}(t)$ whose tangent vector $\dot{\mathbf{x}}$ agrees with the prescribed vector $\mathbf{X}(\mathbf{x})$ at each point on the curve. The solution curves are sometimes called **flow lines** of the vector field.

- In very simple cases, we can solve the system explicitly. Sometimes we can obtain partial information by methods of calculus.
- But we can also get at least a rough idea of what the phase portrait looks like by direct inspection of the signs of the functions $X_1(x_1, x_2)$ and $X_2(x_1, x_2)$.
- Syntax for drawing phase portraits in Mathematica or Wolfram Alpha (www.wolframalpha.com):

StreamPlot[$X_1(x, y), X_2(x, y), \{\mathbf{x}, x_{\min}, x_{\max}\}, \{\mathbf{y}, y_{\min}, y_{\max}\}$]

For example:

StreamPlot[-1-x^2+y, 1+x-y^2, {x, -3, 3}, {y, -3, 3}]

(But don't expect an automatic command like this to produce anything nearly as good as the hand-tuned graphics in the textbook.)

- Arrowsmith & Place use the phrase **fixed point** for a point \mathbf{x}^* such that $\mathbf{X}(\mathbf{x}^*) = \mathbf{0}$, but I will usually say **equilibrium point** or just **equilibrium**, and you may come across many other synonyms too: **rest point**, **critical point**, **steady state**, etc.

The reason for the terminology is of course that $\mathbf{x}(t) = \mathbf{x}^*$ is a constant solution of the system $\dot{\mathbf{x}} = \mathbf{X}(\mathbf{x})$ in this situation. That is, a system starting in the state \mathbf{x}^* remains in the state \mathbf{x}^* forever.

- The **evolution operator** or **flow** φ is the function which maps each point \mathbf{x} in phase space to the place where it will be t units of time later if moving as prescribed by the dynamical system. It is denoted by $\varphi_t(x)$ in the book, but it's also common to write $\varphi(t, x)$, since it is simply a function of t and x .

A fact which isn't mentioned in the textbook is that the flow φ is as "nice" as the vector field \mathbf{X} . More precisely, there's a theorem which says that if the system is

$$\dot{\mathbf{x}} = \mathbf{X}(\mathbf{x}, \boldsymbol{\lambda}),$$

where $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_N)$ is some vector of parameters, and if \mathbf{X} is of class C^k ($1 \leq k \leq \infty$) as a function of the variables $(\mathbf{x}, \boldsymbol{\lambda})$, then the flow φ is of class C^k as a function of the variables $(t, \mathbf{x}, \boldsymbol{\lambda})$.

Lesson 2

Underlined problems are **homework** problem to be handed in.

- Finding fixed points: 1.19ac.
In these problems, also find the **nullclines** (the curves where $\dot{x}_1 = 0$ or $\dot{x}_2 = 0$) and use this to draw the regions of the phase plane where \dot{x}_1 and \dot{x}_2 are positive/negative).
- And another one for homework: 1.19d (including the follow-up question above about nullclines and signs of \dot{x}_1 and \dot{x}_2 .)
- Drawing families of parametrized curves and finding the corresponding ODEs: 1.20.
- Exact solution of a linear systems: 1.23 (draw the phase portrait both in the $y_1 y_2$ -plane and in the $x_1 x_2$ -plane).
- An easy verification: 1.24.
- A nonlinear system which can be solved using polar coordinates: 1.25.
- Flows: 1.32, 1.36.
(Hint for 1.32, if you want to solve it from scratch: it's a Bernoulli equation, let $y = 1/x^2$. Or just use separation of variables and partial fractions directly. Or else *verify* that the flow is correct, by computing $\frac{d}{dt}\varphi_t(x_0)$.)

Lecture 3. Two-dimensional linear systems

(Arrowsmith & Place, sections 2.1, 2.2, 2.3.)

We will spend some time on understanding **linear** dynamical systems in **two dimensions**:

$$\begin{aligned} \dot{x}_1 &= ax_1 + bx_2 \\ \dot{x}_2 &= cx_1 + dx_2 \end{aligned} \iff \begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \end{pmatrix} = \underbrace{\begin{pmatrix} a & b \\ c & d \end{pmatrix}}_{=A} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}.$$

- Definition of a **simple** linear system: $\det(A) = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$ is **nonzero**. (Equivalently: the eigenvalues of A are nonzero.) This implies that the origin $(x_1, x_2) = (0, 0)$ is the **only** equilibrium point. In this lecture we'll only consider simple systems.
- A linear change of variables $\mathbf{x} = M\mathbf{y}$ turns the system $\dot{\mathbf{x}} = A\mathbf{x}$ into $\dot{\mathbf{y}} = M^{-1}AM\mathbf{y}$.
- "Recipe" for choosing the columns \mathbf{m}_1 and \mathbf{m}_2 in the matrix M so that $J = M^{-1}AM$ simplifies to the **Jordan normal form** of A :

- (a) Suppose A has two distinct real eigenvalues $\lambda_1 > \lambda_2$. Taking \mathbf{m}_1 and \mathbf{m}_2 to be the corresponding eigenvectors gives $J = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$.
- (b) Suppose A has a double real eigenvalue λ_0 . Then either $A = \begin{pmatrix} \lambda_0 & 0 \\ 0 & \lambda_0 \end{pmatrix}$ is already in normal form, or else there is just a one-dimensional eigenspace; in this case let \mathbf{m}_1 be an eigenvector and take any vector not parallel to \mathbf{m}_1 as our preliminary \mathbf{m}_2 . This will give a preliminary $J = \begin{pmatrix} \lambda_0 & C \\ 0 & \lambda_0 \end{pmatrix}$ with some nonzero constant C . Adjust \mathbf{m}_2 by dividing it by this constant C ; this will give the Jordan form $J = \begin{pmatrix} \lambda_0 & 1 \\ 0 & \lambda_0 \end{pmatrix}$.
- (c) Suppose A has a complex-conjugated pair of eigenvalues $\lambda_{1,2} = \alpha \pm i\beta$ with $\beta > 0$. This case is done in a very complicated way in the book (p. 39)! A much simpler way is to let $\mathbf{a} + i\mathbf{b}$ (with \mathbf{a} and \mathbf{b} real) be a complex eigenvector corresponding to the eigenvalue $\lambda_1 = \alpha + i\beta$. Then $\mathbf{m}_1 = \mathbf{b}$ and $\mathbf{m}_2 = \mathbf{a}$ will work, and directly give the Jordan form $J = \begin{pmatrix} \alpha & -\beta \\ \beta & \alpha \end{pmatrix}$. (These vectors are automatically linearly independent – otherwise they would both be *real* eigenvectors of the real matrix A , with a *non-real* eigenvalue λ_1 , impossible.)

- How to solve the system $\dot{\mathbf{y}} = J\mathbf{y}$ with a matrix in Jordan form, and how to draw its phase portrait.

In case (a) the phase portrait is a **node** or a **saddle** depending on the signs of the eigenvalues. In case (b) it is a **star node** or an **improper node**. And in case (c) it is a **spiral** (also called **focus**) if $\alpha \neq 0$, or a **centre** if $\alpha = 0$.

The phase portrait for $\mathbf{x} = M\mathbf{y}$ will be of the same type, only distorted by the linear transformation M . The columns \mathbf{m}_1 and \mathbf{m}_2 give the **principal directions** of the phase portrait.

- A solution method which isn't mentioned in the book, but is sometimes convenient, is to **rewrite the system as a single second-order ODE** with constant coefficients, as follows. The derivative of $\dot{x}_1 = ax_1 + bx_2$ is

$$\ddot{x}_1 = a\dot{x}_1 + b\dot{x}_2.$$

The second term here, $b\dot{x}_2$, can be rewritten using the equations from the system:

$$b\dot{x}_2 = b(cx_1 + dx_2) = bcx_1 + d(bx_2) = bcx_1 + d(\dot{x}_1 - ax_1) = d\dot{x}_1 - (ad - bc)x_1.$$

So we find that $\ddot{x}_1 = a\dot{x}_1 + (d\dot{x}_1 - (ad - bc)x_1)$, or in other words

$$\ddot{x}_1 - (a + d)\dot{x}_1 + (ad - bc)x_1 = 0.$$

This second-order ODE for $x_1(t)$ can now be **solved as in calculus**, with the help of its characteristic polynomial

$$p(\lambda) = \lambda^2 - (a + d)\lambda + (ad - bc),$$

which (perhaps not surprisingly) coincides with the characteristic polynomial of the system's matrix A , as we can easily check:

$$p(\lambda) = \det(A - \lambda I) = \begin{vmatrix} a - \lambda & b \\ c & d - \lambda \end{vmatrix} = (a - \lambda)(d - \lambda) - bc = \lambda^2 - (a + d)\lambda + (ad - bc).$$

And once $x_1(t)$ is known, we also get $x_2(t)$ from $x_2 = (\dot{x}_1 - ax_1)/b$, assuming that $b \neq 0$. (If $b = 0$, the system is easy to solve right away: the first equation $\dot{x}_1 = ax_1 + 0x_2$ gives $x_1(t) = x_1(0)e^{at}$; plug this into the second equation and solve it using an integrating factor or $(x_2)_{\text{hom}} + (x_2)_{\text{part}}$.)

Lesson 3

- A little reminder of how linear transformations work: 2.1.
- Transformation to canonical form: 2.3, 2.4.
- Solving canonical linear systems (and drawing the phase portrait): 2.8, 2.9.


Lecture 4. More about linear systems

(Arrowsmith & Place, sections 2.4, 2.5, 2.6, 2.7.)

- How to tell directly from the coefficients of the matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ what the type of the phase portrait for $\dot{\mathbf{x}} = A\mathbf{x}$ is, without computing the eigenvalues: compute the **trace** and the **determinant**,

$$\beta = \text{tr}(A) = a + d, \quad \gamma = \det(A) = ad - bc,$$

and locate the point (β, γ) in the following diagram (cf. Figure 2.7 in the book):



For systems on the parabola $\gamma = (\beta/2)^2$, the type is a star node if the matrix A equals a constant times the identity matrix, otherwise it's an improper node.

Centres are always stable, and saddles are always unstable (according to the general definitions of “stable equilibrium” and “unstable equilibrium” that will be given later in the course).

- In particular, the zeros of the polynomial $p(z) = z^2 - \beta z + \gamma$ both have *negative real part* if and only if the coefficients satisfy $\beta < 0$ and $\gamma > 0$, i.e., if the point (β, γ) lies in the second quadrant in the diagram above. (By the way, this is a special case of the **Routh–Hurwitz criterion**, which determines for a polynomial of arbitrary degree whether all its zeros have negative real part.)
- The word “type” above refers to a kind of “algebraic type”, two linear systems being of the same type if and only if they are related via a **linear** change of variables. Such changes preserve the eigenvalues, so (for example) an unstable node with $\lambda_1 = 5$ and $\lambda_2 = 3$ cannot be transformed into an unstable node with $\lambda_1 = 43$ and $\lambda_2 = 17$.

We might want to introduce some looser type of equivalence instead, which will allow (for example) any two unstable nodes to be considered equivalent.

- Classification of **qualitatively equivalent** types of phase portraits for two-dimensional simple linear systems. Under this equivalence relation (**Definition 2.4.1**) there are only four different types:
 - Stable node, star node, improper node or spiral. (“Stable but not centre.”)
 - Centre.
 - Saddle.
 - Unstable node, star node, improper node or spiral. (“Unstable but not saddle.”)

(One subtle detail regarding Definition 2.4.1: They say that two systems are qualitatively equivalent if there is a continuous bijection f which maps the phase portrait of one system onto the phase portrait of the other and preserves the orientation of the trajectories. But how do we know that this relation is *symmetric*? (If system A is equivalent to system B , then we want system B to be equivalent to system A as well.) For this to hold, the inverse f^{-1} must also be a *continuous* bijection. This isn't necessarily true for continuous bijections in general; consider for example the function from the interval $(-\pi, \pi]$ in \mathbf{R} to the unit circle in \mathbf{R}^2 given by $f(t) = (\cos t, \sin t)$. But there is a famous and rather difficult theorem by Brouwer which says that for a continuous bijection between two open sets in \mathbf{R}^n the inverse is automatically continuous, so the definition is actually correct as it stands.)

- Qualitative equivalence is also called **topological equivalence**.

(A stronger condition is C^k -**equivalence**, where one requires the map f and its inverse f^{-1} to be of class C^k instead of just continuous.)

- As an aside, we may also mention the slightly different notion of **conjugacy**, where one also requires the *time parametrization* of the trajectories to be respected. More precisely, two systems with flows φ and ψ are said to be **topologically conjugate** if there's a continuous map f with continuous inverse f^{-1} such that

$$f \circ \varphi_t = \psi_t \circ f$$

for all t . That is, following the flow φ of one system for t units of time and then mapping over to the other phase portrait is always the same as first mapping to the other phase portrait and then following *that* flow ψ for t units of time.

(And the systems are C^k -**conjugate** if f and f^{-1} are of class C^k .)

For example, a system whose phase portrait consists of concentric circles all traversed with the same period T is *topologically equivalent* to a system whose phase portrait consists of concentric circles traversed with different periods, but the systems are *not topologically conjugate*.

- The **exponential function** for square matrices P :

$$e^P = \exp(P) = I + P + \frac{1}{2!} P^2 + \frac{1}{3!} P^3 + \dots$$

- The solution to $\dot{\mathbf{x}} = A\mathbf{x}$ is

$$\mathbf{x}(t) = e^{At} \mathbf{x}(0).$$

(Here it's important that A is a *constant* matrix, not time-dependent.)

- **Affine** systems $\dot{\mathbf{x}} = A\mathbf{x} + \mathbf{h}(t)$, also called **non-homogeneous linear** systems.

It's easy if \mathbf{h} is time-independent and $A\mathbf{x}_0 = \mathbf{h}$ for some \mathbf{x}_0 : just set $\mathbf{y}(t) = \mathbf{x}(t) + \mathbf{x}_0$ to get $\dot{\mathbf{y}} = A\mathbf{y}$. But even if this doesn't hold, an affine system can be solved using e^{-At} as an integrating factor.

- Linear systems and Jordan form in n dimensions.

Lesson 4

- Using the trace–determinant criterion: 2.13.
- Phase portrait via change of variables: 2.14.
- Computing matrix exponentials: 2.22, 2.23ab, 2.23cd.
- Affine systems: 2.29abcd, 2.30.
- A three-dimensional linear system: 2.33.
- Solution corresponding to a 4×4 Jordan block: 2.35a.

Lecture 5. Nonlinear systems, linearization at an equilibrium point

(Arrowsmith & Place, sections 3.1, 3.2, 3.3, 3.4.)

Now back to nonlinear systems $\dot{\mathbf{x}} = \mathbf{X}(\mathbf{x})$, in the plane \mathbf{R}^2 for simplicity, but the results are valid in \mathbf{R}^n too. We assume that the vector field $\mathbf{X}(\mathbf{x})$ is of class C^1 , so that we have existence and uniqueness of solutions; this assumption is also needed for the linearization theorem below to be valid.

- Suppose that $\mathbf{x}^* = (a_1, a_2)$ is an equilibrium point: $X_1(a_1, a_2) = X_2(a_1, a_2) = 0$. Since the functions X_1 and X_2 are assumed to be of class C^1 , they are also differentiable, which by definition means that

$$X_1(a_1 + h_1, a_2 + h_2) = \underbrace{X_1(a_1, a_2)}_{=0} + \frac{X_1}{\partial x_1}(a_1, a_2) h_1 + \frac{X_1}{\partial x_2}(a_1, a_2) h_2 + \text{remainder},$$

$$X_2(a_1 + h_1, a_2 + h_2) = \underbrace{X_2(a_1, a_2)}_{=0} + \frac{X_2}{\partial x_1}(a_1, a_2) h_1 + \frac{X_2}{\partial x_2}(a_1, a_2) h_2 + \text{remainder},$$

where the remainders tend to zero *faster** than $\sqrt{h_1^2 + h_2^2}$ as $(h_1, h_2) \rightarrow (0, 0)$.

- If we discard the remainders, we get a **linear** system for $\mathbf{h}(t) = \mathbf{x}(t) - \mathbf{x}^*$:

$$\dot{h}_1 = \frac{X_1}{\partial x_1}(a_1, a_2) h_1 + \frac{X_1}{\partial x_2}(a_1, a_2) h_2,$$

$$\dot{h}_2 = \frac{X_2}{\partial x_1}(a_1, a_2) h_1 + \frac{X_2}{\partial x_2}(a_1, a_2) h_2,$$

or in matrix notation,

$$\dot{\mathbf{h}} = \underbrace{\frac{\partial \mathbf{X}}{\partial \mathbf{x}}(\mathbf{x}^*)}_{=A} \mathbf{h}.$$

This system is called the **linearization** of the original system at the equilibrium point \mathbf{x}^* . Its matrix A is the **Jacobian matrix** of $\mathbf{X}(x)$, *evaluated* at the equilibrium point \mathbf{x}^* (so it's really just a *constant* matrix).

- Our hope is that the linear system $\dot{\mathbf{h}} = \mathbf{A}\mathbf{h}$ (which we know how to analyze completely) will tell us something about the behaviour of the original nonlinear system $\dot{\mathbf{x}} = \mathbf{X}(\mathbf{x})$ *near* the point \mathbf{x}^* .

And this is indeed the case, provided that \mathbf{x}^* is a **hyperbolic† equilibrium point**, meaning that the Jacobian matrix $A = \frac{\partial \mathbf{X}}{\partial \mathbf{x}}(\mathbf{x}^*)$ has **no eigenvalues on the imaginary axis** in the complex plane.


Under that condition, **Theorem 3.3.1** (the **linearization theorem**) says that the nonlinear system $\dot{\mathbf{x}} = \mathbf{X}(\mathbf{x})$ is indeed topologically equivalent‡ to its linearization $\dot{\mathbf{h}} = \mathbf{A}\mathbf{h}$ in a neighbourhood of \mathbf{x}^* .

In terms of the trace-determinant diagram of $A = \frac{\partial \mathbf{X}}{\partial \mathbf{x}}(\mathbf{x}^*)$:

*What this means is that the quotient $R(h, k) / \sqrt{h^2 + k^2}$ tends to zero, where $R(h, k)$ is the remainder.

†The word “hyperbolic” is very over-used in mathematics, and it is perhaps not a very good choice in this context, but unfortunately it has become standard terminology. Arrowsmith & Place avoid this word initially, and express the condition by saying that the linearized system should be **simple** ($\lambda = 0$ is not an eigenvalue) and **not a centre** (we don't have $\lambda = 0 \pm ik$ either). But they introduce it a little later, on p. 80.

‡ Actually, it's even topologically conjugate, but the book doesn't introduce that concept.



The linearization theorem is also called the **Hartman–Grobman theorem**, proved independently by Philip Hartman in the U.S.A. and D. M. Grobman in the Soviet Union around 1960. The proof is rather difficult, and way beyond the scope of this course.[§]

(However, a simpler theorem, not dealing with topological equivalence but only with determining *stability* based on linearization, can be proved using Liapunov’s theorems that we will learn about in the next lecture.)

Lesson 5

- Linearization: 3.5 (3.5eg), 3.6, 3.7 (3.7b).
- An isolated, but non-simple, fixed point: 3.8.
- Non-isolated fixed points: 3.11.
(To avoid confusion: here “a *line* of fixed points” rather means a *curve*.)
- An explicit example of a topological conjugacy: A4*.
- An example regarding smoothness in the Hartman–Grobman theorem: A5*.

Additional problems for lesson 5

A4 Find a topological conjugacy $(u, v, w) = h(x, y, z)$ between the nonlinear system

$$\dot{x} = -x, \quad \dot{y} = -y + xz, \quad \dot{z} = z$$

and the corresponding linearized system at the origin,

$$\dot{u} = -u, \quad \dot{v} = -v, \quad \dot{w} = w.$$

[Answer: For example, $h(x, y, z) = (x, y - xz, z)$ works (and it’s actually even a global diffeomorphism, not just a local homeomorphism).]

[§]For two-dimensional systems, a stronger result holds (also proved by Hartman): if the vector field $\mathbf{X}(\mathbf{x})$ is of class C^2 , then the nonlinear system is actually C^1 -conjugate (not just topologically conjugate) to its linearization, and moreover the derivative of the conjugating map at the origin is the identity. This means, for example, that if the linearization is a saddle, the eigenvectors will tell us the correct incoming and outgoing *directions* of the trajectories of the nonlinear system.

It is not true in three or more dimensions that we can always get C^1 -conjugacy. However, it *is* true (as was proved as recently as 2003) that the continuous conjugation map is differentiable *at the origin*, with the derivative there equal to the identity. (The proof assumes that $\mathbf{X}(\mathbf{x})$ is of class C^∞ , but the claim is conjectured to be true already for class C^2 .)

It’s also not true (even in two dimensions) that one can get C^k -conjugacy with $k \geq 2$ by assuming the vector field to be nicer; see problem A5 for a counterexample with a vector field of class C^∞ where one doesn’t get more than C^1 -conjugacy.

A5 Solve the system $\dot{x} = -x$, $\dot{y} = -2y + x^2$ exactly and draw the phase portrait. Eliminate t to get expressions $y = y(x)$ for the solution curves; what degree of smoothness do they have at the origin? Compare to the solution of the linearized system $\dot{x} = -x$, $\dot{y} = -2y$. Are the phase portraits C^2 -equivalent?

Lecture 6. Stability theorems

(Arrowsmith & Place, sections 3.5, 3.6, 3.7.)

- **Definitions 3.5.1–4:** The definition of a **stable** equilibrium (sometimes called **Liapunov stable**). Some stable equilibria are **asymptotically stable**. And those which are stable but not asymptotically stable are called **neutrally stable**. So there are exactly those two types of stable equilibria. **Unstable** equilibrium simply means an equilibrium which is not stable.
- Russian names can be transliterated into the Latin alphabet in many ways. Here I'm writing **Liapunov** like in the textbook, but a very common alternative in English is **Lyapunov**, and one may also come across **Ljapunov**, **Liapounoff**, and so on. Anyway, it's pronounced with the stress on the last syllable: "-OFF".
- **Theorem 3.5.1** is **Liapunov's stability theorem** (1892).

This theorem is useful for showing stability in situations where linearization is inconclusive. Even more importantly, it also provides a **domain of stability**; that is, a neighbourhood N of an asymptotically stable equilibrium \mathbf{x}^* , such that any solution which starts in N stays in N and converges to \mathbf{x}^* as $t \rightarrow \infty$. (From linearization we can only say that if all eigenvalues have negative real part, then there is *some* domain of stability, but we don't get any clue about the *size* of that domain.)

However, Arrowsmith & Place don't state exactly how to find a domain of stability, and their proof of the theorem is also rather unclear. I will try to give more precise statements here.

First let us fix some terminology.

Definition. Let I be a proper* interval in \mathbf{R} , and let f be a real-valued function whose domain of definition contains I . The function f is said to be **strictly decreasing** on I if

$$f(t_1) > f(t_2)$$

whenever $t_1 \in I$, $t_2 \in I$ and $t_1 < t_2$. It is **weakly decreasing** on I if

$$f(t_1) \geq f(t_2)$$

whenever $t_1 \in I$, $t_2 \in I$ and $t_1 < t_2$.

Remark. Any function which is strictly decreasing is weakly decreasing as well.

Remark. In English, it is more common to say **decreasing** and **non-increasing** instead of **strictly decreasing** and **weakly decreasing**, but I have chosen the latter option here to reduce the risk of confusion with the usual terminology in Swedish, which is **strängt/strikt avtagande** and **avtagande**, respectively. (And similarly in German and French.)

From now on we consider some fixed dynamical system $\dot{\mathbf{x}} = \mathbf{X}(\mathbf{x})$ in \mathbf{R}^n , where the vector field \mathbf{X} is defined in some open set $S \subseteq \mathbf{R}^n$, and we assume that $V: \Omega \rightarrow \mathbf{R}$ is a differentiable function defined on some open set $\Omega \subseteq S$. (And of course we assume that Ω and S are not the empty set, since that wouldn't be very interesting.)

* A **proper interval** is an interval (in \mathbf{R}) which contains infinitely many points, as opposed to the *degenerate* intervals $[a, a] = \{a\}$ and $\emptyset = \{\}$.

Definition. The function $\dot{V}: \Omega \rightarrow \mathbf{R}$ is the dot product of the gradient ∇V and the vector field \mathbf{X} :

$$\dot{V}(\mathbf{x}) = \nabla V(\mathbf{x}) \cdot \mathbf{X}(\mathbf{x}), \quad \mathbf{x} \in \Omega.$$

Theorem. Suppose $\dot{V}(\mathbf{x}) < 0$ for all $\mathbf{x} \in \Omega$. Then, for any solution $\mathbf{x}(t)$ of the system which stays in the set Ω during some nonempty open time interval I , the function

$$f(t) = V(\mathbf{x}(t)), \quad t \in I$$

is strictly decreasing on I . If instead $\dot{V}(\mathbf{x}) \leq 0$ for all $\mathbf{x} \in \Omega$, then f is weakly decreasing on I . And if $\dot{V}(\mathbf{x}) = 0$ for all $\mathbf{x} \in \Omega$, then f is constant on I .

Proof. The chain rule gives

$$\begin{aligned} f'(t) &= \nabla V(\mathbf{x}(t)) \cdot \dot{\mathbf{x}}(t) \\ &= \nabla V(\mathbf{x}(t)) \cdot \mathbf{X}(\mathbf{x}(t)) = \dot{V}(\mathbf{x}(t)), \quad t \in I. \end{aligned}$$

So if we know that $\dot{V} < 0$ everywhere in Ω , and that $\mathbf{x}(t)$ stays in Ω for $t \in I$, then we have $f'(t) < 0$ for $t \in I$, which implies that f is strictly decreasing on I (by a very basic calculus theorem). Similarly if $\dot{V} \leq 0$ or $\dot{V} = 0$. \square

Theorem (Liapunov's stability theorem, weak version). Let \mathbf{x}^* be an equilibrium point of the dynamical system $\dot{\mathbf{x}} = \mathbf{X}(\mathbf{x})$, $\mathbf{x} \in S \subseteq \mathbf{R}^n$. Suppose that there is a **weak Liapunov function**, i.e., a differentiable function $V: \Omega \rightarrow \mathbf{R}$ defined on some open set $\Omega \subseteq S$ containing \mathbf{x}^* and satisfying the conditions

1. $V(\mathbf{x}^*) = 0$ and $V(\mathbf{x}) > 0$ for all $\mathbf{x} \in \Omega \setminus \{\mathbf{x}^*\}$,
2. $\dot{V}(\mathbf{x}) \leq 0$ for all $\mathbf{x} \in \Omega$.

Then the equilibrium \mathbf{x}^* is **stable**.

Remark. As the theorem is formulated, the function $V: \Omega \rightarrow \mathbf{R}$ has *exactly* the set Ω as its domain of definition. Usually we have some nice function V (typically a polynomial) which is defined *everywhere* to begin with. What we actually do then is that we compute \dot{V} , use that to locate some open set Ω where the assumptions of the theorem are fulfilled, and then apply the theorem with V equal to the *restriction* of the original function V to the set Ω .

Outline of proof. If U is any neighbourhood of \mathbf{x}^* , let $B \subset U \cap \Omega$ be a closed ball centered at \mathbf{x}^* and set

$$U' = \{\mathbf{x} \in B : V(\mathbf{x}) < \alpha\},$$

where $\alpha > 0$ is the minimum of V on the boundary sphere ∂B . Then $U' \subset B \subset U$, and U' is a neighbourhood of \mathbf{x}^* such that trajectories starting in U' can't leave U (in fact, they can't even leave U'). \square

Detailed proof. Let U be an arbitrary neighbourhood of \mathbf{x}^* . To prove stability, we need to find another neighbourhood U' such that solutions starting in U' will never leave U . To find U' we begin by taking a closed ball

$$B = \overline{B(\mathbf{x}^*, \varepsilon)} = \{\mathbf{x} \in \mathbf{R}^n : |\mathbf{x} - \mathbf{x}^*| \leq \varepsilon\}$$

centered at \mathbf{x}^* , with radius $\varepsilon > 0$ small enough for B to be contained inside both U and Ω . (This is possible since U and Ω are neighbourhoods of \mathbf{x}^* .) The boundary

$$\partial B = \{\mathbf{x} \in \mathbf{R}^n : |\mathbf{x} - \mathbf{x}^*| = \varepsilon\}$$

is a sphere of radius ε centered at \mathbf{x}^* . This sphere is a compact set (closed and bounded), and V is continuous by assumption, so according to the extreme value theorem V has a smallest value on ∂B :

$$\alpha = \min_{\mathbf{x} \in \partial B} V(\mathbf{x}).$$

In other words, there is a point $\mathbf{x}_0 \in \partial B$ such that

$$\alpha = V(\mathbf{x}_0) \leq V(\mathbf{x}) \quad \text{for all } \mathbf{x} \in \partial B.$$

Since V is positive definite on Ω and we have chosen B small enough to be a subset of Ω , we have $V(\mathbf{x}) > 0$ for all $\mathbf{x} \in \partial B$, in particular

$$\alpha = V(\mathbf{x}_0) > 0.$$

Now set

$$U' = \{\mathbf{x} \in B : V(\mathbf{x}) < \alpha\}.$$

Then U' contains \mathbf{x}^* , since $V(\mathbf{x}^*) = 0 < \alpha$. And U' is an open set, since V is continuous.[†] In other words, U' is an open neighbourhood of \mathbf{x}^* . Moreover, a trajectory $\mathbf{x}(t)$ starting in U' (at $t = 0$, say) can't leave U . Here's why: to leave U , the trajectory would have to leave B to begin with (since $U' \subset B \subset U$), and it's a continuous curve so it would have to intersect the boundary sphere ∂B in order to get out. But $f(t) = V(\mathbf{x}(t))$ is a weakly decreasing function of t as long as $\mathbf{x}(t)$ stays in B (since $B \subset \Omega$, and $\dot{V} \leq 0$ in Ω by assumption). Since we start in U' , we have $f'(0) < \alpha$, and hence $f(t) < \alpha$ for $t \geq 0$. So it's impossible for the trajectory to reach ∂B , since that would mean $f(t) \geq \alpha$ for some $t > 0$. (In fact, the trajectory can't even leave U' – as soon as it did, it would mean that $f(t) \geq \alpha$.)

(One more technical detail: since the trajectory stays inside the compact set B , it must exist for all $t \geq 0$; there can't be any "blowup in finite time". We haven't proved that theorem in this course, so we'll just accept this fact on faith here.) \square

Theorem (Liapunov's stability theorem, strong version). Let \mathbf{x}^* be an equilibrium point of the dynamical system $\dot{\mathbf{x}} = \mathbf{X}(\mathbf{x})$, $\mathbf{x} \in S \subseteq \mathbf{R}^n$. Suppose that there is a **strong Liapunov function**, i.e., a differentiable function $V: \Omega \rightarrow \mathbf{R}$ defined on some open set $\Omega \subseteq S$ containing \mathbf{x}^* and satisfying the conditions

1. $V(\mathbf{x}^*) = 0$ and $V(\mathbf{x}) > 0$ for all $\mathbf{x} \in \Omega \setminus \{\mathbf{x}^*\}$,
2. $\dot{V}(\mathbf{x}) < 0$ for all $\mathbf{x} \in \Omega \setminus \{\mathbf{x}^*\}$.

Then the equilibrium \mathbf{x}^* is **asymptotically stable**. In fact, for any closed ball $B = \overline{B(\mathbf{x}^*, r)}$ contained in Ω , the set

$$N = \{\mathbf{x} \in B : V(\mathbf{x}) < \alpha\}, \quad \text{where } \alpha = \min_{\mathbf{x} \in \partial B} V(\mathbf{x}),$$

is a **domain of stability**: solutions starting in N stay in N , and converge to \mathbf{x}^* as $t \rightarrow \infty$.

Remark. If $\Omega = \mathbf{R}^n$ and if the additional condition

$$V(\mathbf{x}) \rightarrow \infty \quad \text{as } |\mathbf{x}| \rightarrow \infty$$

holds, then for *any* given point $\mathbf{x}_0 \in \mathbf{R}^n$ it is true that

$$\min_{\mathbf{x} \in \partial B} V(\mathbf{x}) > V(\mathbf{x}_0)$$

if we take B large enough. This means that $\mathbf{x}_0 \in N$ for that choice of B , causing the trajectory starting at \mathbf{x}_0 to converge to \mathbf{x}^* . So in this case \mathbf{x}^* is stable and *every* trajectory of the system converges to \mathbf{x}^* , which is expressed by saying that \mathbf{x}^* is **globally asymptotically stable**.

[†] Take any $\mathbf{x} \in U'$, or in other words any $\mathbf{x} \in B$ with $V(\mathbf{x}) < \alpha$. Then actually \mathbf{x} is in the interior of B , since $V \geq \alpha$ on the boundary ∂B . Continuity of V at \mathbf{x} means that there is an open ball $B_2 = B(\mathbf{x}, \delta)$ where $V < \alpha$, and this ball must also be contained in the interior of B , for the same reason. So $B_2 \subseteq U'$. Thus any $\mathbf{x} \in U'$ has an open neighbourhood contained in U' , and this is exactly what it means for U' to be open.

Remark. If one studies the proof, it should be clear that the set B in the definition of N doesn't really have to be precisely a closed ball, just something which is topologically equivalent to a ball. For example, if our Liapunov function is $V(x, y) = x^6 + y^4$, then for $k > 0$ the sublevel set $B = \{x^6 + y^4 \leq k\}$ is sufficiently "ball-like" for the proof to work: no continuous curve can pass from the interior to the exterior without crossing the closed level curve $\partial B = \{x^6 + y^4 = k\}$. The same arguments as in the proof then show that trajectories can't leave the set $N = \{\mathbf{x} \in B : V(x, y) < k\} = \{x^6 + y^4 < k\}$, so provided that B is contained in the region Ω , N is a domain of stability for the equilibrium $(0, 0)$.

Outline of proof. Stability follows from the weak version of Liapunov's theorem. Any trajectory starting in N must stay in N , and along such a trajectory the function V decreases strictly towards some limit $L \geq 0$. But $L > 0$ would contradict the continuity of the flow φ_t , so $L = 0$, which in turn implies that the trajectory converges to \mathbf{x}^* (the only point where $V = 0$). \square

Detailed proof. Stability follows from the weak version of Liapunov's theorem. As in the proof of that theorem, we see that N (as defined above) is an open neighbourhood of \mathbf{x}^* , and that any trajectory $\mathbf{x}(t)$ starting in N stays in N and is defined for all $t \geq 0$. For the constant solution $\mathbf{x}(t) = \mathbf{x}^*$ there is nothing to prove – of course it converges to \mathbf{x}^* ! So suppose $\mathbf{x}(t)$ is some *other* solution starting in N . Then, by the assumption $\dot{V} < 0$, $V(\mathbf{x}(t))$ is strictly decreasing function of t on the interval $t \geq 0$, and it's bounded below (since $V \geq 0$), so it has a limit $L \geq 0$ as $t \rightarrow \infty$.

We want to show that $L = 0$, so assume $L > 0$ in order to get a contradiction. Take any sequence of positive numbers $t_n \nearrow \infty$; then $\mathbf{x}_n = \mathbf{x}(t_n)$ is a sequence of points in the compact set B . According to the Bolzano–Weierstrass theorem, a standard theorem about compact sets in \mathbf{R}^n , this sequence of points must have a convergent subsequence, i.e., there is a point $\mathbf{y} \in B$ and an integer sequence $n_k \nearrow \infty$ such that $\mathbf{x}_{n_k} \rightarrow \mathbf{y}$ as $k \rightarrow \infty$. Since V is continuous, we can move the limit outside V and obtain

$$V(\mathbf{y}) = V\left(\lim_{k \rightarrow \infty} \mathbf{x}_{n_k}\right) = \lim_{k \rightarrow \infty} V(\mathbf{x}_{n_k}) = L.$$

We are assuming $L > 0$, which means that $\mathbf{y} \neq \mathbf{x}^*$ (since V is positive definite), and V will thus continue to decrease *strictly* along the trajectory starting at \mathbf{y} . So the flow φ_1 , for example (or φ_t for any fixed $t > 0$), will map \mathbf{y} to a point where $V < L$. But φ_1 is a continuous function, so it will also map all sufficiently nearby points \mathbf{x}_{n_k} to points where $V < L$:

$$V(\varphi_1(\mathbf{x}_{n_k})) = V(\mathbf{x}(1 + t_{n_k})) < L, \quad \text{for all sufficiently large } k.$$

But we know that $V(\mathbf{x}(t)) > L$ for all $t \geq 0$, since $V(\mathbf{x}(t))$ is *decreasing* towards the limit L . This contradiction shows that the assumption $L > 0$ must have been incorrect. Hence $L = 0$.

Now we know that $V(\mathbf{x}(t)) \searrow L = 0$ as $t \rightarrow \infty$. It remains to show that this implies $\mathbf{x}(t) \rightarrow \mathbf{x}^*$, i.e., that for any $\varepsilon > 0$ there is a time τ such that $\mathbf{x}(t) \in B_\varepsilon$ for all $t > \tau$, where $B_\varepsilon = B(\mathbf{x}^*, \varepsilon)$ is the open ball of radius ε centered at \mathbf{x}^* . We may assume that $0 < \varepsilon < r$, where r is the radius of the closed ball B . Then

$$B \setminus B_\varepsilon = \overline{B(\mathbf{x}^*, r)} \setminus B(\mathbf{x}^*, \varepsilon)$$

is a compact nonempty set, so the continuous function V has a smallest value β on this set (and $\beta > 0$ since V is positive definite). What this means is that if $\mathbf{x} \in B$ and $V(\mathbf{x}) < \beta$, then $\mathbf{x} \in B_\varepsilon$. But we have $\mathbf{x}(t) \in B$ for all $t \geq 0$, and since $V(\mathbf{x}(t)) \searrow 0$ as $t \rightarrow \infty$ there is a τ such that $V(\mathbf{x}(t)) < \beta$ for $t > \tau$. Consequently $\mathbf{x}(t) \in B_\varepsilon$ for $t > \tau$, as desired. \square

- The very useful **Theorem 3.5.2** is an improvement of Liapunov's theorem which is due to LaSalle (1960). It allows us to conclude *asymptotic* stability using only a *weak* Liapunov function, provided an additional condition is satisfied. The proof is not given in the book, but it is a consequence of something called **LaSalle's invariance principle** (see the [next lecture](#)).

Theorem (LaSalle’s stability theorem). Let \mathbf{x}^* be an equilibrium point of the dynamical system $\dot{\mathbf{x}} = \mathbf{X}(\mathbf{x})$, $\mathbf{x} \in S \subseteq \mathbf{R}^n$. Suppose that there is a **weak Liapunov function** $V : \Omega \rightarrow \mathbf{R}$ on some open set $\Omega \subseteq S$ containing \mathbf{x}^* , and in addition suppose that the set

$$\{\mathbf{x} \in \Omega : \dot{V}(\mathbf{x}) = 0\}$$

contains no complete trajectory except the constant solution \mathbf{x}^* .

Then the equilibrium \mathbf{x}^* is **asymptotically stable**, and N (defined as in the strong version of Liapunov’s theorem) is a **domain of stability**.

(And if $\Omega = \mathbf{R}^n$ and $V(\mathbf{x}) \rightarrow \infty$ as $|\mathbf{x}| \rightarrow \infty$, then \mathbf{x}^* is **globally asymptotically stable**.)

Proof. See the [next lecture](#). □

- Here is a somewhat subtle point concerning the above theorems and weak Liapunov functions. If we have a function V which is everywhere defined and nice (continuously differentiable), then the set $\Omega_1 = \{\mathbf{x} \in \mathbf{R}^n : \dot{V}(\mathbf{x}) \leq 0\}$ will be *closed*. But the theorems, as they are formulated above, require us to restrict V to an *open* set Ω . So we have to shrink Ω_1 “by hand” to get an *open* set Ω where $\dot{V} \leq 0$ holds. The purpose of this, as well as all the business with closed balls contained inside Ω , is to avoid accidentally making plausible-sounding claims which are actually false. We know that V is weakly decreasing along trajectories, but only as long as they stay in Ω , so we need to take some precautions to prevent the trajectories from sneaking out of Ω !

Example. If the system is $\dot{x} = y$, $\dot{y} = -x - y(1 - x^2)$, and if $V(x, y) = x^2 + y^2$, then $\dot{V} = -2y^2(1 - x^2)$. Thus, the set $\Omega_1 = \{\dot{V} \leq 0\}$ is the union of the closed strip $-1 \leq x \leq 1$ and the line $y = 0$. So any trajectory will be moving closer to the origin (or at least not further away from it) as long as it is inside the strip, but trajectories for $y > 3$ (or so) will enter the strip from the left, *leave it again on the right* (a little further down), and then go off steeply upwards towards infinity instead of converging towards the equilibrium $(0, 0)$. So for example, a set like $\{(x, y) \in \Omega_1 : V(x, y) \leq 100\}$ is not forward invariant despite V being weakly decreasing on trajectories in Ω_1 ! But we can take Ω to be the open strip $-1 < x < 1$, let B be any closed ball $x^2 + y^2 \leq k$ with $0 < k < 1$ so that it fits inside Ω , and then the set N given by $x^2 + y^2 < k$ will be a domain of attraction by LaSalle’s theorem, since there are no trajectories contained in the line $y = 0$ except the equilibrium solution $(x(t), y(t)) = (0, 0)$. And since this is true for any $0 < k < 1$, in fact the open unit disk $x^2 + y^2 < 1$ is a domain of attraction.[‡]

- **Theorem 3.5.3 is Liapunov’s instability theorem.**

- We have relied upon the rather deep Hartman–Grobman theorem to show that an equilibrium is asymptotically stable if the linearization there is asymptotically stable. This fact can be proved more directly using Liapunov’s stability theorem. The simplest case is when the Jacobian matrix $A = \frac{\partial \mathbf{X}}{\partial \mathbf{x}}(\mathbf{x}^*)$ has distinct real eigenvalues (assumed negative, in order for $d\mathbf{h}/dt = A\mathbf{h}$ to be asymptotically stable):

$$0 > \lambda_1 > \dots > \lambda_n.$$

Make the usual linear change of coordinates $\mathbf{x} = M\mathbf{y}$, where the columns of M form a basis of eigenvectors of A . In terms of these coordinates, we have a system $\dot{\mathbf{y}} = \mathbf{Y}(\mathbf{y})$ with an equilibrium \mathbf{y}^* where the Jacobian is diagonal:

$$J = \frac{\partial \mathbf{Y}}{\partial \mathbf{y}}(\mathbf{y}^*) = \text{diag}(\lambda_1, \dots, \lambda_n).$$

[‡]We can actually do yet a little better: the closed unit disk $x^2 + y^2 \leq 1$ is forward invariant since the open unit disk is; this follows from the continuity of the flow, since if φ_t (for some $t > 0$) would map some point on the unit circle to a point outside the circle, then by continuity it would also have to map some nearby point inside the circle out of the circle, which we know it doesn’t. So the closed unit disk is compact and forward invariant, and then we can try applying LaSalle’s invariance principle (see next lecture) to it. This turns out to be successful (exercise), so actually the *closed* unit disk is a domain of attraction.

So $\mathbf{k} = \mathbf{y} - \mathbf{y}^*$ satisfies

$$\dot{\mathbf{k}} = J\mathbf{k} + \text{remainder},$$

where the remainder tends to zero faster than $|\mathbf{k}|$. It's not too difficult to check that $V(\mathbf{k}) = \sum k_i^2$ is a strict Liapunov function for this system in a neighbourhood of $\mathbf{k} = \mathbf{0}$, which proves asymptotic stability. With repeated and/or non-real eigenvalues things are a bit more complicated, but if all eigenvalues have negative real part one can find a strong Liapunov function in the form of a sum of squares in those cases too.

Similarly, one can use Liapunov's instability theorem to prove that if some eigenvalue has positive real part, then the equilibrium is unstable.

- The **flow box theorem**. Global phase portraits.
- **First integrals**, also known as **constants of motion**, **integrals of motion**, **conserved quantities**, **invariants**, etc.

(Can sometimes be found by writing $dy/dx = \dot{y}/\dot{x} = Y(x, y)/X(x, y)$ and solving the resulting ODE for $y = y(x)$.)

- If $H(p, q)$ is any C^1 function, then the **Hamiltonian system**

$$\begin{pmatrix} \dot{p} \\ \dot{q} \end{pmatrix} = \begin{pmatrix} \partial H / \partial q \\ -\partial H / \partial p \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \nabla H$$

automatically has H as a first integral. A fundamental fact in mechanics is that the Hamiltonian system generated by $H(p, q) = \frac{1}{2}p^2 + V(q)$ is equivalent to the Newton-type equation $\ddot{q} = -V'(q)$. (This works also in higher dimensions, with vectors \mathbf{p} and \mathbf{q} instead.)

Lesson 6

- Using strong Liapunov functions: 3.13abe, 3.14abe.
- Using weak Liapunov functions: 3.15, 3.17bc.
- Finding a domain of stability using a less obvious Liapunov function: [3.19b](#), 3.18*.
- An example showing that the assumption $V(\mathbf{x}) \rightarrow \infty$ is important in order to get global asymptotic stability: [A6](#).
- Liapunov's instability theorem: [3.22](#).
- Illustration of an explicit transformation which straightens out a nonlinear vector field: 3.24.
- First integrals (constants of motion): 3.28ab, 3.29*.

In 3.28b, you can do better than the answer in the book, and find a constant of motion which is actually defined for *all* x_1 and x_2 !

In 3.29, you should do much better than the answer in the book, which is actually wrong. A correct constant of motion is $F(x_1, x_2) = x_1(x_1^2 - x_2)/(x_1^2 + x_2)^2$. The level curves of this function are not easy to plot by hand, but you can of course do it on the computer if you want to see what the phase portrait looks like.

- An example^S illustrating why the requirement about stability in the definition of asymptotic stability is necessary: [A7](#).

^SThe system (3.33) in the book is another such example, but that one is much more difficult to analyze rigorously (Exercise 3.12); for details see Section 40 of the book *Stability of Motion* by W. Hahn (Springer, 1967).

Additional problems for lesson 6

A6 Show that

$$V(x, y) = \frac{x^2}{1+x^2} + y^2$$

is a global strong Liapunov function for the system

$$\dot{x} = -x(1-3x^2y^2), \quad \dot{y} = -y(1+x^2y^2),$$

but that the origin is **not** globally asymptotically stable (although it is locally asymptotically stable).

Note that $V(x, y)$ does *not* satisfy the condition that $V(x, y) \rightarrow \infty$ as $\sqrt{x^2 + y^2} \rightarrow \infty$. What do the level sets of V look like, in particular the level set $V = 1$?

[Hints: One can write \dot{V} as

$$\dot{V} = -2 \frac{x^2(1-x^2y^2)^2 + y^2(1+2x^2)(1+x^2y^2)}{(1+x^2)^2},$$

and $(x(t), y(t)) = (e^{2t}, e^{-2t})$ is a particular solution of the system.]

A7 Consider the system

$$\dot{x} = x - y - xr^2 + \frac{xy}{r}, \quad \dot{y} = x + y - yr^2 - \frac{x^2}{r},$$

where $r = \sqrt{x^2 + y^2}$, and where the right-hand sides are interpreted as zero when $(x, y) = (0, 0)$.[†] Rewrite this system in polar coordinates, and use this to draw the phase portrait.

Deduce that, with the exception of the equilibrium solution $(x, y) = (0, 0)$, all solutions $(x(t), y(t))$ approach the point $(x, y) = (1, 0)$ as $t \rightarrow \infty$, but $(1, 0)$ is still an **unstable** equilibrium.

[Answer: $\dot{r} = r - r^3$, $\dot{\theta} = 1 - \cos\theta$.]

Lecture 7. Limit sets

(Arrowsmith & Place, sections 3.8, 3.9.)

- Definition of the α -**limit set** $L_\alpha(\mathbf{x})$ and the ω -**limit set** $L_\omega(\mathbf{x})$ of a point \mathbf{x} . (Definition 3.8.1 and the paragraph just below it.)
- In three or more dimensions, limit sets can be extremely complicated, since trajectories have room to wind around in space in very strange ways. But in the plane, the possibilities are much more restricted, as shown by **Theorem 3.9.1**, the **Poincaré–Bendixson theorem** (given without proof in the textbook): if a compact nonempty limit set in the plane contains no equilibrium points, then it must be a periodic orbit.
- Some properties of ω -limit sets:
 - $L_\omega(\mathbf{x})$ is always a **closed** and **invariant** set. (See **Definition 3.9.2**.)
 - Limit sets may be empty, unbounded, disconnected (see problem A8). But if the forward orbit of \mathbf{x} is **bounded**, then $L_\omega(\mathbf{x})$ is **connected, compact** and **non-empty**, and*

$$\varphi_t(\mathbf{x}) \rightarrow L_\omega(\mathbf{x}) \quad \text{as } t \rightarrow \infty.$$

[†]This system is a bit nasty at the origin, since the right-hand sides are not differentiable there, only continuous. If you prefer, you can get a system of class C^1 by multiplying both right-hand sides by r^2 ; this rescaling of the vector field doesn't change what the phase portrait looks like.

*The notation $\varphi_t(\mathbf{x}) \rightarrow L_\omega(\mathbf{x})$ means that for any neighbourhood U of the limit set $L_\omega(\mathbf{x})$ there is a time T such that $\varphi_t(\mathbf{x}) \in U$ for all $t > T$.

The corresponding properties hold of course for α -limit sets (as $t \rightarrow -\infty$).

Proof of the invariance property. Suppose $\mathbf{y} \in L_\omega(\mathbf{x})$. By definition, this means that $\varphi_{t_n}(\mathbf{x}) \rightarrow \mathbf{y}$ for some sequence $t_n \rightarrow \infty$. Fix an arbitrary $t \in \mathbf{R}$. Applying the continuous function φ_t to both sides gives $\varphi_{t+t_n}(\mathbf{x}) \rightarrow \varphi_t(\mathbf{y})$, so $\varphi_t(\mathbf{y}) \in L_\omega(\mathbf{x})$. \square

(I omit the proofs of the other properties, although they are not very difficult.)

- A **limit cycle (Definition 3.8.2)** is a periodic orbit which lies in the α - or ω -limit set of some point not on the orbit.

To show the existence of a limit cycle for a planar system using the Poincaré–Bendixson theorem, one tries to find a **trapping region containing no equilibria**. A **trapping region** for a system with flow φ_t is a compact, connected set $D \subset \mathbf{R}^2$ such that $\varphi_t(D) \subset D$ for $t > 0$. When reading this definition, it's important to note that Arrowsmith & Place use the symbol “ \subset ” for *strict* set inclusion. The point is that if we only require D to be forward invariant (that is, $\varphi_t(D) \subseteq D$ for $t > 0$, with non-strict set inclusion “ \subseteq ”), then it may be the case that D is a union of periodic orbits[†], in which case there are no limit cycles in D . But with strict set inclusion, the points in the nonempty set $D \setminus \varphi_t(D)$ (for any fixed $t > 0$) can't lie on a periodic orbit[‡], and the ω -limit set of such a point \mathbf{x}_0 is a nonempty compact subset of D . If we have chosen the trapping region D such that it contains no equilibria, the Poincaré–Bendixson theorem says that $L_\omega(\mathbf{x}_0)$ must be a periodic orbit. Since \mathbf{x}_0 was not on any periodic orbit, $L_\omega(\mathbf{x}_0)$ doesn't contain \mathbf{x}_0 , so it is a periodic orbit which is the ω -limit set of a point not on the orbit – in other words, it's an ω -limit cycle. Conclusion: the trapping region D contains *at least one* limit cycle.

(But there may be many limit cycles in D ! To prove that there is *at most one* limit cycle is usually much more difficult.)

- Now that we know what an ω -limit set $L_\omega(\mathbf{x})$ is, we can state **LaSalle's invariance principle** that was mentioned in the previous lecture. We consider a system $\dot{\mathbf{x}} = \mathbf{X}(\mathbf{x})$ with flow φ_t .

Theorem. Suppose $V : \Omega \rightarrow \mathbf{R}$ is differentiable on the open set $\Omega \subseteq \mathbf{R}^n$, and satisfies $\dot{V}(\mathbf{y}) \leq 0$ for each \mathbf{y} in some closed set $M \subseteq \Omega$.

1. If \mathbf{x} is a point in M whose forward orbit $\mathcal{O}^+(\mathbf{x})$ never leaves M , then there is an $\alpha \in \mathbf{R}$ such that

$$L_\omega(\mathbf{x}) \subseteq \{\mathbf{y} \in M : V(\mathbf{y}) = \alpha\}.$$

This implies that $L_\omega(\mathbf{x})$ is an invariant set contained in the set

$$C = \{\mathbf{y} \in M : \dot{V}(\mathbf{y}) = 0\},$$

and hence $L_\omega(\mathbf{x}) \subseteq E$, where E is the *largest invariant subset* of C (i.e., E is the union of all trajectories which stay in C for all $t \in \mathbf{R}$).

Proof. This is trivially true if $L_\omega(\mathbf{x}) = \emptyset$, since the empty set is a subset of *every* set. So assume $L_\omega(\mathbf{x}) \neq \emptyset$. (In particular, this assumption entails that the solution $\varphi_t(\mathbf{x})$ exists for all $t \geq 0$.) To begin with,

$$\begin{aligned} L_\omega(\mathbf{x}) &\subseteq \overline{\mathcal{O}^+(\mathbf{x})} && \text{(follows from def. of } L_\omega(\mathbf{x})\text{)} \\ &\subseteq \overline{M} && \text{(since } \mathcal{O}^+(\mathbf{x}) \subseteq M \text{ by assumption)} \\ &= M && \text{(since } M \text{ is closed).} \end{aligned}$$

[†]For example (in polar coordinates), if the system is $\dot{r} = 0, \dot{\theta} = 1$, and D is the annulus $1 \leq r \leq 2$.

[‡]Suppose that $\mathbf{x}_0 \in D \setminus \varphi_t(D)$ for some $t > 0$ and that \mathbf{x}_0 lies on a periodic orbit with period $T > 0$. Let n be a positive integer such that $nT > t$. Then

$$\mathbf{y} = \varphi_{nT-t}(\mathbf{x}_0) \in \varphi_{nT-t}(D) \subset D$$

so that

$$\mathbf{x}_0 = \varphi_{nT}(\mathbf{x}_0) = \varphi_t(\mathbf{y}) \in \varphi_t(D),$$

which is a contradiction.

Next, the assumptions that $\dot{V} \leq 0$ on M and that $\varphi_t(\mathbf{x})$ stays in M for $t \geq 0$ imply that $V(\varphi_t(\mathbf{x}))$ is a weakly decreasing function of t for $t \geq 0$, so the limit

$$\alpha = \lim_{t \rightarrow \infty} V(\varphi_t(\mathbf{x}))$$

exists, either as a real number $\alpha \in \mathbf{R}$ or in the improper sense $\alpha = -\infty$. But if \mathbf{y} is any element in the nonempty set $L_\omega(\mathbf{x})$, meaning that $\varphi_{t_n}(\mathbf{x}) \rightarrow \mathbf{y}$ for some sequence $t_n \nearrow \infty$, then

$$V(\mathbf{y}) = V\left(\lim_{n \rightarrow \infty} \varphi_{t_n}(\mathbf{x})\right) = \lim_{n \rightarrow \infty} V(\varphi_{t_n}(\mathbf{x})) = \alpha,$$

since V is continuous. This shows that α equals the real number $V(\mathbf{y})$, not $-\infty$.

The above calculation holds for an arbitrary $\mathbf{y} \in L_\omega(\mathbf{x})$, so $V = \alpha$ on all of $L_\omega(\mathbf{x})$. And $L_\omega(\mathbf{x})$ is an invariant set (general property of limit sets), so $\varphi_t(\mathbf{y}) \in L_\omega(\mathbf{x})$ for all t if $\mathbf{y} \in L_\omega(\mathbf{x})$. Thus $V(\varphi_t(\mathbf{y})) = \alpha$ for all t , and hence

$$\dot{V}(\mathbf{y}) = 0 \quad \text{if } \mathbf{y} \in L_\omega(\mathbf{x}).$$

What we have shown now is that $L_\omega(\mathbf{x})$ is an invariant set which is contained in the set $C \subseteq M$ where $\dot{V} = 0$. Therefore, it must trivially be contained in E , the *largest* invariant set contained in C . □

2. If moreover the forward orbit $\mathcal{O}^+(\mathbf{x})$ is **bounded**, then $L_\omega(\mathbf{x})$ is nonempty and $\varphi_t(\mathbf{x}) \rightarrow L_\omega(\mathbf{x})$ as $t \rightarrow \infty$. So $\varphi_t(\mathbf{x}) \rightarrow E$ as $t \rightarrow \infty$.

Proof. The first sentence was one of the general properties of limit sets stated at the beginning of the lecture. The second sentence follows at once from the property $L_\omega(\mathbf{x}) \subseteq E$ that we proved in item 1. (The conclusion that $\varphi_t(\mathbf{x}) \rightarrow E$ is of course a bit weaker than $\varphi_t(\mathbf{x}) \rightarrow L_\omega(\mathbf{x})$, but the point is that the set E does not depend on \mathbf{x} .) □

3. If M is **compact** and **forward invariant**, then items 1 and 2 apply to *every* point $\mathbf{x} \in M$. So in this case, $\varphi_t(\mathbf{x}) \rightarrow E$ as $t \rightarrow \infty$, for every $\mathbf{x} \in M$.

Proof. Trivial. □

The point of this theorem is that the set E is often quite easy to determine. We find the set C simply by computing \dot{V} and checking where it's zero. This is typically some curve, if we are in \mathbf{R}^2 . Then we study what the vector field \mathbf{X} is doing at each point of the set C – if the vector field is pointing out from C at some point, then that point can't be part of a trajectory completely contained in C , so it can't belong to E .

A typical application is the situation described in **Theorem 3.5.2** (see the [previous lecture](#)), where we have only managed to find a *weak* Liapunov function V , but the “bad” set C where we have $\dot{V} = 0$ instead of $\dot{V} < 0$ doesn't contain any trajectories except the equilibrium point \mathbf{x}^* .

Proof of Theorem 3.5.2. As usual, let $B = \overline{B(\mathbf{x}^*, r)}$ be a closed ball (or some other neighbourhood of \mathbf{x}^* topologically equivalent to a closed ball) contained in Ω , and define

$$N = \{\mathbf{x} \in B : V(\mathbf{x}) < \alpha\}, \quad \text{where } \alpha = \min_{\mathbf{x} \in \partial B} V(\mathbf{x}) > 0.$$

Stability of \mathbf{x}^* follows from the weak version of Liapunov's theorem, so we just need to show that N is a domain of stability. To apply LaSalle's invariance principle, we need a compact and forward invariant set M , so N (which is open) won't do. Instead, take β with $0 \leq \beta < \alpha$, and let

$$M = \{\mathbf{x} \in B : V(\mathbf{x}) \leq \beta\}.$$

Then M is closed (and hence compact) since V is continuous.[§] And it's forward invariant; indeed, a trajectory starting in M can't leave M , since then it would enter the part of B where $V > \beta$, so V wouldn't be weakly decreasing along that trajectory, and that would contradict the assumption that $\dot{V} \leq 0$ in Ω . Now the invariance principle says that every trajectory starting in M converges to E , the largest invariant subset of $C = \{\mathbf{x} \in M : \dot{V}(\mathbf{x}) = 0\}$. But by assumption, there are no trajectories even in the larger set $C_2 = \{\mathbf{x} \in \Omega : \dot{V}(\mathbf{x}) = 0\}$ except for the equilibrium \mathbf{x}^* . Hence $E = \{\mathbf{x}^*\}$, and every trajectory starting in M converges to \mathbf{x}^* . So every such set M is a domain of stability.

To show that N is a domain of stability, just note that any $\mathbf{x}_0 \in N$ belongs to the set $M \subseteq N$ defined using $\beta = V(\mathbf{x}_0) < \alpha$. Therefore the trajectory starting at \mathbf{x}_0 stays in N and converges to \mathbf{x}^* . \square

- The **Poincaré map** (or **first-return map**) associated with a closed orbit.
- **Theorem 3.9.2** is called the **Bendixson criterion**.

A simple generalization is the **Bendixson–Dulac criterion**, which gives the same conclusion provided that there is a function $f(x_1, x_2)$ of class C^1 such that the divergence of the rescaled vector field $f\mathbf{X}$,

$$\nabla \cdot (f\mathbf{X}) = \frac{\partial}{\partial x_1}(fX_1) + \frac{\partial}{\partial x_2}(fX_2),$$

is of constant sign in D .

Lesson 7

- α - and ω -limit sets: 3.35, [A8*](#).
(There's an error in the answer to 3.35b; it should say $L_\alpha(\mathbf{x}) = \{\mathbf{0}\}$ for $0 < r < 1$.)
- Poincaré map: 3.36.
(In this problem, it's understood that $a > 0$.)
- Invariant sets, trapping regions: 3.42 ([3.42e](#)), 3.43.
- The Bendixson criterion: [A9](#).

Additional problems for lesson 7

- A8 (a) Show that for the equation $\dot{x} = 1$, the ω -limit set of each point is the **empty** set.
(b) Sketch the phase portrait for the system

$$\dot{x} = -y + \frac{x}{1+x^2}, \quad \dot{y} = x(1-y^2).$$

Show that the ω -limit set of any point $(x, y) \neq (0, 0)$ in the strip $|y| < 1$ is the union of the lines $y = \pm 1$, and hence is **unbounded** and **disconnected**. (Note also that for these points it's *not* true that $\varphi_t(x, y) \rightarrow L_\omega(x, y)$ as $t \rightarrow \infty$.)

[Hint: To show that the solution curves spiral outwards, consider in what direction they deviate from the closed solution curves of the conservative system $\dot{x} = -y, \dot{y} = x(1-y^2)$.]

- A9 Show that the following systems have no closed orbits:

(a) $\dot{x} = y + x^3, \dot{y} = x + y + y^3$.

(b) $\dot{x} = 1 - x^3 + y^2, \dot{y} = 2xy$.

(c) $\dot{x} = y, \dot{y} = -x - y + x^2 + y^2$.

[Hint: Use Bendixson–Dulac with $f(x, y) = e^{-2x}$.]

[§] If \mathbf{x}_n is a sequence of points in M converging to \mathbf{x} , then $\mathbf{x} \in \overline{B} = B$, and by continuity

$$\underbrace{V(\mathbf{x}_n)}_{\leq \beta} \rightarrow V(\mathbf{x}),$$

so $V(\mathbf{x}) \leq \beta$. Hence $\mathbf{x} \in M$, which means that M is closed.

Lecture 8. Some applications

(Arrowsmith & Place, sections 5.1, 5.2, 5.3, 5.4.)

- In this lecture we will have a look at a selection of the applications from Chapter 5, but there will not be time to cover everything, so you'll have to read the rest for yourself.
- The analysis in Section 5.3.3 of the **Holling-Tanner predator-prey model**

$$\frac{dx}{dt} = rx \left(1 - \frac{x}{K}\right) - \frac{wxy}{D+x}, \quad \frac{dy}{dt} = sy \left(1 - \frac{y}{J}\right)$$

can be made simpler by writing the system in **dimensionless variables** (s, u, v) instead of (t, x, y) . This **nondimensionalization** is a very useful technique for reducing the number of parameters in a system, and since it's not described in the textbook, I'll explain it here instead. Let

$$s = c_0 t, \quad x = c_1 u, \quad y = c_2 v,$$

where c_0 , c_1 and c_2 are constants that we will specify soon. Inserting this into the differential equations, we get

$$\frac{c_1}{c_0} \frac{du}{ds} = rc_1 u \left(1 - \frac{c_1 u}{K}\right) - \frac{wc_1 uc_2 v}{D + c_1 u}, \quad \frac{c_2}{c_0} \frac{dv}{ds} = sc_2 v \left(1 - \frac{c_2 v}{c_1 u/J}\right),$$

which we can simplify to

$$\frac{du}{ds} = \underset{\uparrow}{(rc_0)} u \left(1 - \frac{c_1}{K} u\right) - \frac{\frac{wc_0 c_2}{c_1} uv}{\frac{D}{c_1} + u}, \quad \frac{dv}{ds} = (sc_0) v \left(1 - \frac{c_2 J}{c_1} \frac{v}{u}\right).$$

At this stage, we have some freedom of choice, but for example we can get rid of the coefficients indicated by the arrows, if we choose c_0 , c_1 and c_2 such that

$$rc_0 = 1, \quad \frac{c_1}{K} = 1, \quad \frac{c_2 J}{c_1} = 1.$$

In other words, we take

$$c_0 = \frac{1}{r}, \quad c_1 = K, \quad c_2 = \frac{K}{J}. \quad (1)$$

Then the equations become

$$\frac{du}{ds} = u(1-u) - \frac{\frac{w}{rJ} uv}{\frac{D}{K} + u}, \quad \frac{dv}{ds} = \frac{s}{r} v \left(1 - \frac{v}{u}\right),$$

and if we now give names to the remaining coefficients appearing in the formulas, for example

$$\alpha = \frac{w}{rJ}, \quad \beta = \frac{s}{r}, \quad \delta = \frac{D}{K}, \quad (2)$$

then the system in its final form is

$$\frac{du}{ds} = u(1-u) - \frac{\alpha uv}{\delta + u}, \quad \frac{dv}{ds} = \beta v \left(1 - \frac{v}{u}\right). \quad (3)$$

Note that this system contains only three parameters (α, β, δ) , instead of the original six parameters (r, K, w, D, s, J) . By using the possibility of rescaling the three variables, we have reduced the number of parameters by three.

Our choice (1) of the constants c_k means that the new variables the we have introduced are actually

$$s = \frac{t}{c_0} = r t, \quad u = \frac{x}{c_1} = \frac{x}{K}, \quad v = \frac{y}{c_2} = \frac{Jy}{K}.$$

Considering that the parameter r in the original ODEs must have the dimension $[\text{time}]^{-1}$ (it's a per capita growth rate), and that t of course has dimension $[\text{time}]$, we see that the rescaled time variable $s = r t$ is actually dimensionless. Similarly, the carrying capacity K for the prey has the same dimension as the prey population size x (whatever unit we happen to use for this, like number of millions of individuals, or biomass in kilograms, or something else), so the variable $u = x/K$ is dimensionless. And so is v , as you can check.

Moreover, the new parameters given by (2) are also dimensionless. For example, r and s are both per capita growth rates and have the same units, so $\beta = s/r$ is a dimensionless quantity which measures the *ratio* between the intrinsic growth rates of the two species. It is rather meaningless to say something like “ r is small”, since this depends on what time unit we are using – if we switched from measuring time in nanoseconds to measuring it in centuries, we would get a very different numerical value for r . But the statement “ β is small” (say much less than 1) expresses a fact which is meaningful regardless of scale, namely that the predators reproduce much slower than the prey.

With the system in the simpler form (3), we can now carry out the same analysis as in Section 5.3.3, but it will be cleaner, since there is less to write, and we also don't get all the original parameters scattered among our formulas, but we always keep them gathered in the meaningful combinations α , β and δ . In the textbook, they do a bit of rescaling at the end, namely taking x/x^* and y/y^* as new variables, where (x^*, y^*) is the nontrivial equilibrium point, but they don't use the possibility of rescaling time to get rid of one more parameter.

Lesson 8

- Damped harmonic oscillator: 5.2, 5.3, 5.10.
- Population models: 5.15, 5.16, 5.17.

The system in 5.15 is the same as in 1.19d that you have done before. Since we view it as population model here, we can restrict ourselves to investigating the region $x_1 \geq 0$, $x_2 \geq 0$.

- Epidemics: 5.21.

(Hint: Find a constant of motion.)

Lecture 9. More about existence and uniqueness

(Not covered in Arrowsmith & Place; see notes below instead.)

Our goal this time is to use **Picard iteration**, also known as the **method of successive approximations**, to prove the **Picard–Lindelöf theorem**, the fundamental existence and uniqueness theorem for a system of (non-autonomous) first order ODEs $\dot{\mathbf{x}} = \mathbf{X}(t, \mathbf{x})$ with a given initial condition $\mathbf{x}(t_0) = \mathbf{c}$.

Preliminaries from analysis: uniform convergence

We will need some theorems about convergence of sequences and series of *functions* (not just *numbers*).

Definition (Pointwise and uniform convergence). Suppose f and f_0, f_1, f_2, \dots are real-valued functions all defined on the same set I (for example an interval).

- The sequence $(f_n)_{n=0}^{\infty}$ converges to the function f **pointwise** on I if

$$\lim_{n \rightarrow \infty} f_n(x) = f(x)$$

is true for each $x \in I$.

[Equivalently: for each $x \in I$ and for each $\varepsilon > 0$ there is an N (which may depend on x and ε) such that $|f_n(x) - f(x)| < \varepsilon$ for all $n \geq N$.]

- The sequence $(f_n)_{n=0}^\infty$ converges to the function f **uniformly** on I if

$$\lim_{n \rightarrow \infty} \sup_{x \in I} |f_n(x) - f(x)| = 0.$$

[Equivalently: for each $\varepsilon > 0$ there is an N (which may depend on ε) such that $|f_n(x) - f(x)| < \varepsilon$ for all $x \in I$ and all $n \geq N$.]

- The function series $\sum_{n=0}^\infty f_n(x)$ converges pointwise/uniformly to the function $s(x)$ if the *sequence of partial sums* $s_n(x) = \sum_{k=0}^n f_k(x)$ converges pointwise/uniformly to $s(x)$.

Remark. The same definitions also apply to complex-valued or vector-valued functions, etc., with the suitable interpretation of what $|f_n(x) - f(x)|$ means.

Theorem. Uniform convergence implies pointwise convergence, but not the other way around.

Proof. If $f_n \rightarrow f$ uniformly, then for a given $\varepsilon > 0$ one can find an N which works for all x , so the same number N will work for each particular x in the definition of pointwise convergence.

An example showing that the converse fails is the sequence $f_n(x) = x^n$ on the interval $[0, 1]$, which converges pointwise, but *not uniformly*, to the discontinuous function

$$f(x) = \begin{cases} 0, & 0 \leq x < 1, \\ 1, & x = 1. \end{cases} \quad \square$$

Theorem (The uniform limit theorem). If each f_n is continuous on I , and $f_n \rightarrow f$ *uniformly*, then f is continuous on I .

Proof. Suppose $a \in I$. Let $\varepsilon > 0$. Since $f_n \rightarrow f$ uniformly, there is an N such that $|f_N(x) - f(x)| < \varepsilon/3$ for all $x \in I$. Since f_N is continuous, there is a $\delta > 0$ such that $|f_N(x) - f_N(a)| < \varepsilon/3$ for all $x \in I$ such that $|x - a| < \delta$. The triangle inequality gives

$$\begin{aligned} |f(x) - f(a)| &= |f(x) - f_N(x) + f_N(x) - f_N(a) + f_N(a) - f(a)| \\ &\leq |f(x) - f_N(x)| + |f_N(x) - f_N(a)| + |f_N(a) - f(a)| \\ &< \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon \end{aligned}$$

for all $x \in I$ such that $|x - a| < \delta$. Thus f is continuous at a . □

Theorem (The Weierstrass M-test). If the numerical series $\sum_{n=0}^\infty M_n$ converges, and if $|f_n(x)| \leq M_n$ for all $x \in I$, then the function series $\sum_{n=0}^\infty f_n(x)$ converges uniformly (and absolutely) on I to some function $S(x)$.

Proof. (Omitted.) □

Remark. In the M-test, if each f_n is *continuous*, then the sum S is also a continuous function. This follows by applying the uniform limit theorem to the sequence of partial sums.

Equivalent integral equation

Lemma. Let $I \subseteq \mathbb{R}$ be an open interval (bounded or unbounded), and assume that $\mathbf{X}: I \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous. Let $t_0 \in I$. Then the function $\mathbf{x}(t)$ ($t \in I$) is a continuously differentiable solution of the initial value problem

$$\begin{aligned} \dot{\mathbf{x}}(t) &= \mathbf{X}(t, \mathbf{x}(t)) \quad \text{for } t \in I, \\ \mathbf{x}(t_0) &= \mathbf{c}, \end{aligned} \quad (\text{A})$$

if and only if it is a continuous solution of the integral equation

$$\mathbf{x}(t) = \mathbf{c} + \int_{t_0}^t \mathbf{X}(s, \mathbf{x}(s)) ds \quad \text{for } t \in I. \quad (\text{B})$$

The same statement holds also for closed intervals I , provided that the derivative $\dot{\mathbf{x}}(t)$ is interpreted as a *one-sided* derivative when t is an *endpoint* of I .

Proof. This is an immediate consequence of the fundamental theorem of calculus. □

Picard iteration

Picard's idea for proving the existence of a solution to problem (A) is to recursively define an infinite sequence of functions

$$\mathbf{x}_0(t), \quad \mathbf{x}_1(t), \quad \mathbf{x}_2(t), \quad \dots \quad (t \in I)$$

by the formulas

$$\begin{aligned} \mathbf{x}_0(t) &= \mathbf{c}, \\ \mathbf{x}_n(t) &= \mathbf{c} + \int_{t_0}^t \mathbf{X}(s, \mathbf{x}_{n-1}(s)) ds \quad \text{for } n \geq 1, \end{aligned}$$

and to show that this sequence converges (under some conditions) to a continuous function $\mathbf{x}(t)$ which satisfies the integral equation (B), and hence also the initial value problem (A). The uniqueness of this solution is proved by separate argument (but under the same conditions).

We may note right away that each function $\mathbf{x}_n(t)$ in the sequence is differentiable on I . This is obvious for $n = 0$ since \mathbf{x}_0 is just a constant function, and for $n \geq 1$ it follows from the fundamental theorem of calculus:

$$\frac{d\mathbf{x}_n}{dt}(t) = \mathbf{X}(t, \mathbf{x}_{n-1}(t)).$$

And since the functions $\mathbf{x}_n(t)$ are differentiable, they are automatically continuous as well.

The Lipschitz condition

How to prove that the sequence defined by Picard iteration converges? Answer: We write

$$\begin{aligned} \mathbf{x}_n &= (\mathbf{x}_n - \mathbf{x}_{n-1}) + \dots + (\mathbf{x}_2 - \mathbf{x}_1) + (\mathbf{x}_1 - \mathbf{x}_0) + \mathbf{x}_0 \\ &= \mathbf{c} + \sum_{k=1}^n (\mathbf{x}_k - \mathbf{x}_{k-1}), \end{aligned}$$

and apply the Weierstrass M -test to show that the function series

$$\mathbf{c} + \sum_{k=1}^{\infty} (\mathbf{x}_k - \mathbf{x}_{k-1})$$

converges (uniformly, on some interval).

For this, we will need to estimate the differences $\mathbf{x}_k - \mathbf{x}_{k-1}$. For $k \geq 2$ we have

$$\begin{aligned} \mathbf{x}_k(t) - \mathbf{x}_{k-1}(t) &= \left(\mathbf{c} + \int_{t_0}^t \mathbf{X}(s, \mathbf{x}_{k-1}(s)) ds \right) \\ &\quad - \left(\mathbf{c} + \int_{t_0}^t \mathbf{X}(s, \mathbf{x}_{k-2}(s)) ds \right) \\ &= \int_{t_0}^t (\mathbf{X}(s, \mathbf{x}_{k-1}(s)) - \mathbf{X}(s, \mathbf{x}_{k-2}(s))) ds. \end{aligned}$$

To get anything interesting out of this expression, it's necessary to assume something about the function \mathbf{X} . The natural assumption in this context is that \mathbf{X} satisfies the following so-called **Lipschitz condition** with respect to \mathbf{x} : there is some set $\Omega \subseteq \mathbf{R}^n$ and some constant $L > 0$ such that*

$$|\mathbf{X}(t, \mathbf{a}) - \mathbf{X}(t, \mathbf{b})| \leq L|\mathbf{a} - \mathbf{b}| \quad \text{for all } t \in I \text{ and for all } \mathbf{a}, \mathbf{b} \in \Omega. \quad (\text{Lip})$$

This assumption allows us to make an estimate where we get rid of the terms containing \mathbf{X} , as follows: if

$$\mathbf{x}_{k-1}(t) \in \Omega \quad \text{and} \quad \mathbf{x}_{k-2}(t) \in \Omega \quad \text{for all } t \in I,$$

then for $t_0 \leq t \in I$ we have

$$\begin{aligned} |\mathbf{x}_k(t) - \mathbf{x}_{k-1}(t)| &= \left| \int_{t_0}^t \left(\mathbf{X}(s, \mathbf{x}_{k-1}(s)) - \mathbf{X}(s, \mathbf{x}_{k-2}(s)) \right) ds \right| \quad (\text{put } |\dots| \text{ around the equality above}) \\ &\leq \int_{t_0}^t \left| \mathbf{X}(s, \mathbf{x}_{k-1}(s)) - \mathbf{X}(s, \mathbf{x}_{k-2}(s)) \right| ds \quad (\text{triangle inequality for integrals}) \\ &\leq L \int_{t_0}^t |\mathbf{x}_{k-1}(s) - \mathbf{x}_{k-2}(s)| ds \quad (\text{because of the Lipschitz condition}), \end{aligned}$$

and similarly for $t_0 \geq t \in I$ with the bounds of integration in the opposite order:

$$|\mathbf{x}_k(t) - \mathbf{x}_{k-1}(t)| \leq L \int_t^{t_0} |\mathbf{x}_{k-1}(s) - \mathbf{x}_{k-2}(s)| ds.$$

In the proofs below, these inequalities will allow us to use knowledge about one difference $\mathbf{x}_{k-1} - \mathbf{x}_{k-2}$ to say something about the next difference $\mathbf{x}_k - \mathbf{x}_{k-1}$.

The Picard–Lindelöf theorem

Theorem (Picard–Lindelöf theorem, **global** version). Let $I \subseteq \mathbf{R}$ be an open interval, and assume that $\mathbf{X}: I \times \mathbf{R}^n \rightarrow \mathbf{R}^n$ is continuous and satisfies the Lipschitz condition (Lip) on the whole space \mathbf{R}^n :

$$|\mathbf{X}(t, \mathbf{a}) - \mathbf{X}(t, \mathbf{b})| \leq L|\mathbf{a} - \mathbf{b}| \quad \text{for all } t \in I \text{ and for all } \mathbf{a}, \mathbf{b} \in \mathbf{R}^n.$$

Then for any $t_0 \in I$ and any $\mathbf{c} \in \mathbf{R}^n$, the initial value problem (A),

$$\begin{aligned} \dot{\mathbf{x}}(t) &= \mathbf{X}(t, \mathbf{x}(t)) \quad \text{for } t \in I, \\ \mathbf{x}(t_0) &= \mathbf{c}, \end{aligned}$$

has exactly one solution $\mathbf{x}(t)$. (Note that the interval I may be bounded or unbounded, and that the solution is defined on the whole interval $t \in I$.)

Theorem (Picard–Lindelöf theorem, **local** version). Let $I \subseteq \mathbf{R}$ be an open interval, and assume that $\mathbf{X}: I \times \Omega \rightarrow \mathbf{R}^n$ is continuous and satisfies the Lipschitz condition (Lip) on some open set $\Omega \subseteq \mathbf{R}^n$:

$$|\mathbf{X}(t, \mathbf{a}) - \mathbf{X}(t, \mathbf{b})| \leq L|\mathbf{a} - \mathbf{b}| \quad \text{for all } t \in I \text{ and for all } \mathbf{a}, \mathbf{b} \in \Omega.$$

*For example, in the one-dimensional case, if the partial derivative $\frac{\partial X}{\partial x}(t, x)$ exists and satisfies the boundedness condition

$$\left| \frac{\partial X}{\partial x}(t, x) \right| \leq L \quad \text{for all } t \in I \text{ and for all } a, b \in \Omega,$$

then the mean value theorem for derivatives implies that

$$|X(t, a) - X(t, b)| = \left| (b - a) \frac{\partial X}{\partial x}(t, \xi) \right| \leq L|b - a| \quad \text{for all } t \in I \text{ and for all } a, b \in \Omega,$$

so that the Lipschitz condition holds. Similarly in higher dimensions. For simplicity, one often uses the stronger assumption that $\mathbf{X}(t, \mathbf{x})$ is of class C^1 ; this gives the Lipschitz condition automatically.

Given any $t_0 \in I$ and any $\mathbf{c} \in \Omega$, take $h > 0$ and $r > 0$ small enough that the interval $J = [t_0 - h, t_0 + h]$ is contained in I and the closed ball $B = \overline{B(\mathbf{c}, r)}$ is contained in Ω , and let

$$C = \max_{(t, \mathbf{x}) \in J \times B} |\mathbf{X}(t, \mathbf{x})|.$$

(This maximum exists by the extreme value theorem.) Then the initial value problem (A),

$$\begin{aligned} \dot{\mathbf{x}}(t) &= \mathbf{X}(t, \mathbf{x}(t)) \quad \text{for } t \in [t_0 - \varepsilon, t_0 + \varepsilon] \text{ where } \varepsilon = \min(h, r/C), \\ \mathbf{x}(t_0) &= \mathbf{c}, \end{aligned}$$

has exactly one solution $\mathbf{x}(t)$. (Note that we cannot in general guarantee that the solution is defined on the whole interval I , only on a subinterval.)

Proof of the global version. Define the sequence $(\mathbf{x}_n(t))_{n=0}^\infty$ for $t \in I$ by Picard iteration as above.

Take any $S \in I$ and $T \in I$ with $S < t_0 < T$. We will show that there is a unique solution defined on the interval $[S, T]$, and since S and T are arbitrary, this implies that there is a unique solution on the whole interval I .

Since the function \mathbf{X} is continuous and the interval $[S, T]$ is closed and bounded, the maximum

$$M = \max_{t \in [S, T]} |\mathbf{X}(t, \mathbf{c})|$$

exists, by the extreme value theorem.

Let $t \in [t_0, T]$ to begin with. Then we have

$$|\mathbf{x}_1(t) - \mathbf{x}_0(t)| = \left| \left(\mathbf{c} + \int_{t_0}^t \mathbf{X}(s, \mathbf{c}) ds \right) - \mathbf{c} \right| \leq \int_{t_0}^t |\mathbf{X}(s, \mathbf{c})| ds \leq M(t - t_0),$$

Now that we have an estimate for the first difference $\mathbf{x}_1 - \mathbf{x}_0$, we can start estimating the other differences $\mathbf{x}_k - \mathbf{x}_{k-1}$ successively, using the inequality that we derived in the section about the Lipschitz condition above. This gives (still for $t \in [t_0, T]$):

$$\begin{aligned} |\mathbf{x}_2(t) - \mathbf{x}_1(t)| &\leq L \int_{t_0}^t |\mathbf{x}_1(s) - \mathbf{x}_0(s)| ds \leq L \int_{t_0}^t M(s - t_0) ds = \frac{LM}{2} (t - t_0)^2, \\ |\mathbf{x}_3(t) - \mathbf{x}_2(t)| &\leq L \int_{t_0}^t |\mathbf{x}_2(s) - \mathbf{x}_1(s)| ds \leq L \int_{t_0}^t \frac{L^2 M}{2} (s - t_0)^2 ds = \frac{L^2 M}{2 \cdot 3} (t - t_0)^3, \\ |\mathbf{x}_4(t) - \mathbf{x}_3(t)| &\leq L \int_{t_0}^t |\mathbf{x}_3(s) - \mathbf{x}_2(s)| ds \leq L \int_{t_0}^t \frac{L^2 M}{2 \cdot 3} (s - t_0)^3 ds = \frac{L^3 M}{2 \cdot 3 \cdot 4} (t - t_0)^4, \end{aligned}$$

and so on, with an obvious pattern emerging. To get a uniform estimate, let $t = T$:

$$|\mathbf{x}_k(t) - \mathbf{x}_{k-1}(t)| \leq \frac{L^{k-1} M}{k!} (T - t_0)^k \quad \text{for all } t \in [t_0, T] \text{ and } k \geq 1.$$

If we instead consider $t \in [S, t_0]$, we find in the same way that

$$|\mathbf{x}_k(t) - \mathbf{x}_{k-1}(t)| \leq \frac{L^{k-1} M}{k!} (t_0 - S)^k \quad \text{for all } t \in [S, t_0] \text{ and } k \geq 1.$$

We can combine these two estimates into a single uniform estimate over the whole interval $[S, T]$, less sharp but still good enough for our purposes:

$$|\mathbf{x}_k(t) - \mathbf{x}_{k-1}(t)| \leq \frac{L^{k-1} M}{k!} (T - S)^k \quad \text{for all } t \in [S, T] \text{ and } k \geq 1.$$

The numerical series

$$\sum_{k=1}^{\infty} \frac{L^{k-1} M}{k!} (T - S)^k = \frac{M}{L} \sum_{k=1}^{\infty} \frac{(L(T - S))^k}{k!} = \frac{M}{L} (e^{L(T-S)} - 1)$$

converges, so the Weierstrass M -test shows the uniform convergence on $[S, T]$ of the function series that we have majorized,

$$\mathbf{c} + \sum_{k=1}^{\infty} (\mathbf{x}_k(t) - \mathbf{x}_{k-1}(t)).$$

The partial sums of this series are just the functions

$$\mathbf{x}_n(t) = \mathbf{c} + \sum_{k=1}^n (\mathbf{x}_k(t) - \mathbf{x}_{k-1}(t)),$$

so what we have shown is that the function sequence $(\mathbf{x}_n)_{n=0}^{\infty}$ converges uniformly on $[S, T]$ to some function \mathbf{x} . And since each function \mathbf{x}_n is continuous, the uniform limit theorem shows that this function \mathbf{x} is continuous.

Moreover, for $t \in [S, T]$,

$$\begin{aligned} \mathbf{x}(t) - \mathbf{c} &= \int_{t_0}^t \mathbf{X}(s, \mathbf{x}(s)) ds \\ &= \mathbf{x}(t) - \mathbf{x}_n(t) + \mathbf{x}_n(t) - \mathbf{c} - \int_{t_0}^t \mathbf{X}(s, \mathbf{x}(s)) ds && \text{(add and subtract } \mathbf{x}_n) \\ &= \mathbf{x}(t) - \mathbf{x}_n(t) + \int_{t_0}^t \mathbf{X}(s, \mathbf{x}_{n-1}(s)) ds - \int_{t_0}^t \mathbf{X}(s, \mathbf{x}(s)) ds && \text{(use definition of } \mathbf{x}_n) \\ &= \mathbf{x}(t) - \mathbf{x}_n(t) + \int_{t_0}^t (\mathbf{X}(s, \mathbf{x}_{n-1}(s)) - \mathbf{X}(s, \mathbf{x}(s))) ds, \end{aligned}$$

so

$$\begin{aligned} \left| \mathbf{x}(t) - \mathbf{c} - \int_{t_0}^t \mathbf{X}(s, \mathbf{x}(s)) ds \right| &\leq |\mathbf{x}(t) - \mathbf{x}_n(t)| + \left| \int_{t_0}^t (\mathbf{X}(s, \mathbf{x}_{n-1}(s)) - \mathbf{X}(s, \mathbf{x}(s))) ds \right| \\ &\leq |\mathbf{x}(t) - \mathbf{x}_n(t)| + L \int_{t_0}^t |\mathbf{x}_{n-1}(s) - \mathbf{x}(s)| ds, \end{aligned}$$

where the right-hand side can be made arbitrarily small by taking n large enough, due to the *uniform* convergence

$$\max_{t \in [S, T]} |\mathbf{x}_n(t) - \mathbf{x}(t)| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Therefore, since the left-hand side is nonnegative and independent of n , it must be zero:

$$\mathbf{x}(t) - \mathbf{c} - \int_{t_0}^t \mathbf{X}(s, \mathbf{x}(s)) ds = 0.$$

In other words, the continuous function $\mathbf{x}(t)$ satisfies the integral equation (B) on the interval $[S, T]$, and thus it also satisfies the equivalent initial value problem (A) on $[S, T]$.

To show uniqueness, assume that the function $\mathbf{y}(t)$ is also a continuous solution of (B) on $[S, T]$. Then the maximum

$$A = \max_{t \in [S, T]} |\mathbf{y}(t) - \mathbf{c}|$$

exists, by the extreme value theorem. For $t \in [t_0, T]$ we first estimate

$$\begin{aligned} |\mathbf{y}(t) - \mathbf{x}_1(t)| &= \left| \left(\mathbf{c} + \int_{t_0}^t \mathbf{X}(s, \mathbf{y}(s)) ds \right) - \left(\mathbf{c} + \int_{t_0}^t \mathbf{X}(s, \mathbf{x}_0(s)) ds \right) \right| && \text{(def. of } \mathbf{y} \text{ and } \mathbf{x}_1) \\ &\leq \int_{t_0}^t |\mathbf{X}(s, \mathbf{y}(s)) - \mathbf{X}(s, \mathbf{c})| ds && \text{(triangle inequality for integrals)} \\ &\leq L \int_{t_0}^t |\mathbf{y}(s) - \mathbf{c}| ds && \text{(Lipschitz condition)} \\ &\leq LA(t - t_0) && \text{(definition of } A), \end{aligned}$$

and then successively (in a manner very similar to what we did earlier)

$$\begin{aligned} |\mathbf{y}(t) - \mathbf{x}_2(t)| &\leq \frac{L^2 A (t - t_0)^2}{2}, \\ |\mathbf{y}(t) - \mathbf{x}_3(t)| &\leq \frac{L^3 A (t - t_0)^3}{2 \cdot 3}, \\ |\mathbf{y}(t) - \mathbf{x}_4(t)| &\leq \frac{L^4 A (t - t_0)^4}{2 \cdot 3 \cdot 4}, \end{aligned}$$

and so on. Together with the similar estimates for $t \in [S, t_0]$ we get

$$|\mathbf{y}(t) - \mathbf{x}_n(t)| \leq \frac{L^n A (T - S)^n}{n!} \quad \text{for all } t \in [S, T] \text{ and } n \geq 0.$$

The right-hand side tends to zero as $n \rightarrow \infty$, so the limit of the left-hand side, $|\mathbf{y}(t) - \mathbf{x}(t)|$, must also be zero. Thus $\mathbf{y}(t) = \mathbf{x}(t)$ for all $t \in [S, T]$, and uniqueness is proved. \square

Idea of proof of the local version. Do more or less the same thing, but also use the restrictions to make sure that the Picard iteration doesn't take us outside of the region where the Lipschitz condition holds. \square

Lesson 9

- Integral equations and Picard iteration: [A10](#).
- Non-uniqueness and non-existence: [A11](#), [A12](#).
- Grönwall's lemma: [A13](#), [A14](#).

Additional problems for lesson 9

A10 (a) Solve the integral equation

$$x(t) = 1 + \int_0^t x(s) ds$$

exactly. For comparison, also compute the sequence of Picard approximations

$$x_n(t) = 1 + \int_0^t x_{n-1}(s) ds$$

starting with the constant function $x_0(t) = 1$.

[Answer: Exact solution $x(t) = e^t$. Picard iterates $x_n(t) = \sum_{k=0}^n \frac{t^k}{k!}$.]

(b) Do the same for

$$x(t) = 3 + \int_0^t 4s x(s) ds.$$

A11 Consider the ODE $t \dot{x} = 2x$. (Notice that the coefficient of \dot{x} equals zero at $t = 0$, so we might expect some "trouble" there; it's a *singular point* of the equation.)

(a) Verify that

$$x(t) = \begin{cases} t^2, & t \geq 0, \\ Ct^2, & t < 0 \end{cases}$$

satisfies the ODE for any constant C . Thus there are infinitely many solutions satisfying the condition $x(1) = 1$.

(b) Show that there are also infinitely many solutions satisfying the condition $x(0) = 0$, but no solutions satisfying $x(0) = b$ with $b \neq 0$.

A12 Find *all* functions $x(t)$, $t \in \mathbf{R}$ which satisfy

$$\dot{x} = 2\sqrt{|x|}, \quad x(0) = 0.$$

A13 There are several variants of **Grönwall's lemma** (or **Grönwall's inequality**), which all boil down to the fact that if a function satisfies a differential or integral *inequality* of a certain form, then it can be no bigger than the solution of the corresponding differential or integral *equation*.

(The inequality limits how fast the function can grow, and to push those limits to the maximum and *grow as fast as possible*, the function should satisfy the inequality *with equality*.)

Here your task is to prove (with guidance) the following version of Grönwall's lemma:

Theorem. Let $a(t)$ and $b(t)$ be continuous functions with $b(t) \geq 0$, and suppose that $u(t)$ satisfies the integral inequality

$$u(t) \leq a(t) + \int_0^t b(s) u(s) ds \quad \text{for } t \geq 0.$$

Then

$$u(t) \leq y(t) \quad \text{for } t \geq 0,$$

where $y(t)$ is the solution of the corresponding integral equation

$$y(t) = a(t) + \int_0^t b(s) y(s) ds,$$

namely

$$y(t) = a(t) + \int_0^t a(s) b(s) e^{B(t)-B(s)} ds, \quad \text{where } B(t) = \int_0^t b(\tau) d\tau.$$

Outline of proof. Follow these steps:

- First rewrite the integral equation for $y(t)$ as an ODE for the integral $I(t) = \int_0^t b(s) y(s) ds = y(t) - a(t)$ appearing on the right-hand side:

$$I'(t) - b(t) I(t) = a(t) b(t), \quad I(0) = 0.$$

- Multiply both sides by the integrating factor $e^{-B(t)}$ and integrate from 0 to t .
- After having solved for $I(t)$ in this way, you also know what $y(t)$ is, namely $y(t) = a(t) + I(t)$. Verify that your expression for $y(t)$ agrees with the formula for $y(t)$ given in the theorem above.
- Next consider the inequality for $u(t)$. Let $J(t) = \int_0^t b(s) u(s) ds$ be the integral appearing on the right-hand side. It satisfies $J'(t) = b(t) u(t)$. By assumption we have $u \leq a + J$ and $b \geq 0$, and therefore $J' = bu \leq b(a + J)$. So we know that J satisfies

$$J'(t) - b(t) J(t) \leq a(t) b(t), \quad J(0) = 0.$$

- Convince yourself that you can now perform exactly the same steps as when you solved the equation for $I(t)$ (multiply by the integrating factor, integrate, etc.), but with J instead of I and “ \leq ” instead of “ $=$ ”. (Here it's important that $t \geq 0$.)
- And since you do the same steps, you will get the same result in the end, except with “ $u(t) \leq$ ” instead of “ $y(t) =$ ”.
- This proves that $u(t) \leq y(t)$ for $t \geq 0$. Done! □

As a bonus question, see if you can prove this variant of Grönwall's lemma in a similar way:

Theorem. Let g be a continuous function (which need not be positive). If $u(t)$ is continuous for $t \geq 0$ and differentiable for $t > 0$, and satisfies the differential inequality

$$\begin{aligned} u'(t) &\leq g(t) u(t) && \text{for } t > 0, \\ u(0) &= c, \end{aligned}$$

then

$$u(t) \leq y(t) \quad \text{for } t \geq 0,$$

where $y(t)$ is the solution of the corresponding differential equation

$$\begin{aligned} y'(t) &= g(t) y(t) && \text{for } t > 0, \\ y(0) &= c, \end{aligned}$$

namely

$$y(t) = c e^{G(t)}, \quad G(t) = \int_0^t g(s) ds.$$

- A14 (a) Compute the solution $x(t)$ of the initial value problem $\dot{x} = x^2$, $x(0) = c > 0$, and note that it blows up after finite time (as $t \nearrow 1/c$).
- (b) In part (a), the right-hand side of the ODE was obviously quadratic in x . In contrast, show that if the right-hand side of the system $\dot{\mathbf{x}} = \mathbf{X}(\mathbf{x})$ is **linearly bounded**, meaning that there are constants $a \geq 0$ and $b \geq 0$ such that

$$|\mathbf{X}(\mathbf{x})| \leq a + b|\mathbf{x}| \quad \text{for all } \mathbf{x} \in \mathbf{R}^n,$$

then the solution $\mathbf{x}(t)$ with initial value $\mathbf{x}(0) = \mathbf{x}_0$ cannot blow up in finite time, so it is defined for all $t \geq 0$ (regardless of \mathbf{x}_0).

(We are assuming that the vector field $\mathbf{X}(\mathbf{x})$ is nice enough for the flow to exist at least locally, say of class C^1 . It can be proved that if $\mathbf{X}(\mathbf{x})$ is defined for all $\mathbf{x} \in \mathbf{R}^n$, then the only way for solutions to cease existing after finite time is that $|\mathbf{x}| \rightarrow \infty$. Let us take this fact for granted here.)

[Hint: Assume, for a contradiction, that some solution $\mathbf{x}(t)$ only exists for $0 \leq t < t_0$. From

$$\mathbf{x}(t) = \mathbf{x}_0 + \int_0^t \mathbf{X}(\mathbf{x}(s)) ds,$$

deduce that

$$|\mathbf{x}(t)| \leq |\mathbf{x}_0| + at + \int_0^t b|\mathbf{x}(s)| ds \quad \text{for } 0 \leq t < t_0,$$

and apply Grönwall's lemma from problem A13 to show that $|\mathbf{x}(t)|$ cannot tend to ∞ as $t \nearrow t_0$.]

Lecture 10. Linear equations with non-constant coefficients

(Not covered in Arrowsmith & Place; see notes below instead.)

Second-order linear ODEs

Many ODE books with a more "classical" flavour allocate plenty of space to the topic of second order (inhomogeneous) linear ODEs,

$$\ddot{x} + p_1(t) \dot{x} + p_0(t) x = f(t),$$

which appear in many applications, and are also of historical importance. Some particular such ODEs have been studied so much that one could easily spend several courses on them alone, like the **Bessel equation** (with parameter $\alpha \in \mathbf{C}$),

$$\ddot{x} + \frac{1}{t} \dot{x} + \frac{t^2 - \alpha^2}{t^2} x = 0.$$

Any second-order ODE can be rewritten as a system of two first-order ODEs, for example by letting $x_1 = x$ and $x_2 = \dot{x}$:

$$\ddot{x} + p_1(t)\dot{x} + p_0(t)x = f(t) \iff \begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ f(t) - p_1(t)x_2 - p_0(t)x_1 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -p_0(t) & -p_1(t) \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} 0 \\ f(t) \end{pmatrix}.$$

Here we will focus more on systems of first-order (inhomogeneous) linear ODEs in general:

$$\dot{\mathbf{x}} = A(t)\mathbf{x} + \mathbf{f}(t),$$

where $\mathbf{x}(t) \in \mathbf{R}^n$ and $A(t)$ is some $n \times n$ matrix. From the general results about systems we can then obtain some of the basic facts about second-order linear ODEs simply by considering the special case when $n = 2$ and $\dot{x}_1 = x_2$.

First-order systems of linear ODEs

Theorem. Assume that $A(t)$ and $\mathbf{f}(t)$ are *continuous* on some time interval I (bounded or not) containing $t = 0$. Then the initial value problem

$$\dot{\mathbf{x}} = A(t)\mathbf{x} + \mathbf{f}(t), \quad \mathbf{x}(0) = \mathbf{x}_0$$

has a unique solution defined on all of I .

Proof. Let $J = [t_1, t_2]$ be an arbitrary compact subinterval of I containing $t = 0$. By the extreme value theorem, each matrix entry $A_{ij}(t)$ is bounded on J , and since there are only finitely many matrix entries, there must be a common constant which bounds all of them:

$$|A_{ij}(t)| \leq C \quad \text{for all } i \text{ and } j, \text{ and all } t \in J.$$

Then the i th entry in the matrix product $A(t)\mathbf{y}$ can be estimated using the triangle inequality and the Cauchy–Schwarz inequality:

$$|(A(t)\mathbf{y})_i| = |A_{i1}(t)y_1 + \dots + A_{in}(t)y_n| \leq C|y_1| + \dots + C|y_n| \leq C\sqrt{n}|\mathbf{y}|.$$

Squaring, summing over i , and taking the square root, we get

$$|A(t)\mathbf{y}| \leq Cn|\mathbf{y}|,$$

for any vector \mathbf{y} .

Using this property we can show that the right-hand side $\mathbf{X}(t, \mathbf{x}) = A(t)\mathbf{x} + \mathbf{f}(t)$ satisfies the global Lipschitz condition with Lipschitz constant $L = Cn$. Indeed,

$$\mathbf{X}(t, \mathbf{a}) - \mathbf{X}(t, \mathbf{b}) = (A(t)\mathbf{a} + \mathbf{f}(t)) - (A(t)\mathbf{b} + \mathbf{f}(t)) = A(t)\mathbf{a} - A(t)\mathbf{b} = A(t)(\mathbf{a} - \mathbf{b})$$

implies that

$$|\mathbf{X}(t, \mathbf{a}) - \mathbf{X}(t, \mathbf{b})| = |A(t)(\mathbf{a} - \mathbf{b})| \leq Cn|\mathbf{a} - \mathbf{b}| \quad \text{for all } t \in J \text{ and for all } \mathbf{a}, \mathbf{b} \in \mathbf{R}^n.$$

By the global Picard–Lindelöf theorem there is therefore a unique solution defined on all of J , and since the subinterval J was arbitrary, we can extend this solution as far as we like inside I . \square

Theorem. The general solution of the inhomogeneous system $\dot{\mathbf{x}} - A(t)\mathbf{x} = \mathbf{f}(t)$ has the form

$$\mathbf{x}(t) = \mathbf{x}_{\text{hom}}(t) + \mathbf{x}_{\text{part}}(t),$$

where \mathbf{x}_{part} is some particular solution, and \mathbf{x}_{hom} is the general solution of the corresponding homogeneous system $\dot{\mathbf{x}} - A(t)\mathbf{x} = \mathbf{0}$.

Proof. This is just a fact about linearity, and not really about differential equations. If $\mathbf{x} = \mathbf{x}_1$ and $\mathbf{x} = \mathbf{x}_2$ are two particular solutions, then by linearity their difference $\mathbf{x} = \mathbf{x}_1 - \mathbf{x}_2$ satisfies the homogeneous system. \square

The homogeneous case

Let's look at the homogeneous system $\dot{\mathbf{x}} = A(t)\mathbf{x}$ first. If $A(t) = A$ is a *constant* matrix, we have seen before that the solution is simply

$$\mathbf{x}(t) = e^{tA}\mathbf{x}(0),$$

but if $A(t)$ is *time-dependent*, we can't in general find an explicit solution formula like that. But we can still say a few things in principle about the structure of the solution. To formulate the theorem, let

$$(\mathbf{e}_1, \dots, \mathbf{e}_n)$$

denote the standard basis for \mathbf{R}^n , i.e., $\mathbf{e}_k = (0, \dots, 0, 1, 0, \dots, 0)^T$ with a 1 in the k th position.

Theorem (Solution space). The homogeneous system $\dot{\mathbf{x}} = A(t)\mathbf{x}$ has an n -dimensional solution space. For example, a basis is given by the functions $\mathbf{g}_1(t), \dots, \mathbf{g}_n(t)$, where $\mathbf{x}(t) = \mathbf{g}_k(t)$ is the (unique) solution of the initial value problem starting at \mathbf{e}_k :

$$\dot{\mathbf{x}} = A(t)\mathbf{x}, \quad \mathbf{x}(0) = \mathbf{e}_k.$$

In terms of the $n \times n$ -matrix

$$\Phi(t) = [\mathbf{g}_1(t), \dots, \mathbf{g}_n(t)],$$

any solution has the form

$$\mathbf{x}(t) = \Phi(t)\mathbf{x}(0). \tag{SOL}$$

Proof. Let $\mathbf{x}(t) = \mathbf{z}(t)$ be any solution of $\dot{\mathbf{x}} = A(t)\mathbf{x}$, and let $\mathbf{c} = \mathbf{z}(0)$. Then the functions $\mathbf{x}(t) = \mathbf{z}(t)$ and

$$\mathbf{x}(t) = c_1\mathbf{g}_1(t) + \dots + c_n\mathbf{g}_n(t)$$

both satisfy the initial value problem

$$\dot{\mathbf{x}} = A(t)\mathbf{x}, \quad \mathbf{x}(0) = \mathbf{c},$$

so by uniqueness they must be the same function:

$$\mathbf{z}(t) = c_1\mathbf{g}_1(t) + \dots + c_n\mathbf{g}_n(t) = \Phi(t)\mathbf{c}.$$

Thus an arbitrary solution \mathbf{z} can be written as a linear combination of the functions \mathbf{g}_k , which shows that they span the solution space, and the formula (SOL) also follows.

To show that the functions \mathbf{g}_k are linearly independent, suppose that the linear combination $\mathbf{x}(t) = \sum c_k\mathbf{g}_k(t)$ is the zero function; then in particular it's zero when $t = 0$:

$$\mathbf{0} = \mathbf{x}(0) = \sum c_k\mathbf{g}_k(0) = \sum c_k\mathbf{e}_k = (c_1, \dots, c_n)^T.$$

In other words, $c_1 = \dots = c_n = 0$. □

Proposition. The columns of an $n \times n$ matrix $\Phi(t)$ are solutions of the linear system $\dot{\mathbf{x}} = A(t)\mathbf{x}$ if and only if the matrix itself is a solution of the matrix-valued linear ODE

$$\dot{\Phi}(t) = A(t)\Phi(t).$$

Proof. This is an immediate consequence of how matrix multiplication works: the k th column in $A\Phi$ equals A times the k th column in Φ . □

Definition. Any time-dependent $n \times n$ matrix whose columns are *linearly independent* solutions of $\dot{\mathbf{x}} = A(t)\mathbf{x}$ is called a **fundamental matrix** for the system.

Remark. The matrix $\Phi(t)$ in the theorem above is thus one particular fundamental matrix, distinguished by the property that $\Phi(0) = I$. Any other fundamental matrix $\Psi(t)$ has the form $\Psi(t) = \Phi(t)M$ where $M = \Psi(0)$ is a nonsingular constant $n \times n$ matrix; indeed, this is just a change of basis in the solution space. In terms of such a Ψ , the general solution of the initial value problem is $\mathbf{x}(t) = \Phi(t)\mathbf{x}(0) = \Psi(t)M^{-1}\mathbf{x}(0)$.

Rephrasing what we said before the theorem: If A is *constant*, then $\Phi(t) = e^{tA}$ is the fundamental matrix satisfying $\Phi(0) = I$, but if A is *time-dependent* we cannot in general compute the fundamental matrix $\Phi(t)$ explicitly; we only know from the Picard–Lindelöf theorem that it exists and is unique. But curiously enough, we can always compute its determinant:

Theorem (Liouville's identity). If $\Phi(t)$ is a solution of the matrix ODE $\dot{\Phi}(t) = A(t)\Phi(t)$, then its determinant*

$$W(t) = \det \Phi(t)$$

satisfies the scalar ODE

$$\dot{W}(t) = \operatorname{tr}(A(t)) W(t),$$

and hence

$$W(t) = \exp\left(\int_0^t \operatorname{tr}(A(s)) ds\right) W(0).$$

Proof. A down-to-earth way of proving this is to just compute. Consider the 3×3 case, for ease of notation, so that

$$W(x) = \det(\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3) = \begin{vmatrix} g_{11} & g_{21} & g_{31} \\ g_{12} & g_{22} & g_{32} \\ g_{13} & g_{23} & g_{33} \end{vmatrix} = g_{11}g_{22}g_{33} + \dots$$

When differentiating this using the product rule for derivatives, each of the $n!$ terms in the determinant will give rise to n terms, and the resulting sum can be rearranged back into a sum of n determinants:

$$\begin{aligned} \dot{W} &= \frac{d}{dt} \det(\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3) \\ &= \left(\frac{dg_{11}}{dt} g_{22} g_{33} + g_{11} \frac{dg_{22}}{dt} g_{33} + g_{11} g_{22} \frac{dg_{33}}{dt} \right) + \dots \\ &= \det\left(\frac{d\mathbf{g}_1}{dt}, \mathbf{g}_2, \mathbf{g}_3\right) + \det\left(\mathbf{g}_1, \frac{d\mathbf{g}_2}{dt}, \mathbf{g}_3\right) + \det\left(\mathbf{g}_1, \mathbf{g}_2, \frac{d\mathbf{g}_3}{dt}\right) \\ &= \det(A\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3) + \det(\mathbf{g}_1, A\mathbf{g}_2, \mathbf{g}_3) + \det(\mathbf{g}_1, \mathbf{g}_2, A\mathbf{g}_3). \end{aligned}$$

If we expand all these determinants, what terms would we get that contain A_{11} ? Answer: the terms appearing in the expression

$$\begin{vmatrix} A_{11}g_{11} & 0 & 0 \\ 0 & g_{22} & g_{32} \\ 0 & g_{23} & g_{33} \end{vmatrix} + \begin{vmatrix} 0 & A_{11}g_{21} & 0 \\ g_{12} & 0 & g_{32} \\ g_{13} & 0 & g_{33} \end{vmatrix} + \begin{vmatrix} 0 & 0 & A_{11}g_{31} \\ g_{12} & g_{22} & 0 \\ g_{13} & g_{23} & 0 \end{vmatrix} = A_{11} \det(\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3).$$

And the terms that contain A_{12} are

$$\begin{vmatrix} A_{12}g_{12} & 0 & 0 \\ 0 & g_{22} & g_{32} \\ 0 & g_{23} & g_{33} \end{vmatrix} + \begin{vmatrix} 0 & A_{12}g_{22} & 0 \\ g_{12} & 0 & g_{32} \\ g_{13} & 0 & g_{33} \end{vmatrix} + \begin{vmatrix} 0 & 0 & A_{12}g_{32} \\ g_{12} & g_{22} & 0 \\ g_{13} & g_{23} & 0 \end{vmatrix} = A_{12} \begin{vmatrix} g_{12} & g_{22} & g_{32} \\ g_{12} & g_{22} & g_{32} \\ g_{13} & g_{23} & g_{33} \end{vmatrix},$$

but this is zero since two rows are equal, so there will be no terms containing A_{12} in the expansion. Similarly, we get contributions $A_{22} \det(\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3)$ and $A_{33} \det(\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3)$, but no terms containing an A_{ij} with $i \neq j$. Thus,

$$\dot{W} = (A_{11} + A_{22} + A_{33}) \det(\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3) = \operatorname{tr}(A) W,$$

as desired. The formula for $W(t)$ is obtained by solving this ODE for W using an integrating factor. \square

Remark. See math.stackexchange.com/questions/10212/ for a higher-level point of view on this.

* Called the **Wronskian determinant**, or simply the **Wronskian**.

Remark. Liouville's identity shows that if the columns of the matrix $\Phi(t)$ are *solutions of some linear system* $\dot{\mathbf{x}} = A(t)\mathbf{x}$, then $W(t) = \det \Phi(t)$ is either **identically zero** or **never zero**. The case $W(t) = 0$ occurs when the columns of Φ are linearly dependent functions, and $W(t) \neq 0$ when they are linearly independent. But beware that if we just look at the determinant of some arbitrary time-dependent matrix $\Phi(t)$, then the question of linear independence of the columns is not this simple! (See problem A18.)

Consider the special case of second-order linear homogeneous ODEs

$$\ddot{x} + p_1(t)\dot{x} + p_0(t)x = 0,$$

which can be written as $\dot{\mathbf{x}} = A\mathbf{x}$ with

$$\mathbf{x}(t) = \begin{pmatrix} x(t) \\ \dot{x}(t) \end{pmatrix}, \quad A(t) = \begin{pmatrix} 0 & 1 \\ -p_0(t) & -p_1(t) \end{pmatrix},$$

as in the beginning of this lecture. Since $\text{tr}(A(t)) = -p_1(t)$ in this case, Liouville's identity takes the following form:

Theorem (Abel's identity). If $x = y(t)$ and $x = z(t)$ are two solutions of $\ddot{x} + p_1\dot{x} + p_0x = 0$, then their Wronskian

$$W(t) = \begin{vmatrix} y(t) & z(t) \\ \dot{y}(t) & \dot{z}(t) \end{vmatrix} = y(t)\dot{z}(t) - \dot{y}(t)z(t)$$

satisfies

$$\dot{W}(t) = -p_1(t)W(t),$$

and hence

$$W(t) = \exp\left(-\int_0^t p_1(s) ds\right) W(0).$$

The inhomogeneous case

Now to the question of finding a **particular solution** $\mathbf{x}_{\text{part}}(t)$ of the *inhomogeneous* system

$$\dot{\mathbf{x}} - A(t)\mathbf{x} = \mathbf{f}(t),$$

supposing that we already know the general solution $\mathbf{x}_{\text{hom}}(t)$ of the *homogeneous* equation. (In other words, supposing that we know a fundamental matrix.) This can be done by a method called **variation of constants**, **variation of parameters** or **Lagrange's method**, as follows:

Theorem (Variation of constants). If $\Phi(t)$ is a fundamental matrix for the homogeneous system, then

$$\mathbf{x}_{\text{part}}(t) = \Phi(t) \int_0^t \Phi(s)^{-1} \mathbf{f}(s) ds$$

is a particular solution of the inhomogeneous system.

Proof. The fundamental matrix satisfies $\dot{\Phi} = A\Phi$ (by definition). Make the change of variables

$$\mathbf{x}(t) = \Phi(t)\mathbf{y}(t).$$

When we substitute this into the system, together with $\dot{\mathbf{x}} = \dot{\Phi}\mathbf{y} + \Phi\dot{\mathbf{y}}$, we obtain

$$\begin{aligned} & \dot{\mathbf{x}}(t) - A(t)\mathbf{x}(t) = \mathbf{f}(t) \\ \Leftrightarrow & \dot{\Phi}(t)\mathbf{y}(t) + \Phi(t)\dot{\mathbf{y}}(t) - A(t)\Phi(t)\mathbf{y}(t) = \mathbf{f}(t) \\ \Leftrightarrow & \underbrace{(\dot{\Phi}(t) - A(t)\Phi(t))}_{=0}\mathbf{y}(t) + \Phi(t)\dot{\mathbf{y}}(t) = \mathbf{f}(t) \\ \Leftrightarrow & \Phi(t)\dot{\mathbf{y}}(t) = \mathbf{f}(t) \\ \Leftrightarrow & \dot{\mathbf{y}}(t) = \Phi(t)^{-1}\mathbf{f}(t). \end{aligned}$$

(We know that $\Phi(t)^{-1}$ exists for every t , since the Wronskian $W(t) = \det \Phi(t)$ is never zero, by Liouville's identity.) Now just change t to s , integrate both sides from (say) 0 to t to find a particular $\mathbf{y}(t)$ which works, and go back to the old variables \mathbf{x} . Done! \square

Remark. There is no need to memorize the solution formula; just repeat the procedure in the proof every time you need it! The reason for the funny name “variation of constants” is that the general solution of the homogeneous system is

$$\mathbf{x}_{\text{hom}}(t) = \Phi(t) \mathbf{c},$$

where $\mathbf{c} = (c_1, \dots, c_n)^T$ is an arbitrary *constant* vector, and the idea here is to try to find a particular solution by “letting the constants c_k vary”, i.e., by replacing them with the *time-dependent* quantities $\mathbf{y}(t) = (y_1(t), \dots, y_n(t))^T$:

$$\mathbf{x}_{\text{part}}(t) = \Phi(t) \mathbf{y}(t).$$

Back to single second-order (or higher-order) ODEs

Now consider again a single linear ODE of order n , with time-dependent coefficients:

$$\frac{d^n x}{dt^n} + p_{n-1}(t) \frac{d^{n-1} x}{dt^{n-1}} + \dots + p_1(t) \frac{dx}{dt} + p_0(t) x = f(t).$$

The general solution of this inhomogeneous equation equals one particular solution plus the general solution of the homogeneous equation

$$\frac{d^n x}{dt^n} + p_{n-1}(t) \frac{d^{n-1} x}{dt^{n-1}} + \dots + p_1(t) \frac{dx}{dt} + p_0(t) x = 0.$$

The first question is how we may find a basis for the n -dimensional solution space of the homogeneous equation. Sometimes we might get lucky and find one solution (by inspired guessing or power series methods or something else). This can then be used for finding *other* solutions, by the following simple trick:

Theorem (Reduction of order). Suppose $x_0(t)$ is a known solution of the homogeneous equation. Then the substitution $x(t) = Y(t) x_0(t)$ leads to a homogeneous equation of order $n - 1$ for $y(t) = \dot{Y}(t)$.

Proof. Substitute

$$\begin{aligned} x &= Y x_0, \\ \dot{x} &= \dot{Y} x_0 + Y \dot{x}_0, \\ \ddot{x} &= \ddot{Y} x_0 + 2\dot{Y} \dot{x}_0 + Y \ddot{x}_0, \\ &\vdots \end{aligned}$$

into the homogeneous ODE for x . Then the coefficient of Y will be $x_0^{(n)} + p_{n-1} x_0^{(n-1)} + \dots + p_1 \dot{x}_0 + p_0 x_0$, which equals zero by assumption. Thus only $\dot{Y}, \ddot{Y}, \dots, Y^{(n)}$ appear in the equation, so if we let $y = \dot{Y}$ we get an equation involving only $y, \dot{y}, \dots, y^{(n-1)}$. \square

Remark. The reduced equation always has the trivial solution $y(t) = 0$, which gives $Y(t) = C$. But this is quite uninteresting, since $x(t) = Y(t) x_0(t) = C x_0(t)$ is then just a constant multiple of the already known solution $x_0(t)$.

Remark. If we manage to find a solution of a *second-order* homogenous linear ODE, then reduction of order gives a *first-order* equation, which means that we can find a nontrivial solution with the help of an integrating factor. We can then integrate this solution $y(t)$ to find a non-constant $Y(t)$, and hence find a second *linearly independent* solution $x(t) = Y(t) x_0(t)$. So in this case $x_0(t)$ and $Y(t) x_0(t)$ will be a basis of the solution space.

For finding a particular solution of the inhomogeneous equation, we have the method of **variation of constants**, as a special case of what we did for systems. Rewrite the ODE as a first-order system by letting $x_1 = x, x_2 = \dot{x}$, etc.:

$$\begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \vdots \\ \dot{x}_{n-1} \\ \dot{x}_n \end{pmatrix} = \begin{pmatrix} x_2 \\ x_3 \\ \vdots \\ x_n \\ f(t) - \sum_{k=1}^n p_{k-1}(t) x_k \end{pmatrix} = \begin{pmatrix} 0 & 1 & & & \\ & 0 & 1 & & \\ & & \ddots & \ddots & \\ & & & 0 & 1 \\ -p_0(t) & -p_1(t) & \cdots & -p_{n-2}(t) & -p_{n-1}(t) \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_{n-1} \\ x_n \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ f(t) \end{pmatrix}$$

Let's look at the case $n = 3$ to simplify notation:

$$\begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \end{pmatrix} = \begin{pmatrix} x_2 \\ x_3 \\ f(t) - p_0(t)x_1 - p_1(t)x_2 - p_2(t)x_3 \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -p_0(t) & -p_1(t) & -p_2(t) \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ f(t) \end{pmatrix}$$

The method requires that we already know the general solution x_{hom} of the homogeneous equation, say

$$x_{\text{hom}}(t) = c_1 g_1(t) + c_2 g_2(t) + c_3 g_3(t)$$

where (g_1, g_2, g_3) is a known basis for the solution space. We are going to seek a particular solution by "letting the constants c_k vary", i.e., replacing c_k by $y_k(t)$:

$$x_{\text{part}}(t) = y_1(t) g_1(t) + y_2(t) g_2(t) + y_3(t) g_3(t).$$

In terms of the first-order system, this means that

$$\Phi(t) = \begin{pmatrix} g_1 & g_2 & g_3 \\ \dot{g}_1 & \dot{g}_2 & \dot{g}_3 \\ \ddot{g}_1 & \ddot{g}_2 & \ddot{g}_3 \end{pmatrix}$$

is a fundamental matrix, and we are making the substitution $\mathbf{x}(t) = \Phi(t) \mathbf{y}(t)$. Simply remembering what we did for systems above, we know that this leads to $\Phi(t) \dot{\mathbf{y}} = \mathbf{f}(t)$:

$$\begin{pmatrix} g_1 & g_2 & g_3 \\ \dot{g}_1 & \dot{g}_2 & \dot{g}_3 \\ \ddot{g}_1 & \ddot{g}_2 & \ddot{g}_3 \end{pmatrix} \begin{pmatrix} \dot{y}_1 \\ \dot{y}_2 \\ \dot{y}_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ f(t) \end{pmatrix}.$$

This system of equations determines $\dot{\mathbf{y}}$, which we can then integrate to find \mathbf{y} , and hence x_{part} .

Remark. Books which deal only with single higher-order ODEs usually present the method of variation of constants in the following way, which in my opinion is rather obscure! Start with

$$x = y_1 g_1 + y_2 g_2 + y_3 g_3.$$

Take the first derivative:

$$\dot{x} = (\dot{y}_1 g_1 + \dot{y}_2 g_2 + \dot{y}_3 g_3) + (y_1 \dot{g}_1 + y_2 \dot{g}_2 + y_3 \dot{g}_3).$$

For some mysterious reason ("just because it works"), we require the first bracket to be zero: $\dot{y}_1 g_1 + \dot{y}_2 g_2 + \dot{y}_3 g_3 = 0$. So only the second bracket remains in $\dot{x} = y_1 \dot{g}_1 + y_2 \dot{g}_2 + y_3 \dot{g}_3$, and taking the derivative of this gives

$$\ddot{x} = (\dot{y}_1 \dot{g}_1 + \dot{y}_2 \dot{g}_2 + \dot{y}_3 \dot{g}_3) + (y_1 \ddot{g}_1 + y_2 \ddot{g}_2 + y_3 \ddot{g}_3).$$

Again we require the first bracket to be zero: $\dot{y}_1 \dot{g}_1 + \dot{y}_2 \dot{g}_2 + \dot{y}_3 \dot{g}_3 = 0$. This leaves $\ddot{x} = y_1 \ddot{g}_1 + y_2 \ddot{g}_2 + y_3 \ddot{g}_3$, so

$$\ddot{x} = (\dot{y}_1 \ddot{g}_1 + \dot{y}_2 \ddot{g}_2 + \dot{y}_3 \ddot{g}_3) + (y_1 \dddot{g}_1 + y_2 \dddot{g}_2 + y_3 \dddot{g}_3)$$

Now using that g_1, g_2 and g_3 satisfy the homogeneous equation, we find after some computation that if we want x to satisfy the inhomogeneous equation, then we must require the first bracket here to satisfy $\dot{y}_1 \ddot{g}_1 + \dot{y}_2 \ddot{g}_2 + \dot{y}_3 \ddot{g}_3 = f$. These three requirements are exactly the matrix equation $\Phi \dot{\mathbf{y}} = \mathbf{f}$ that we found above in a much simpler way, using systems and matrix algebra.

Lesson 10

- Reduction of order: [A15](#), [A16](#) ([A16b](#)).
- Variation of constants: [A17](#) ([A17c](#)).
- Wronskians: [A18](#).

Additional problems for lesson 10

A15 Determine α^2 so that that the Bessel equation

$$t^2 \ddot{x} + t \dot{x} + (t^2 - \alpha^2)x = 0 \quad (t > 0)$$

has a solution $x_0(t) = t^{-1/2} \sin t$, and use reduction of order to find another (linearly independent) solution.

$$[\text{Answer: } \alpha^2 = 1/4, x(t) = t^{-1/2} \cos t.]$$

A16 (a) The usual method with the characteristic polynomial says that $x(t) = Ae^{2t} + Be^{3t}$ is the general solution of $\ddot{x} - 5\dot{x} + 6x = 0$. Give a direct proof that this *really* is the most general solution, by applying reduction of order using the known solution $x_0(t) = e^{2t}$.

(b) Similarly, show that $x(t) = (At + B)e^{-3t}$ is the general solution of $\ddot{x} + 6\dot{x} + 9x = 0$, by applying reduction of order with $x_0(t) = e^{-3t}$.

(If you look back in your calculus book, you will probably find arguments like these in the section explaining the theory behind the characteristic polynomial.)

A17 Find the general solution. (Variation of constants is useful for finding a particular solution, at least in the more difficult cases. But just for practice, use it in the simpler cases as well.)

(a) $\ddot{x} + 2\dot{x} + x = 2 \sin t$ [Answer: $x(t) = (At + B)e^{-t} - \cos t$.]

(b) $\ddot{x} + 9x = \cos 3t$ [Answer: $x(t) = A \cos 3t + B \sin 3t + \frac{1}{6} t \sin 3t$.]

(c) $\ddot{x} + x = \tan t$ [Answer: $x(t) = A \cos t + B \sin t + \frac{1}{2} \cos t \cdot \ln \left| \frac{1 - \sin t}{1 + \sin t} \right|$.]

(d) $\dot{x}_1 = 2x_1 - 5x_2 + 4t, \dot{x}_2 = x_1 - 2x_2 + 1$ [Answer: $x_1(t) = A(\cos t + 2 \sin t) + B(-5 \sin t) + 8t - 1,$
 $x_2(t) = A \sin t + B(\cos t - 2 \sin t) + 4t - 2$.]

(e) $(t^2 - 1)\ddot{x} - 2t\dot{x} + 2x = t^2 - 1$ (To find $x_{\text{hom}}(t)$, try a power series solution.)
[Answer: $x(t) = At + B(1 + t^2) - t^2 + t \ln \left| \frac{1+t}{1-t} \right| + \frac{1}{2}(1 + t^2) \ln |1 - t^2|$.]

(And if you haven't had enough, you can look back at problem A1.)

A18 (a) Show that there is no second-order ODE $\ddot{x} + p_1(t)\dot{x} + p_0(t)x = 0$ whose general solution has the form $x(t) = At + B \cos t$.

[Hint: Consider the Wronskian $W = y\dot{z} - \dot{y}z$ of $y(t) = t$ and $z(t) = \cos t$.]

(b) Find a 2×2 matrix $\Phi(t)$ such that its columns are linearly independent (as functions of t), but $\det \Phi(t)$ is zero at some points. [Hint: Same as for (a).]

(c) Show that things can be even worse than in (b): the functions $y(t) = t^3$ and $z(t) = |t|^3$ ($t \in \mathbf{R}$) are linearly independent, but their Wronskian is *identically* zero.

Lecture 11. Outlook: Poincaré maps, attractors, chaotic systems

(Arrowsmith & Place, selected parts of chapters 4 & 6.)

This material is not strictly a part of the course, but is provided as “entertainment”, and to give a rough idea of some additional topics which are important in the theory of dynamical systems.

Lesson 11

- Leftovers.